

QV W875t


61841020R



NLM 05063383 0

NATIONAL LIBRARY OF MEDICINE





A

TREATISE ON THERAPEUTICS,

COMPRISING

MATERIA MEDICA AND TOXICOLOGY,

WITH

ESPECIAL REFERENCE TO THE APPLICATION OF THE
PHYSIOLOGICAL ACTION OF DRUGS TO
CLINICAL MEDICINE.



BY

H. C. WOOD, JR., M.D.;

PROFESSOR OF BOTANY AND CLINICAL LECTURER ON DISEASES OF THE NERVOUS SYSTEM IN THE
MEDICAL DEPARTMENT OF THE UNIVERSITY OF PENNSYLVANIA; PHYSICIAN TO THE
PHILADELPHIA HOSPITAL; CORRESPONDING MEMBER OF THE NEW YORK
MEDICO-LEGAL SOCIETY, ETC., AND AUTHOR OF "A TREATISE ON
THERMIC FEVER, OR SUNSTROKE," "FRESH-WATER ALGÆ
OF NORTH AMERICA,—SMITHSONIAN CONTRI-
BUTIONS," ETC. ETC.

PHILADELPHIA:

J. B. LIPPINCOTT & CO.

1874.

QV

W875t

1874

Film no. 10966

Item 1

Entered, according to Act of Congress, in the year 1874, by

J. B. LIPPINCOTT & CO.,

In the Office of the Librarian of Congress at Washington.

TO
DR. GEORGE B. WOOD, LL.D.,

THIS BOOK

IS DEDICATED BY THE AUTHOR, HIS NEPHEW,
AS A TOKEN OF RESPECT AND AFFECTION. .

PREFACE.

AT the present time, when the shelves of private and public libraries are groaning beneath their ever-increasing loads, when a thousand presses in every city send forth day and night their printed messages until the earth is filled with them, it seems almost presumptuous for any one to offer new volumes to the world. Indeed, art is so long, life is so short, that every student has the right to demand of an author by what authority he doeth these things, and to challenge every memoir for its *raison d'être*. This being so, it assuredly will not appear egotistical for the author to state that his voluntary task was first suggested by his own wants, and that to its performance he has brought the training, labor, and experience of years spent in the laboratory, the study, the class-room, and the hospital ward.

There are a number of excellent treatises upon materia medica and therapeutics; yet in various attempts at original research, as well as in the ward and the lecture-room of the hospital, I have keenly felt the want of something more. There are many points of view from which a subject can be looked at; there are many paths by which it may be approached; and to me, other points of view, other modes of approach, have been far more enticing than those adopted in our standard treatises.

The old and tried method in therapeutics is that of empiricism, or, if the term sound harsh, of clinical experience. As stated by one of its most ardent supporters, the best possible development of this plan of investigation is to be found in a close and careful analysis of cases before and after the administration of a remedy, and, if the results be favorable, the continued use of the drug in similar cases. It is evident that this is not a new path, but a highway already worn with the eager but weary feet of the profession for two thousand years.

That very much has been thus accomplished it were folly to deny. Leaving out of sight the growth of the last two decades, almost all of the current therapeutic knowledge has been gained in this way.

Therapeutics developed in this manner cannot, however, rest upon a secure foundation. What to-day is believed is to-morrow to be cast aside, certainly has been the law of advancement, and seemingly must continue to be so. What has clinical therapeutics established permanently and indisputably? Scarcely anything beyond the primary facts that quinia will arrest an intermittent, that salts will purge, and that opium will quiet pain and lull to sleep.

To established therapeutic facts the profession clings as with the heart

and hand of one man,—clings with a desperation and unanimity whose intensity is the measure of the unsatisfied desire for something fixed. Yet with what a Babel of discordant voices does it celebrate its two thousand years of experience!

This is so well known that it seems superfluous to cite examples of the therapeutic discord; and one only shall be mentioned, namely, rheumatism. In this disease, bleeding, nitrate of potash, quinine, mercurials, flying blisters, purgation, opium, the bromides, veratria, and a host of other remedies, all have their advocates clamorous for a hearing; and above all the tumult are to be heard the trumpet-tones of a Chambers, "Wrap your patients in blankets and let them alone."

Experience is said to be the mother of wisdom. Verily she has been in medicine rather a blind leader of the blind; and the history of medical progress is a history of men groping in the darkness, finding seeming gems of truth one after another, only in a few minutes to cast each back to the vast heap of forgotten baubles that in their day had also been mistaken for verities. In the past, there is scarcely a conceivable absurdity that men have not tested by experience and for a time found to be the thing desired; in the present, homœopathy and other similar delusions are eagerly embraced and honestly believed in by men who rest their faith upon experience.

Narrowing our gaze to the regular profession and to a few decades, what do we see? Experience teaching that not to bleed a man suffering from pneumonia is to consign him to an unopened grave, and experience teaching that to bleed a man suffering from pneumonia is to consign him to a grave never opened by nature. Looking at the revolutions and contradictions of the past,—listening to the therapeutic Babel of the present,—is it a wonder that men should take refuge in nihilism, and, like the lotos-eaters, dream that all alike is folly,—that rest and quiet and calm are the only human fruition?

Since the profession has toiled so long and found so little, if further progress is to be made we must question the old methods and search out new ones, which haply may lead to more fruitful fields. In the ordinary affairs and business of life, when anything is to be accomplished, the effort always is to discover what is to be done, and then what are the means at command. A primary knowledge of the end to be accomplished, and a secondary acquaintance with the instruments, are a necessity for successful human effort; and until the sway of this law is acknowledged by physicians, medicine can never rise from the position of an empirical art to the dignity of applied science. Until within a comparatively recent period, it has been impossible to comply with this law. But, through the advances made by the pathologists and by the students of the natural history of disease, we are fast learning the methods in which nature brings the body back to health. When this is done,—when disease is thoroughly understood,—we shall have wrought out the first element of the problem, shall have complied with the first requirement of the law.

It is scarcely within the province of the therapist, and certainly is not possible within the scope and limits of this work, to discuss at length the natural history of disease; but it is allowable to point out evident indications for relief; and this I have done to a greater or less extent throughout the book.

The work of the therapist is chiefly with the second portion of the law. Evidently, it is his especial province to find out what are the means at command, what the individual drugs in use do when put into a human system. It is seemingly self-evident that the physiological action of a remedy can never be made out by a study of its use in disease. Under all circumstances, the problem is one of the most complex with which the human mind has to grapple; and to introduce into this problem the new and ever-varying factors of the effect of disease and its natural vibrations on the system is to put the matter beyond human prescience.

In spite, then, of Dr. Niemeyer's assertion, that experiments made with medicaments upon the lower animals or upon healthy human beings have, as yet, been of no direct service to our means of treating disease, and that a continuation of such experiments gives no prospect of such service, it is certain that in these experiments is the only rational scientific groundwork for the treatment of disease. We must discover what influence a drug exerts when put into the body of a patient before we can use it rationally; and we can gain this coveted knowledge only in the method indicated.

It has been strenuously objected, especially to experiments upon animals, that drugs do not act upon the lower creatures in the same manner as they do upon man. When I first commenced the studies whose outcome is the present volume, I was profoundly impressed with the truth of this oft-repeated assertion and with the difficulties which it put in the way. To-day I do not believe that, stated in its broad sense, it is true. Indeed, more strongly, I assert that it is not true; that, in the vast majority of cases, the actions of drugs upon man and upon the lower animals are, though seemingly different, in reality similar; that the more knowledge we acquire, the fewer exceptions remain unexplained; and that the whole matter is in all probability subject to laws whose development will greatly aid in our explanation of various obscure clinical phenomena.

The general proofs of these assertions are sufficiently obvious, I think, in the following pages to render it unnecessary for me to dwell upon them at length here: moreover, if they be not so obvious to others as to myself, space is here wanting for a full discussion of the subject. I can only make a few general remarks, and point out some of what I believe to be the governing laws.

In the first place, degree and quality are distinct things, and should not be confounded. Yet they frequently are; and because it requires as much morphia to kill a pigeon of a pound weight as to destroy a man, we are told that medicines act differently upon man and the lower animals. Evidently the

conclusion is a non-sequitur, and difference of susceptibility is no proof of difference in the mode of impression. A teaspoonful of Epsom salts may purge one man, whilst it may require ounces to affect another. Evidently there is a difference of susceptibility; but when the impression is once made it is of the same character in each case. As with man and man, so with man and the pigeon,—susceptibility is no measure or gauge of the character of the impression.

A large number of drugs—indeed, it may be said, the larger number of important drugs—exert in the system antagonistic actions. Thus, atropia stimulates the spinal cord, but destroys the conducting power of the nerve-trunks. It is evident that as one or the other of these influences predominates, will there be convulsions or paralysis. Now, if for any reason one animal be exceedingly sensitive to the spinal action of atropia, that animal will in belladonna-poisoning suffer from convulsions, whilst its fellow, which is affected chiefly by the nerve-action of the drug, will, under like circumstances, have paralysis. Here the mere clinician, with his superficial knowledge, seeing the paralyzed and the convulsed lying side by side, says, What a hopeless muddle! Poor fools, these vivisectors! they will never come to any good! In truth, the differences in symptoms in these and in many other cases simply depend upon differences in susceptibility; and the only lesson that the circumstance teaches is the importance of discovering the laws which govern these susceptibilities.

A law which governs the susceptibility to the action of drugs is, that the more highly specialized any system is, the more readily affected is it by a medicine. Thus, the cerebrum of a man is far more highly organized than that of any other animal, and consequently he is far more sensitive to the action of drugs which affect the cerebrum than are the lower forms. Again, in the frog the spinal system is especially developed,—probably, in proportion to the cerebrum, more so than in any other of the animals commonly experimented with: consequently the batrachian is excessively sensitive to remedies which, like strychnia, affect the spinal cord. In obedience to this law, we have resulting the action of opium,—an action which has been considered the strongest proof of the hopelessness of any attempt to explain the effects of drugs upon a man by experiments upon the lower animals. In man, opium causes deep stupor and general relaxation; in the frog, it causes tetanic convulsions. The explanation of these seeming inconsistencies is, however, very evident when the whole subject is looked at. Opium in all animals has a double action, one upon the cerebrum and one upon the spinal centres. In the frog, the latter being the more highly organized, the spinal action overcomes the cerebral. In man, the cerebrum being the more sensitive, stupor replaces the convulsions; yet in man convulsions sometimes occur in opium-poisoning, and in the frog the dose can be so managed as to cause stupor.

A second law which seems to hold sway over the action of drugs upon different animals is that great differences of function in a system affect its

relation to drugs: thus, in a herbivorous animal the alimentary canal is very different from what it is in the carnivora, whose digestive organs in turn differ from those of man,—the omnivore. Medicines which act upon the alimentary canal are apt to vary in their effects upon different orders of animals.

Converse to the above law is that which renders systems which are little specialized similarly acted upon by drugs in different classes of animals.

Thus, the general structure and the functions of the circulatory system are very uniform among vertebrates, as is also the action of those drugs which affect chiefly the circulation: thus, aconite, or digitalis, or potash, influences in the one way the heart of the frog, of the rabbit, and of man.

There are a very few apparent exceptions to the uniformity of the action of drugs upon all animals, which seemingly contravene the laws that have been mentioned. These exceptions are so few, however, that without doubt advancing knowledge will by-and-by explain them all and show what are the laws which for the time being hold in abeyance or overcome those already stated.

An asserted fact, which has recently been brought forward as revealing the worthlessness of animal experimentation, is that some monkeys are not susceptible to the action of strychnia, whilst others are. Granting the truth of the asserted fact, it certainly is explainable. It is at least conceivable that a given species of animal may, by the gradually-acquired habit of feeding upon a substance containing a narcotic poison, acquire an insusceptibility to the influence of that poison, which shall, as it were, belong to its specific type, or, in other words, be an acquired specific character. The nervous system of the opium-eater becomes accustomed to the stimulant, and it is not impossible that a measure of the habit should be transmitted. If the Darwinian law of the gradual evolution by the survival of the fittest have any force, these curious apparent freaks of medicines in regard to their physiological action may be the result of this law, especially since it is species which are affected. It is not all monkeys that are proof against strychnia, but, as we are distinctly told, only one species of monkey; and, so far as I know, it is not all deer that are said to thrive when fed upon tobacco, but only the Virginia deer. Whether this conception be or be not a mere fancy, this much is to my mind very clear, that the few scattered exceptions ought not to outweigh the immense mass of evidence upon the other side, and that it is inconceivable that drugs, in their relations to animal organisms, differ from all other created things in not being subject to law.

In the early portion of this preface I stated that the work had grown out of a need felt by myself: that need was for a book into which should be gathered the many scattered facts in regard to the physiological action of medicine,—a book in which an attempt should be made to sift the true from the false, to reconcile seeming differences, to point out what we know and what we do not know, and to give a platform from which investigators might start forward without the necessity of being, as is so often the case, ignorant of what was already achieved, or of spending a great deal of time in a wild

hunt through the almost boundless, but often scattered and inaccessible, ranges of Continental literature.

The plan of the present work has been to make the physiological action of remedies the principal point in discussion. A thoroughly scientific treatise would in each article simply show what the drug does when put into a healthy man, and afterwards point out to what diseases or morbid processes such action is able to afford relief. Unfortunately, in the great majority of cases our knowledge is not complete enough for this, and the clinical method has to be used to supplement the scientific plan.

I have added to the book a consideration of toxicology, so far as it is of interest to the physician. This has been done for several reasons. First, it was necessary to study the action of poisonous drugs upon man, in order to make out their physiological action; secondly, physicians are constantly required to diagnose and to treat cases of poisoning; thirdly, it is often of the greatest importance for a medical man in a court of law to be able to state what are the symptoms and post-mortem appearances produced by a given poison, what diseases they simulate, and how far and in what they differ from the phenomena of these diseases. That part of the science of toxicology which treats of the recognition of poisons in the cadaver, or in food and drink, belongs to the domain of the chemist, and I have avoided it altogether. For a similar reason, in the sections on *materia medica*, the chemical relations of mineral substances have not been discussed at all.

Imperfect as is our knowledge, in reviewing the literature I have been surprised to find how much work has been done; how many facts there are, hitherto, not garnered into systematic treatises, at least in the English tongue. Another fact which has been very apparent is the great extent of medical literature, and the still greater difficulty of reviewing the whole of it satisfactorily. Much that is most valuable is most scattered and difficult of access; especially is this true of the work of our German confrères. In the city of Philadelphia, medical literature, at least so far as concerns the matter in hand, is better represented than anywhere else in America; and yet, when thwarted and worried by my inability to procure some desired book, how often have I coveted the complete libraries of the Old World! So serious has been this want of books and of original memoirs that I have generally avoided all questions as to the priority of discovery in regard to facts. In this connection, I desire to thank most heartily Prof. A. Stillé for the magnanimous liberality with which he has placed his really magnificent private library at my disposal,—a liberality without which the book could not have been written.

To my former pupil, Dr. J. Wm. White, I am indebted for aid in preparing the manuscript for press, and for many suggestions. I feel also that it is but just to express my thanks to Mr. J. McCreery, to whose intelligent care and great watchfulness whatever freedom from typographical errors the book may possess is chiefly due, and to whom I am under great obligations for many suggestions as to matters of style, and for saving me much labor.

TABLE OF CONTENTS.

INTRODUCTION.—Pharmacy — Therapeutics — Materia Medica—Pharmacopœia — Preparations—Effects of Medicines—Classification.

PART I.—SYSTEMIC REMEDIES.

DIVISION I.—GENERAL REMEDIES.

ASTRINGENTS.—*Vegetable Astringents*.—Tannic Acid—Gallic Acid—Gall—Catechu—Kino—Hæmatoxylon—Rhatany—Oak Barks—Roses—Geranium—Blackberry Roots—Tormentil—Persimmon—Pomegranate.

Mineral Astringents.—Alums—Sulphate of Aluminium—Lead and its Preparations—Bismuth and its Preparations—Zinc and its Preparations—Copper and its Preparations—Silver and its Preparations.

TONICS.—*Simple Bitters*.—Quassia—Simaruba—Gentian—Barberry—Columbo—Boneset—Goldthread—American Centaury—Dogwood—Salicin.

Peculiar Bitters.—Wild Cherry Bark—Cinchona—Sulphate of Quinine—Sulphate of Cinchonina—Chinoidine.

Aromatic Bitters.—Chamomile—Virginia Snakeroot—Cascarilla—Angustura.

True Aromatics.—Cinnamon—Cloves—Nutmeg—Allspice—Cardamom—Ginger—Black Pepper—Red Pepper—Oil of Cajeput, etc.

Mineral Tonics.—Iron and its Preparations—Sulphuric Acid—Muriatic Acid—Nitric Acid—Nitromuriatic Acid—Phosphorus.

CARDIAC STIMULANTS.—Ammonia, Carbonate and Nitrate—Alcohol—Brandy—Whisky—Wine—Turpentine—Digitalis.

CARDIAC SEDATIVES.—Antimony and its Preparations—Veratrum Viride—Veratrum Album—Veratria—Aconite Leaves and Root—Prussic Acid—Cyanide of Potassium—Vegetable Acids.

ANTISPASMODICS.—Musk—Castor—Valerian—Valerianic Acid—Assafetida—Camphor—Brominated Camphor—Amber—Oil of Amber—Hoffman's Anodyne—Hops—Black Snakeroot.

ANALGESICS.—Opium—Morphia—Narcenin—Codeia—Narcotina—Thebaia—Papaverina—Porphyroxin—Meconia—Meconic Acid—Indian Hemp—American Hemp.

MYDRIATICS.—Belladonna—Atropia—Hyoscyamus—Stramonium.

ANÆSTHETICS.—Nitrous Oxide—Ether—Chloroform—Bichloride of Methyl.

EXCITO-MOTORS.—Strychnia and Drugs containing it.

DEPRESSO-MOTORS.—Calabar Bean—Bromide of Potassium—Bromide of Sodium—Bromide of Ammonium—Chloral—Nitrite of Amyl—Lobelia—Conium.

ALTERATIVES.—Arsenic and its Preparations—Mercury and its Preparations—Iodine and its Preparations—Iodoform—Cod-Liver Oil—Phosphoric Acid and the Phosphates—Colchicum—Sarsaparilla—Guaiac—Mezereum—Sassafras—Dandelion.

DIVISION II.—LOCAL REMEDIES.

EMETICS.—*Vegetable Emetics*.—Ipecacuanha—Bloodroot—Apomorphia—Gillenia—Mustard—Squill.

Mineral Emetics.—Antimony—Sulphate of Zinc—Sulphate of Copper—Alum.

- CATHARTICS.—*Laxatives*.—Foods—Tamarinds—Cassia Fistula—Magnesia—Sulphur.
Purges.—Castor Oil—Mercury—Rhubarb—Butternut—Aloes—Senna—Saline Purgatives.
Drastics.—Scammony—Colocynth—Compound Cathartic Pills—Podophyllum—Ela-
 terium—Gamboge—Croton Oil—Black Hellebore.
- DIURETICS.—*Hydragogue Diuretics*.—Squill—Digitalis—Broom—Sweet Spirit of Nitre.
Refrigerant Diuretics.—Potash and its Preparations.
Stimulating Diuretics.—Buchu—Pareira—Uva Ursi—Pipsissewa—Juniper—Carota
 —Erigeron—Turpentine—Copaiba—Cubeb—Matico—Cantharides.
- DIAPHORETICS.—*Nauseating Diaphoretics*.—*Refrigerant Diaphoretics*.—*Simple Diaphoretics*.
 —Spirit of Mindererus—Sweet Spirit of Nitre—Alcohol.
- EXPECTORANTS.—Atomization.
Nauseating Expectorants.—Lobelia—Ipecacuanha—Tartar Emetic.
Stimulating Expectorants.—Muriate of Ammonia—Seneka—Ammoniac—Benzoin—
 Benzoic Acid—Balsams of Tolu and Peru—Garlic—Squill—Tar.
- EMMENAGOGUES.—*Tonic Emmenagogues*.—*Purgative Emmenagogues*.—*Stimulating Emmena-
 gogues*.—Savine—Rue—Parsley—Apiol—Cantharides—Guaiac.
- OXYTICS.—Ergot.
- SIALAGOGUES.—Pellitory.
- ERRHINES.
- EPISTASTICS.—Counter-irritation—Cantharides.
- RUBEFACTANTS.—Mustard—Spices—Turpentine—Burgundy and Canada Pitch.
- ESCHAROTICS.—Potash—Vienna Paste—Arsenic—Chloride of Zinc—Canquoin's Paste, etc.
- DEMULCENTS.—Gum Arabic—Tragacanth—Slippery Elm—Iceland Moss—Irish Moss—
 Liquorice-Root—Sassafras—Tapioca—Arrowroot—Sago—Barley.
- EMOLLIENTS.—Glycerine.
- DILUENTS.
- PROTECTIVES.—Collodium—Solution of Gutta-Percha.

PART II.—NON-SYSTEMIC REMEDIES.

- ANTACIDS.—Sodium and its Preparations—Lime.
- ANTHELMINTICS.—Pinkroot—Wormseed—Kosso—Santonin—Male Fern—Pumpkin-Seed.
 etc.
- DIGESTANTS.—Pepsin.
- ABSORBENTS.—Charcoal.
- DISINFECTANTS.—*Oxidizing Disinfectants*.—Permanganate of Potassium—Chlorine and its
 Preparations—Iodine—Bromine.
Desulphurating Disinfectants.—Metallic Salts—Lime.
Antizymotics.—Sulphurous Acid—Carbolic Acid.
Absorbing Disinfectants.
- APPENDIX.—Tables—Preparations omitted by an oversight from the text—Errata.

A TREATISE
ON
THERAPEUTICS,
INCLUDING
MATERIA MEDICA AND TOXICOLOGY.

INTRODUCTION.

ALTHOUGH PHARMACY, or the science of preparing medicines, is entirely distinct from THERAPEUTICS, or the science of the application of medicines to the cure of disease, it is evident that some acquaintance with the former is necessary to the correct appreciation of the latter. Further, as the basis of both these studies, must first come a knowledge of MATERIA MEDICA, or the substances used as medicines. PHARMACOLOGY is the general term employed to embrace these three divisions.

In every civilized country there is some recognized official list of standard drugs and their preparations, known as the *Pharmacopœia*. In most places, this, being prepared with the sanction of the government, partakes of the nature of a law; but with us conformity to it depends upon the voluntary action of the professions of Medicine and Pharmacy, by a representative convention of which it was originally prepared and is decennially revised.

The *United States Pharmacopœia* is divided into three portions: a *primary* and a *secondary materia medica list*, and a chapter on *preparations*, with directions for their manufacture. The primary list contains medicines whose reputation is believed to be assured; the secondary list, those still on trial, or those which experience has shown to be not altogether valueless, but yet of little importance.

It is evident that a knowledge of the officinal or recognized preparations, of their general mode of manufacture, and of their strength, is essential to the therapist. The *United States Pharmacopœia* recognizes twenty-nine classes of them, as follows:

DECOCTA.—*Decoctions* are made by boiling crude drugs for a greater or less time in water. It is evident that this method of preparing is ineligible

when the active principle is volatile or is easily decomposed by heat, or when the drug contains much starch, whose extraction would make the preparation very thick and predispose it to rapid decomposition. The method is especially adapted to woody, hard substances, and to those containing much albumen, which is coagulated by the boiling water and left in the drug.

INFUSA.—*Infusions* are made with water, either cold or hot, without boiling. They are prepared by maceration, or by displacement.

LIQUORES.—*Solutions* are preparations in which an active, *non-volatile* principle is dissolved in water.

AQUÆ.—*Waters* are solutions of *volatile* principles in water.

MISTURÆ.—*Mixtures* are preparations in which one or more medicinal substances are held in suspension in water. Of such nature are emulsions, in which some oily material is suspended by a gummy or an albuminous body.

MUCILAGINES.—*Mucilages* are solutions of gummy substances in water.

SYRUPÆ.—*Syrups* are sugary liquids, the menstruum or basis of which is water, with, in some cases, vinegar or alcohol.

MELLITA.—*Honeys* are preparations whose basis is honey.

ACETA.—*Vinegars* are those preparations in which vinegar, or dilute acetic acid, is used as the menstruum.

TINCTURÆ.—*Tinctures* are alcoholic solutions prepared from the crude drug, or by dissolving *non-volatile* principles. In some of them strong, in others dilute, alcohol is used. They are made by maceration or displacement.

SPIRITUS.—*Spirits* are alcoholic solutions of volatile principles, made by direct solution or by distillation from the crude drugs.

VINA.—*Wines* are preparations whose menstruum is wine.

GLYCERITA.—*Glycerites* are preparations in which glycerine is the solvent.

OLEA DESTILLATA.—*Volatile Oils* are active principles of such nature, prepared by distillation.

OLEORESINÆ.—*Oleoresins* are concentrated preparations, composed generally of a volatile oil and a resin. They are really ethereal extracts, made by the action of ether upon the crude drugs; in the case of ginger, a mixture of alcohol and ether is used.

SUCCI.—*Fresh juices* are obtained by expression of the green plant, enough alcohol being added to preserve them.

EXTRACTA.—*Solid extracts* are of two kinds; one being prepared by the evaporation of the fresh juice, the other being made in various ways from the crude drugs.

EXTRACTA FLUIDA.—*Fluid extracts* are very concentrated fluid preparations, generally so made that one minim represents one grain of the crude drug.

RESINÆ.—*Resins* are purgative resinous principles, obtained by the precipitation of saturated tinctures with water.

CONFECTIONES.—*Confections* are medicinal substances beaten up with sugar into a pasty mass.

TROCHISCI.—*Troches*, or *lozenges*, are gummy pellets or disks, so made as to dissolve slowly in the mouth.

SUPPOSITORIA.—*Suppositories* are conical bodies, prepared for introduction into the rectum, where they melt with the heat of the body. Their basis is generally cacao butter.

UNGUENTA and CERATA.—*Ointments* and *Cerates* are fatty solid preparations for external use. The cerates containing wax (*cera*) are the firmer of the two.

EMPLASTRA.—*Plasters* are solid substances spread by the aid of heat upon muslin, skin, or other similar material, and of such nature as to be adhesive at the temperature of the body.

CHARTÆ.—*Papers* are medicated leaves or sheets of paper for external use. The only official are those of mustard and of cantharides.

LINIMENTA.—*Liniments* are liquid preparations, generally soapy or oily, and always intended to be applied externally by rubbing.

The names of **PILULÆ** (*Pills*) and **PULVERES** (*Powders*) sufficiently indicate the character of the preparations.

In what way medicines produce changes in the life-actions of various parts is, and probably must ever remain, unknown, precisely as it is beyond the limit of the human intellect to know why the nerve-cell or the spermatozoon performs the prodigies of which it is capable. The fact that various substances do possess a selective tendency, so to speak, do act more forcibly on one form of tissue than on another, is, however, indisputable.

The effects of medicine are commonly divided into the *primary* and *secondary*, or the *immediate* and the *remote*. An example will probably show the difference between these in the briefest and most forcible manner. Thus, the immediate effect of a diuretic is increased urination. The secondary effect may be removal of serous effusion in some part of the body. It is evident that the latter is brought about not by the medicine itself, but by the changes it induces. Thus, increased excretion causes lessening of the amount of fluid in the blood-vessels, and this in turn predisposes to absorption.

The term or expression *indication* for a given remedy, being in constant use, ought to be distinctly understood; by it is meant the pointings of nature, or, in other words, the evident needs of the system. Thus, hard fæces collected in the colon are an indication for a purgative of such character as will produce watery discharges to soften them. Relaxation in a part indicates a remedy of such properties that it will awake into new life the natural contractility of the part. Again, the suppression of secretion from over-excitement, or from irritation, is an indication for some drug which will allay irritation; whilst the same suppression, when dependent upon torpor or loss of cell-activity, will call for an excitant,—an irritant. The childish absurdity of treating symptoms by any such law as “*similia similibus curantur*,” or “*dissimilia dissimilibus curantur*,” is at once apparent. Symptoms are merely

the surface-play of disease; and the rational therapist always seeks out their hidden meaning.

By far the greater number of remedies are absorbed into the blood, and thus find access to the part upon which they act. It is necessary, therefore, that they be so placed that they can be taken into the blood-vessels.

There are five paths of entrance for medicines into the circulation,—the stomach, the cellular tissue, the rectum, the skin, the lungs. By far the most frequently employed of these is the first-named. It is evident that, in order to pass readily and rapidly into the absorbents, medicines must be in solution. When administered by the stomach, however, it is equally plain that solubility in an ordinary menstruum, such as water, is not a *sine qua non*, since the varying acidities, alkalinities, and organic contents of the alimentary juices give them a solvent power far above that of less complex and varying fluids. Thus, a medicine insoluble in water may be dissolved by the acids of the gastric juice, while another drug may owe its activity to its solution by the alkalies or by the fatty matters of the intestinal fluids.

The dissolving power of the rectal fluids is very slight: hence, in order to act efficiently when given by the rectum, medicines must be in solution or readily soluble. Absorption, moreover, does not occur so rapidly from the rectum as from the stomach, and hence greater time is needed to impress the system in this way. In the great majority of cases medicines are thus exhibited to obtain peculiar effects more or less local in character. Thus, an opium suppository is given in dysentery, or to quiet irritation of the genito-urinary organs, or to check vomiting.

The subcutaneous tissue affords the most rapid route to the central organs. The substance must, however, always be in *perfect* solution, and not too irritating. It is thrown under the skin by means of a small syringe with a sharp hollow needle for its nozzle. Great care should be taken to avoid throwing a medicine into a vein, and so producing a sudden overwhelming effect. The point of the syringe-needle should not be thrust deeply into the tissues, but be kept just beneath the skin, and be withdrawn a little, after having been pushed along under the skin farther than necessary. The objections to the method are the danger of injecting into a vein, the slight pain of the operation, and the skill required for the latter, so that the administration cannot be left to the nurse, together with the local irritation and, it may be, inflammation, to which many drugs so used give rise. For these reasons the *hypodermic* method is not employed except under certain circumstances. When, as in severe suffering, an immediate decided action is required, it is beyond all value,—the system coming under the influence of the remedy as if by magic. The dose should always be smaller than that by the mouth.

There are several ways in which medicinal principles are introduced through the skin, although the only one in common use is the application of medicated fatty preparations, either with or without friction. Absorption takes place, of course, most rapidly in those positions at which the skin is thinnest,—

the inside of the thighs, the surface of the abdomen, and especially the arm-pits. Almost the only remedy which is introduced into the system in this way is mercury. Absorption will take place through the skin from baths, but so slowly that advantage is never taken of it in the constitutional treatment of disease.—unless the sulphur baths, sometimes employed in rheumatism, impress the general system by absorption, which seems to me doubtful.

Formerly, medicines were sometimes exhibited by placing them on blistered surfaces, *beneath* the raised cuticle; but, except in the instance of morphia, so used in gastric disturbance, at present the *endermic* method is very rarely employed.

To influence the general system, the lungs are only used for the introduction of the vapors of the anæsthetics.

For *local* purposes medicines are applied to various parts,—to the skin, to the ear, nares, fauces, stomach, larynx, lungs, rectum, vagina, urethra, etc. For the last three, liquid preparations known as *injections*, or solid ones known as *suppositories*, or, in case of the urethra, as *bougies*, are employed.

For the purpose of making local applications to the respiratory organs, *atomization* is very commonly practiced. Very many forms of apparatus are in use, but the principle in all of them is the same. A rapid current of air, or of steam, is forcibly projected from a horizontal pipe, through a capillary orifice, directly across a similar opening in a vertical tube. The rush of the vapor over this second orifice forms a vacuum; the fluid into which the base of the vertical tube is set, rushing up to fill this, is sucked or drawn out through the orifice, and as it emerges is broken into a fine spray, and is carried along by the current of air or steam into a mouth-piece, at which sits the patient. It cannot be gainsaid that in this way we are able to carry medicinal substances not merely into the larynx, but into the lungs themselves. Volatile medicines vaporized by heat are also sometimes employed in the treatment of lung affections.

There are various classes of agencies which so modify the action of drugs as to necessitate consideration of them here. The chief of these are disease, climate, habit, temperament, idiosyncrasies, sex, age, time of administration, and emotions.

Diseases often fortify the system against the action of remedies, so that the dose has to be greatly increased to obtain perceptible effects. Thus, pain or delirium tremens will interfere greatly with the production of narcotism by opium; or spinal disease with purgation. Disease may altogether prevent the action of a remedy. In all these cases two rules should never be lost sight of: first, never give the medicine in such doses as would in health cause death; second, always be sure, before giving large amounts, that the remedy will not make matters worse (as a drastic in intussusception).

Climate, by giving physical habits or tendencies to the patient, often greatly influences the proper selection and dose of remedies. It is only necessary to allude to the great consumption of quinine in malarial regions as an example of this.

Habit—including mode of life—seems to alter, as it were, the very constitution of man. Not only does it give type to disease, by producing habitual plethora, or its opposite, but it also fortifies against the action of single remedies, or whole classes of them. Thus, in the opium-eater, a dose sufficiently large to kill an ordinary man serves only to gratify the cravings of appetite. Again, a man accustomed to one narcotic, as alcohol or opium, loses, to a greater or less degree, his susceptibility to all narcotic influence; and the patient whose bowels require daily to be moved by a cathartic, finds himself react more and more slowly to medicines of that class. Again, a nervous system blunted by exposure and toil in the open air is far less susceptible to the action of remedies, and requires larger doses to influence it, than the delicate organization of a woman weakened by indolence and luxury.

Temperaments are peculiarities of organization characterizing classes of individuals; *idiosyncrasies*, peculiarities belonging to single individuals. This is scarcely the place to discuss the subject of temperaments, but it is allowable to state that while the *phlegmatic* person is no more easily moved by medicinal than by other agencies, the *nervous* individual answers as quickly to the one as to the other. Idiosyncrasies seem at present to be beyond law. They are often very remarkable, and a knowledge of them is most important for the practitioner. Thus, a relative of the author's is thrown into the most alarming fainting-fits by eating even so much butter as would be ordinarily used as a dressing for vegetables at dinner. Some persons are poisoned by the slightest touch of turpentine, others are frightfully salivated by a mere particle of a mercurial. These idiosyncrasies are numerous, cannot be foreseen, and are often very important: hence the necessity, in prescribing for an unfamiliar patient, of always asking as to his or her peculiarities.

Sex, of course, modifies all diseases connected with the organs or the process of generation, but it does more. Woman is more impressible, less robust, with less power of resisting external agencies, than is man. Consequently, the dose for her should, as a rule, be less than that for him. It is needless to remark here at length on the necessity for abstinence from strongly perturbing remedies during pregnancy, or at the menstrual periods.

Age, of course, modifies materially the dose. The rule of Dr. Young, the one which seems to me the most practical and generally useful, is to add twelve to the age and divide by the age. Thus, a child one year old would require one-thirteenth, one three years old, one-fifth, of the amount necessary for an adult. Other rules have been invented, but the only one which seems to me practical is the following, proposed by Dr. R. O. Cowling (*American Practitioner*, vol. i.):

"The proportionate dose for any age under adult life is represented by the number of the following birthday divided by twenty-four:" *i.e.*, for one year is $\frac{2}{24} = \frac{1}{12}$; for two years, $\frac{3}{24} = \frac{1}{8}$; for three years, $\frac{4}{24} = \frac{1}{6}$; for five years, $\frac{6}{24} = \frac{1}{4}$; for eleven years, $\frac{12}{24} = \frac{1}{2}$, etc.

Prof. Clarke (*Boston Med. and Surg. Journal*, 1872) has proposed a rule,

which, although probably more accurate than either of those given, seems too cumbersome for ordinary purposes. It is based upon relative weights :

“ Assuming the average weight of an adult to be one hundred and fifty pounds, for whom an appropriate dose is 1, or one drachm, the dose of most medicines must be increased or diminished in the proportion of the weight of the patient to that number of pounds. This proportion is represented by a fraction whose numerator is the patient's weight, and whose denominator is 150. If a child at birth weighs six pounds, the appropriate dose for it would be $\frac{6}{150}$, or $\frac{1}{25}$; if it weighs ten pounds, $\frac{10}{150}$, or $\frac{1}{15}$. A child two years old, weighing twenty pounds, would require $\frac{20}{150}$, or about $\frac{1}{7}$ of an adult dose; or, more precisely, $\frac{1}{7\frac{1}{2}}$. A person whose weight is two hundred pounds should have $\frac{200}{150}$, or $1\frac{1}{3}$ of an average adult dose.”

It must never be lost sight of that children bear narcotics very badly, and that the doses of such remedies for them should always be proportionally smaller than for the adult.

Time of Administration.—Absorption takes place most rapidly in an empty stomach, and consequently, when rapidity of action is desired, the medicine should be given under such circumstances. Thus, a purgative acts soonest when given before breakfast. Substances which are irritating to the stomach should always be administered not only properly diluted, but also when the viscous is filled by a mass of food, which may serve still further to lessen their concentration. Hence, such remedies as iodine and arsenic are preferably exhibited after meals. Again, some drugs, such as iron, are best dissolved by the acid gastric juice, and it is a matter of some importance to place them in the stomach after eating, when the process of digestion is most vigorous.

Mental Emotion.—Space is wanting to discuss at any length the influence of the imagination upon the action of remedies; and the reader must be referred to the delightful book of Dr. Tuke for illustrations. Suffice it to state that a positive announcement that a remedy will have a certain effect has often a most remarkable influence in producing that effect, especially on persons of nervous organization and of not too great culture to have faith. I have given a hypodermic injection of a grain of morphia to a man, inducing a degree of hypnotism, and the next day, doubling the size of the injection but withdrawing all morphia, have caused a much more intense effect.

It is evident that every treatise upon pharmacology must have some plan in accordance with which the various substances treated of are arranged. Most, if not all, of the classifications heretofore adopted may be brought under one of three heads,—alphabetical, natural historical, physiological. In the United States Dispensatory the first is adopted. In Pereira's famous work the drugs are arranged according to the natural history relations of the entities that produce them. It is evident that neither of these methods gives any aid to the memory of the student of therapeutics; and hence both are to be discarded in a treatise not encyclopædic in its aims. The physiological clas-

sification has the advantage of being seemingly scientific, and is certainly also more convenient for the purposes of teaching than the other two; yet it has serious drawbacks. If the reader will refer to almost any two works arranged upon such plan, he will see the same drug placed in two absolutely opposing classes. Thus, in Prof. George B. Wood's *Treatise upon Therapeutics* chloroform is included among the cerebral *sedatives*; and in Prof. Carson's "*Synopsis*" it is a cerebral *stimulant*. Again, by the physiological classification, medicines most closely allied in their clinical uses are often widely separated, whilst others very diverse are closely united. Thus, opium and belladonna are certainly in some sense antagonistic, yet they are always considered together; whilst chloroform and ether, used in practice for the one purpose, are often widely separated and placed in even antagonistic classes. I do not want to occupy too much space upon this matter, but can scarcely refrain from expressing the belief that a scientific, accurate, physiological classification of drugs is, in the present state of our knowledge, impossible; and, further, that it is very probable it will always so remain. So many medicines are stimulants and sedatives at the same time, stimulant to the one function and sedative to the other, the relations between the system and medicines are so complex,—our knowledge in the future will probably remain, as it is in the present, so fragmentary,—that the idea of a fixed physiological classification seems to me little more than a dream.

Certainly, at present, all that can be reasonably demanded of any system is that it be a convenient row of pegs upon which to hang our ideas and facts, so that they may be easily retained, and be easily accessible when wanted. Under these circumstances I venture to offer, as the basis of the present treatise, without further comment, the following clinico-physiological classification, founded upon, but very different from, that of Prof. George B. Wood.

CLASSIFICATION.

I. SUBSTANCES WHICH ACT ON THE SOLIDS AND FLUIDS OF THE BODY.

a. *General Remedies.*

CLASS	I.—Astringents.
"	II.—Tonics.
"	III.—Cardiac Stimulants.
"	IV.—Cardiac Sedatives.
"	V.—Antispasmodics.
"	VI.—Analgesics.
"	VII.—Mydriatics.
"	VIII.—Anæsthetics.
"	IX.—Excito-motors.
"	X.—Depresso-motors.
"	XI.—Alteratives.

b. *Local Remedies.*

CLASS	I.—Emetics.
“	II.—Cathartics.
“	III.—Diuretics.
“	IV.—Diaphoretics.
“	V.—Expectorants.
“	VI.—Emmenagogues.
“	VII.—Oxytocics.
“	VIII.—Sialagogues.
“	IX.—Errhines.
“	X.—Epispastics.
“	XI.—Rubefacients.
“	XII.—Escharotics.
“	XIII.—Demulcents.
“	XIV.—Emollients.
“	XV.—Diluents.

II. SUBSTANCES WHICH ACT EXTERNALLY TO THE BODY.

CLASS	I.—Antacids.
“	II.—Anthelmintics.
“	III.—Digestants.
“	IV.—Absorbents.
“	V.—Disinfectants.

PART I.

SYSTEMIC REMEDIES.

DIVISION I.—GENERAL REMEDIES.

CLASS I.—ASTRINGENTS.

ASTRINGENTS are those drugs which cause contraction of living tissues. That they do not act, as has been supposed, either by coagulating albumen or by calling into action the muscular function, is demonstrated by the transitoriness of their effects, and by the fact that they influence tissues containing no muscular fibre. Every living soft tissue appears to possess a normal degree of condensation, which may be departed from on either hand; when this happens, the part is said to be relaxed in the one case, in the other to have its tonicity increased, or to be astringed.

The action of astringents is always a *local one*, *i.e.* produced not through the intervention of the nervous system, but by direct contact with the part affected.

A pure astringent should be capable of doing nothing beyond inducing contraction; but in reality there is scarcely such a drug. All astringents are, when applied too freely, irritants; indeed, it is doubtful whether their therapeutical property of astringency is not due to the exercise of a mild form of irritation.

It has been customary to divide the astringents into two groups: the vegetable and the mineral. Tannic acid, and the drugs which contain it, constitute the former class, whilst in the latter group are included various substances of diverse therapeutic power.

The indications for the use of an astringent are very evident.

In the first rank among such indications is the *existence of relaxation*. Local relaxation is almost always the result of previous over-excitement. Thus, a throat is relaxed after over-use, or after inflammation.

Astringents are more efficient as local than as general remedies, but in cases of inflammation care must be taken to use them in such way that they shall not act as irritants. Applied too soon or too vigorously, they may do harm. These remarks are scarcely applicable to some of the mineral astringents, such as lead and nitrate of silver, which really appear to have sedative properties, and may with care be used advantageously in all stages of inflam-

mation, whenever there is distention and relaxation of the blood-vessels, although the general action of the part be above *par*.

Closely allied to relaxation is *over-secretion*, and astringents are constantly used to *check morbid discharges*. Indeed, these discharges are often simply the result of relaxation. Thus, Asp has experimentally proven that division of the intestinal nerves and consequent paralysis and relaxation of the vessels are followed by free watery secretion. In such cases the indication for astringents is very plain. But when a morbid discharge really represents a high degree of inflammation, the same care must be practiced in the use of astringents as in treating other local inflammations. Especially is this true since free secretion is often nature's method of relieving local inflammation. Thus, when abnormal alvine discharges are dependent upon intestinal relaxation, the astringents are most valuable, but when they are dependent upon enteritis or colitis, astringents do harm.

If the morbid discharge by its profuseness endanger life, as in serous diarrhœa, astringents are urgently demanded. Very rarely, if ever, are these discharges other than paralytic in their origin; even, however, if they be due to over-action, an astringent may be necessary to check their excessiveness.

Another indication for the use of astringents is to *check hemorrhage*, and the same general reasoning is applicable to this as to the other indications. Hemorrhage dependent upon over-action demands other treatment than by astringents. Sometimes in these cases it is necessary, however, to check the hemorrhage at all hazards, and then astringents may be used in conjunction with other measures, although they may be to some extent contra-indicated. Various astringents are employed locally to check hemorrhage due to traumatic or other ruptures of vessels. In such cases the astringents are employed as *styptics*, and do not act so much by their astringency as by coagulating the albumen of the blood, and thus forming a clot and mechanically arresting the flow.

Under certain circumstances there seems to be a general relaxation or loss of tone throughout the whole system, which may be best met by a consentaneous use of tonics and astringents.

VEGETABLE ASTRINGENTS.

The active principle of the vegetable astringents is tannic acid, and, as it is almost their sole therapeutic principle and represents them very closely, it seems proper first to consider it, and afterwards to point out any especial therapeutic virtues the crude drugs of the class may possess.

ACIDUM TANNICUM—TANNIC ACID.

There are two kinds of tannic acid, the *gallo-* and the *kino-tannic*; of these the former yields, upon exposure to the air in a moist state, *gallic acid*, the latter a *gelatinous, inert* substance. They are further distinguished by the

color of the precipitates which they yield with the persalts of iron; gallo-tannic acid producing a blue-black, kino-tannic a green-black color.

The official tannic acid is derived from galls, and is therefore the gallo-tannic acid; to it subsequent remarks especially refer.

It is obtained by treating powdered galls with washed ether, which on standing separates into two strata, the upper of which is ethereal, and contains chiefly the coloring-matter and other impurities. The lower watery stratum contains the tannic acid, which is recovered by evaporation.

Commercial tannic acid is a light, feathery, *non-crystalline* powder, of a yellowish-white color, a faint odor, and an astringent, somewhat bitter taste. When absolutely pure, it is colorless and free from odor or taste other than that of astringency. Its reaction is strongly acid, and it unites freely with both organic and inorganic bases. It is very freely soluble in water, even more so in glycerine, somewhat so in dilute alcohol, scarcely at all in absolute alcohol, and not at all in ether free from water. By a heat of from 180° C. to 215° C. it is changed into *pyro-gallic acid*, which crystallizes in white, shining plates, of a bitter taste and neutral reaction.

Tests.—Tannic acid produces, with salts of the alkaloids, a whitish precipitate, very soluble in acetic acid; with persalts of iron, a black (bluish or greenish) precipitate; with lime-water, a precipitate which is at first whitish, then gray, dingy greenish, and finally brownish; with gelatine or albumen, a whitish coagulum. All of these secondary products are tannates. Tannic acid also dissolves in concentrated sulphuric acid, with the production of a black color. By prolonged exposure in solution to the air, or by the action of dilute sulphuric acid, it is converted into gallic acid.

PHYSIOLOGICAL ACTION.—Tannic acid is, so far as may be, a pure astringent. Owing to its property of coagulating albumen, it cannot be absorbed into the blood, and its solution injected into the veins causes rapid death, with convulsions, or other nervous phenomena, due to the production of thrombi or emboli. Out of the body it is readily converted into gallic acid, and, when taken into the stomach, a proportion of it undergoes a similar change. In the viscera of a rabbit poisoned with tannic acid, Schroff (*Die Pflanzenstoffe*, von Dr. Aug. und Dr. Theo. Husemann, p. 1005) found only gallic acid; and according to Clarus (*ibid.*) the greater part of ingested tannic acid can be recovered from the stools as tannate of albumen or as gallic acid. Wöhler and Frerichs have found gallic acid with pyro-gallic acid in the urine after the exhibition of tannic acid.

THERAPEUTICS.—As tannic acid must undergo conversion into gallic acid before absorption, it is evident that the latter is to be preferred to it when the part to be acted on can only be reached through the circulation. As a local application, tannic acid is much more powerful than gallic acid. Locally applied, it may be used to overcome relaxation, as in *spongy gums*, *mercurial sore mouth*, *hemorrhoids*, *chronic sore throat*. To check hemorrhage it may be used whenever the source of the flow can be reached directly, as in *epis-*

taxis, *hæmatemesis*, *hemorrhage from the bowels*, etc. To arrest *excessive secretion* it may be employed locally in *leucorrhœa*, *diarrhœa*, *old abscesses*, *chronic ulcers*, *excessive perspiration*, *osmidrosis*, and various diseases of the skin. It is also often very useful for the purpose of hardening parts exposed to friction, as in cases of *sore nipples* and *tender feet*.

TOXICOLOGY.—Tannic acid can scarcely be called poisonous, although Rollett reports the case of a young girl in whom a very large quantity of it induced severe gastric and abdominal pains, with obstinate vomiting and constipation, fever and general malaise. Both Schroff and Judell assert that eighty grains of it produce no symptoms of importance in the rabbit.

As an antidote it is useful in tartar emetic poisoning, forming an insoluble tannate of antimony. It is also the best chemical antidote for the poisonous alkaloids; but as the compounds it makes with them are slowly dissolved by the fluids of the alimentary canal, it must always be followed by emetics and cathartics.

ADMINISTRATION.—When given to act on the stomach, as in hæmatemesis, tannic acid should be in powder (ten to twenty grains). When the bowel is to be influenced, as in diarrhœa, the drug should be administered in pill (three to five grains), so that, if possible, it may pass the pylorus undissolved. For external use two preparations are officinal:

Glyceritum Acidi Tannici—*Glycerite of Tannic Acid* (1 part to 4).

Unguentum Acidi Tannici—*Ointment of Tannic Acid* (grs. xxx to 3i).

ACIDUM GALLICUM—GALLIC ACID.

Galic acid is a white, powdery substance, in fine acicular prisms, soluble in one hundred parts of cold water, in three parts of boiling water, and freely soluble in alcohol and ether. Its taste is acidulous and somewhat astringent.

According to the officinal method, gallic acid is prepared by the exposure of moistened powdered nut-galls in a warm place for a month. A species of fermentation, with the development of a peculiar fungus, is said to occur, during which oxygen is absorbed, carbonic acid evolved, and glucose and gallic acid are produced. M. Sacc (*Chemical News*, July 24, 1871) has recently denied this, affirming that the change is simply one of hydration, tannic acid being an anhydride of gallic acid. Tannic acid also may be rapidly converted into gallic acid by the action of dilute sulphuric acid.

Galic acid produces with salts of the alkaloids whitish precipitates, with persalts of iron a bluish precipitate, with lime-water a whitish precipitate, changing to blue, and then to violet or purplish,—all of these precipitates being gallates. It does *not coagulate gelatine* or *albumen*, and dissolves in concentrated sulphuric acid, with production of a deep red color. It has the power of reducing silver from its solution slowly in the cold, instantaneously if warmed.

PHYSIOLOGICAL ACTION.—Gallic acid as an astringent is similar to, but

much less powerful than, tannic acid. It escapes from the body through the kidneys.

THERAPEUTIC ACTION.—Gallic acid is not nearly so efficient as tannic acid when applied locally, but, because it does not coagulate albumen, should always be preferred when the part is to be reached through the medium of the circulation. It is useful as an astringent in *hæmatemesis*, *hæmaturia*, *colliquative sweats*, etc. It has been recommended in *bronchorrhœa* and in the profuse expectoration of *chronic phthisis*. In my hands, however, it has completely failed in the latter affections. In certain forms of *Bright's disease*, when there was an abnormally large secretion of highly albuminous urine, I have found it to lessen very materially the excretion of albumen.

ADMINISTRATION.—Gallic acid may be given in powder, or sometimes in pill form. The dose of it is from ten to thirty grains, repeated as often as may be necessary. The following preparation is officinal:

Glyceritum Acidi Gallici—*Glycerite of Gallic Acid* (1 part to 4).

GALLA—GALLS.

Galls are vegetable excrescences, which are produced by the deposition of the ova of insects. They occur on almost all kinds of plants, even on fungi; but the officinal gall is developed on the *Quercus infectoria* by the act of the fly *Cynips* (*Diplolepis*) *tinctoria*. Our supply of them is derived chiefly from the Levant.

There are two varieties of galls. The *blue* or *green galls* are globular, solid bodies, from the size of a pea to that of a hickory-nut, externally smooth, or more commonly marked with large tubercles. They are the young galls which have been gathered before the ova of the fly have hatched, or before the caterpillar has eaten out the interior of its birthplace. The *white galls* are large, light, hollow bodies, with a hole, through which the *Cynips* has escaped after having fed upon the interior during its whole larval life. They contain but little tannic acid, and are of comparatively little value.

THERAPEUTICS.—The sole value of galls is as the source of tannic acid. As galls, they should not be used in medicine; but the United States Pharmacopœia indicates the following preparations:

Tinctura Gallæ—*Tincture of Galls* (℥iv to Oj).

Unguentum Gallæ—*Ointment of Galls* (1 part to 7).

CATECHU—CATECHU.

An extract of the wood of an East Indian tree,—the *Acacia Catechu*. It occurs in masses of various shapes, or in small fragments, of a dull reddish-brown color, and having a bitterish, astringent, and, after a time, sweetish taste.

It contains kino-tannic and catechuic acid.

Pale catechu, or *gambir*, which is officinal in the British but not in the United States Pharmacopœia, occurs in small cubes, about an inch in diam-

eter, lighter than water, pale-yellowish within, deep-yellowish or reddish-brown externally.

THERAPEUTICS.—A powerful astringent, which may be used externally or for diarrhœa.

ADMINISTRATION.—The dose of the powder is twenty to thirty grains. The following preparations are official:

Infusum Catechu Compositum—*Compound Infusion of Catechu* (Catechu, ℥ss, Cinnamom., ℥i to Oj). Useful in *diarrhœas* of relaxation. Dose, one to two fluidounces.

Tinctura Catechu—*Tincture of Catechu* (Catechu, ℥iii, Cinnamom., ℥i to Oij). Dose, one to two fluidrachms.

KINO—KINO.

The inspissated juice of *Pterocarpus marsupium* and of other plants. It occurs in small, irregular, angular, shining, reddish, brittle fragments, of a bitterish, highly astringent, and, after a time, sweetish taste. There are four varieties,—the East India, West India, Botany Bay, and African. Of these the first is common, the second rare, and the last two are never seen in our market.

Kino contains kino-tannic acid, and in its therapeutic powers is almost identical with catechu.

ADMINISTRATION.—The dose of the powder is twenty to thirty grains. The following preparation is official:

Tinctura Kino—*Tincture of Kino* (℥vi to Oss). Dose, one fluidrachm.

HÆMATOXYLUM—HÆMATOXYLON.

The heart-wood of *Hæmatoxylon Campechianum*, or logwood-tree, a native of Central America. A dense, heavy wood, of a deep reddish-brown color, containing, besides kino-tannic acid, a crystalline principle, *Hæmatin* or *Hæmatoxylin*, which when pure is yellow, but readily yields red or purple dyes.

THERAPEUTICS.—A mild, efficient astringent, valued on account of its sweetish taste. It is readily taken by children, but is sometimes objected to on account of the staining of the diapers by the blood-red stools which it produces. The following formula offers an efficient and elegant remedy for diarrhœas of relaxation; the proportions may be varied to suit individual cases:

R Ext. hæmatoxyli, ℥ii;
 Acid. sulph. aromat., f℥iii;
 Tinct. opii camph., f℥iiss;
 Syrupi zingiberis, q. s. ad f℥vi.

M.—Dose, a tablespoonful, properly diluted.

ADMINISTRATION.—*Extractum Hæmatoxyli*—*Extract of Hæmatoxylon*. Dose, ten to thirty grains.

KRAMERIA—RHATANY.

The root of *Krameria triandra*, a native shrub of Peru. This woody root, as it occurs in our markets, varies from one-fourth inch to one inch in diameter, and from half a foot to three feet in length. The readily separable bark is of a deep reddish color. The internal woody portion is of a lighter hue, although decidedly reddish. The bark contains a much larger percentage of the active principle, kino-tannic acid, than the wood.

Rhatany is a powerful astringent, similar in virtue to kino and catechu.

ADMINISTRATION.—*Krameria* is never administered in powder but in one of the following official preparations:

Extractum Kramerizæ—*Extract of Rhatany*. Dose, grs. v–x.

Infusum Kramerizæ—*Infusion of Rhatany* (3i to Oj). Dose, one to two fluidounces.

Syrupus Kramerizæ—*Syrup of Rhatany*. Dose, half a fluidounce.

Extractum Kramerizæ Fluidum—*Fluid Extract of Rhatany*. Dose, twenty drops.

The United States Pharmacopœia recognizes the following astringent vegetable drugs not yet mentioned:

QUERCUS ALBA and *QUERCUS TINCTORIA* are the inner barks of the trees whose names they bear,—the *white* and the *black oak* respectively. The latter is a rough, yellowish-brown bark, which is used in dyeing, under the name of *quercitron*. On account of its imparting readily its color, it is rarely, if ever, employed in medicine. White-oak bark also stains, but not nearly so deeply as black-oak bark, and, containing largely of gallo-tannic acid, is used as a means of making cheap astringent infusions for baths, vaginal washes, etc., also in powder for poultices. There is an official decoction of the white oak.

ROSA GALLICA is the dried petals of the generally half-opened flowers of the hundred-leaved rose. They are of a deep red color, of a pleasant scarcely astringent taste, and contain a small percentage of gallo-tannic acid, red coloring-matter, and a trace of volatile oil. Sulphuric acid changes their infusions or tinctures to a bright red color. They are almost destitute of therapeutic virtues, but their preparations are used as elegant vehicles. The following are the official preparations:

Infusum Rosæ Compositum—*Compound Infusion of Rose*. It contains sulphuric acid, and affords an elegant acid astringent vehicle.

Confectio Rosæ—*Confection of Rose*. Used as a vehicle.

Syrupus Rosæ Gallicæ—*Syrup of Red Rose*. An elegant vehicle.

The *pale rose* has no astringency, and will be considered under Emollients.

The rhizome of *Geranium maculatum* Linn., an herbal plant, which grows abundantly in open woods in the middle United States, and may be recognized by its light purplish petals, slender pointed sepals, and five-parted leaves, is

official under the name of *Geranium*. It occurs in pieces from one to three inches long, one-quarter to one-half inch in thickness, wrinkled, contorted, tuberculated, often fibrillated, brownish externally, grayish internally. The taste is a nearly pure astringent one. It contains largely of gallic and tannic acids, and is a somewhat popular astringent, although rarely used by practitioners. It may be boiled in milk for children needing a mild astringent. Dose, grs. xx-xxx.

RUBUS CANADENSIS—DEWBERRY ROOT; RUBUS VILLOSUS—BLACK-BERRY ROOT, both indigenous, and feeble, not much used astringents, containing gallo-tannic acid. TORMENTILLA is the root of *Potentilla Tormentilla* or *Septfoil* of Europe: although it contains kino-tannic acid and is a powerful astringent, it is never used in this country. DIOSPYROS, the unripe fruit of *Diospyros Virginiana* Linn., or Persimmon. GRANATI FRUCTUS CORTEX and GRANATI RADICIS CORTEX, the rind of the fruit and the bark of the root of the *Pomegranate*, powerful astringents, containing gallo-tannic acid, rarely used in the form of decoctions as topical remedies.

MINERAL ASTRINGENTS.

ALUMEN—ALUM.

Formerly the double salt of alumina and potash constituted the ordinary alum as well as the officinal drug. Ammonia as a secondary product in the manufacture of coal gas has become so cheap, however, that it is now used very largely instead of potash, and the ammonia alum is the officinal *alum*, the potash salt being retained under the name of *Aluminii et Potassii Sulphas*. The two salts are identical in physical and medical qualities, but when the ammonia alum is triturated with lime the odor of ammonia is at once evolved. The formulæ of the two salts are, according to the old nomenclature, $\text{Al}_2\text{O}_3 \cdot 3\text{SO}_3 + \text{NH}_4\text{OSO}_3 + 24\text{HO}$, and $\text{Al}_2\text{O}_3 \cdot 3\text{SO}_3 + \text{KOSO}_3 + 24\text{HIO}$; according to the recent system the formulæ are $(\text{SO}_4)_2\text{AlK} + 12\text{OH}_2$ and $(\text{SO}_4)_2\text{AlNH}_4 + 12\text{OH}_2$. Alum occurs in octahedral colorless crystals, which are often aggregated into large masses. Its taste is astringent, acidulous, and sweetish. It is soluble in about fifteen times its weight of cold and in about three-fourths its weight of boiling water. It is slightly efflorescent, and when heated a little beyond 112° Fahr. parts with its water of crystallization, and is converted into a white powder, which is officinal as *Alumen Exsiccatum*, or *Dried Alum*. The alkalies and their carbonates, lime, magnesia and its carbonate, tartrate of potassa, and acetate of lead are incompatible with alum.

PHYSIOLOGICAL ACTION.—As alum, even in very dilute solutions, coagulates albumen, it would appear as though it could not be absorbed. Since, however, both Drs. Geo. B. Wood and A. Stillé assert, on what authority I do not know, that alumina can be detected in the urine of persons taking it, it or its derivatives must find a way into the blood. What changes it undergoes in the alimentary canal, or in what form it enters the blood, is not known.

Applied to a tissue, it acts as a very powerful astringent and irritant. Orfila found that in dogs one or two ounces of it simply induce violent vomiting and purging, whilst in Mitscherlich's experiments two drachms of it produced in rabbits fatal gastritis, evidently on account of their inability to vomit.

THERAPEUTICS.—Alum may be used locally to serve all the purposes of a very active astringent. It has been employed very frequently with success as a styptic to arrest hemorrhage; and, applied by the atomization of its saturated solution, I have found it of signal service in *hæmoptysis* and in *bronchorrhœa*. It also frequently enters into the composition of gargles for *sore throat*; but this practice is to be reprobated, since alum acts very destructively on the teeth. In *colliquative sweats* sponging at bed-time with alum-water, or, still better, the taking of an alum-water bath, will often materially aid in restoring the lost tone to the skin. In *chronic ulcers* with exuberant spongy granulations, and in certain conditions of *conjunctivitis*, it is often applied with benefit in the form of *alum curd*. In *diarrhœa*, and indeed generally when it is desired to exert an astringent action upon the internal organs, alum is not nearly so useful as other members of the class.

Owing to its irritant properties, alum when given in sufficient amount acts as a mechanical emetic, and may be used as an adjuvant to the sulphate of zinc or of copper in narcotic poisoning. Originally introduced by Dr. C. D. Meigs, it is believed in *membranous croup* to be of service not only by its emetic action, but also by modifying the mucous membranes with which it comes in contact in its passage down and up.

So long ago as the last century, Dr. Grashius, of Holland, commended alum in *colica pictonum*, and although for a long time its value was not recognized, abundant confirmative testimony has recently been brought forward. Since it is a soluble sulphate, it is of course a chemical antidote to any lead salt which may be in the alimentary canal. It is, however, of service when there is no lead in the primæ viæ, and must act in some way as yet unknown. It has, indeed, been used with asserted success in other neuroses of the alimentary canal,—in *gastralgia* and in *intestinal neuralgia*. Dr. Aldredge even commends it in *habitual constipation*. *Burnt alum* is used as a very mild escharotic for exuberant granulations in ulcers.

ADMINISTRATION.—As an astringent, the dose of alum is from ten to twenty grains; as an emetic, a teaspoonful of the powder for a child, a table-spoonful for an adult, in syrup, repeated in fifteen minutes; in *colica pictonum*, twenty to forty grains every three or four hours, combined with morphia. Alum curd may be made by dissolving two drachms in a pint of milk, and straining, or by rubbing the alum with white of egg.

ALUMINII SULPHAS—SULPHATE OF ALUMINIUM (Al_2O_3 ; 3SO_3 — Al_2 ; 3SO_4 .)—This salt, which occurs as a white powder, or in lamellated cakes, or in a crystalline cake, is used externally as a powerful astringent and antiseptic. Its solution has also been employed by injection for the preservation of cadavers.

PLUMBUM—LEAD. (Pb.)

When a soluble salt of lead is applied to a part in not too concentrated solution, it acts as an astringent and sedative. Owing to the contraction of the vessels which is induced, the tissue becomes blanched, and any inflammatory action which may be present is remarkably affected. When in concentrated solution, however, the mildest salts of lead are capable of acting as irritants, increasing or even originating inflammation.

When the salts of lead are taken internally in therapeutic doses, no decided symptoms are generally induced, except a diminution of the secretions, especially of those of the alimentary canal. Sometimes, when full therapeutic doses are exhibited, a slight lowering of the frequency and force of the pulse (see Laidlaw's Observations, quoted by Stillé, *Therapeutics*, second edition, p. 177, vol. i.) is said to result, but I have never witnessed this. The insoluble preparations of lead act similarly but less decidedly than the soluble; yet in regard to some of them it is doubtful whether they can under any circumstances become irritant.

TOXICOLOGY.—The salt of lead with which intentional or accidental acute poisoning is most frequently induced is the acetate.* The first result of a toxic dose of this is in most cases a persistent sweet, somewhat metallic taste; this in a few minutes is followed by vomiting, which may or may not be preceded by nausea. The matters vomited are often milky-white, from the presence of chloride of lead. A severe burning persistent pain in the abdomen now comes on, and is accompanied with a craving for drink. There may be obstinate constipation, or diarrhoea may ensue; in either case the stools are generally black, from the sulphuret of lead. In certain cases a state of collapse is developed; the pulse falls to forty or fifty per minute, the voice is lost, the face is deadly pale, the lips livid, and syncope seems imminent. In other instances the nervous symptoms may predominate, or they may accompany those of disordered circulation: cramps in the calves of the legs, severe neuralgic pains in the extremities, paralysis and anaesthesia, vertigo, stupor, may any or all of them be present. In fatal cases, coma, with or without convulsions, finally develops itself. A distinctive mark of lead-poisoning, which sometimes is present very early, is the blue line upon the gums. After death inflammation of the alimentary mucous membrane is sometimes, but not always, found.

The fatal dose of sugar of lead is between one and two ounces. The subacetate of lead is even more poisonous than the acetate, and the nitrate acts as a violent irritant. The carbonate appears to be incapable of causing acute poisoning.

The treatment of acute lead-poisoning consists in the evacuation of the

* According to Husemann (*Handbuch der Toxicologie*), the Poudre de Succession, so famous during the reign of Louis XIV., was composed chiefly of acetate of lead.

stomach, if necessary, the exhibition of the sulphate of soda or of magnesia, and the meeting of the indications as they arise. The Epsom and Glauber's salts act as chemical antidotes, by precipitating the insoluble sulphate of lead, and also, if in excess, empty the bowel of the compound formed. To allay the gastro-intestinal irritation, albuminous drinks should be given and opium freely exhibited.

Of all forms of poisoning, chronic lead-poisoning is the most common. It is almost always accidental, and occurs most frequently among those whose occupation exposes them to daily contact with some compound of the metal, manufacturers of white lead, painters, glaziers, and similar artisans furnishing the greater number of victims. It is seen, however, in persons of all conditions of life, for although neither food nor drink is often purposely adulterated with it, yet it is frequently introduced into the system accidentally along with those necessities. Lead pipes are habitually used for the conveyance of water, and when that water contains salts of lime, even in minute proportion, no evil results, because through the decomposition which ensues insoluble coatings are deposited on the inside of the pipes.* When the water is pure, no such reactions occurring, the lead is slowly dissolved in the form of a carbonate, and poisoning may result. Poisoning has also frequently resulted from the use of cosmetics and hair-dyes containing salts of the metal, from imperfectly burnt pottery (*Schmidt's Jahrbücher*, Bd. cxliv. p. 279), and in other still more curious ways.

The most ordinary form of chronic lead-poisoning is that in which colic is the first decided symptom. After some days of malaise and wretchedness, or sometimes very suddenly, the victim is taken with abdominal colicky pains, which increase in intensity until they become very severe. They are constant, with occasional exacerbations, are sometimes dull, sometimes sharp, are generally described as twisting, and seem to centre around the umbilicus. There is very often repeated retching and vomiting. The walls of the abdomen are retracted, rigid, knotted; the bowels are obstinately costive; the tongue is contracted and whitish; the appetite gone; and the thirst sometimes excessive. Neuralgic pains in the thorax and in the extremities are of frequent occurrence.

These symptoms usually abate after a time, but are very apt to recur with increasing severity, especially among those who are habitually exposed to the cause. In one of these repeated attacks severe cerebral symptoms may come on suddenly, or after some days of headache; delirium, either mild or maniacal, or stupor, is in some such cases the chief manifestation, but epileptiform convulsions are more common. These convulsions are often very severe, and end in coma, in which death may take place. The more ordinary nervous symptoms of lead-poisoning may be developed after the first attack of colic, or they may occur without being preceded by any marked abdominal dis-

* For an elaborate article on the chemical relation of water to lead, see *Schmidt's Jahrbücher*, Bd. cxliv. p. 279.

turbance. The most common of all of them is the local paralysis known as "drop wrist." This is a complete paralysis of the extensor muscles of the forearm, which allows the hand to drop forward. It is often associated with complete or partial anæsthesia of the part affected.

Among the rarer forms of nervous disturbance induced by lead may be mentioned *amaurosis* from atrophy of the optic nerve.

According to Dr. E. Levy (*Schmidt's Jahrbücher*, Bd. clii. p. 250), acute asthma is rarely produced by the inhalation of the dust of white lead, and chronic saturnine asthma is sometimes seen in feeble narrow-chested people. The epileptiform convulsions produced by lead are in some cases secondary to chronic nephritis, but in other instances are due directly to the action of the lead (see paper of Dr. Rosenstein, *Virchow's Archiv für patholog. Anatomie*, Bd. xxxix., 1867, p. 4). Upon pregnant women the influence of the poison is very deleterious, and Dr. Constantine Paul (*Archives Générales*, 5th series, vol. xv., 1860, p. 513) has shown that it very commonly produces the early death of the fœtus.

As any of these obscure manifestations of lead-poisoning may exist, and even prove fatal, without a distinct history of other more characteristic phenomena, great care is sometimes necessary to avoid being misled, and not rarely the true nature of saturnine epilepsy or of saturnine albuminuria is overlooked. Hence the importance of the *blue line upon the gums where they join the teeth*, which probably exists in a greater or less degree in all cases of lead-poisoning, although very great cleanliness is said to lessen or even to eradicate it.

In those cases of lead-poisoning which pursue a slow course to death, the paralysis involves after a time the extensors of the lower as well as of the upper extremities, epileptic paroxysms occur at intervals, racking pains shoot through the limbs, points of cutaneous anæsthesia appear, and often albuminuria aids in producing the fatal issue. Gradually the patient becomes more and more cachectic, general œdema, with the whitened skin, betrays the increasing anæmia, the paralysis extends from muscle to muscle, locomotion becomes impossible, and if a convulsion or other accident does not close the scene, death at last takes place from loss of power in the respiratory muscles.

After death lead has been frequently found in the tissues. Prof. Chatin (*Comptes-Rendus de la Société de Biologie*, tome iv., 3d series, 1862, p. 84) obtained from the deep-gray upper cervical cord of a patient of Prof. Trouseau three milligrammes of the sulphuret of lead in one hundred and fifty grammes of nerve-tissue. When persistent albuminuria has existed during life, the kidneys are found in a state of fatty or albuminous degeneration.

The paralyzed muscles are not only exceedingly wasted, but their structure may be so totally destroyed that scarcely a single striated fibre can be found. The nerve-trunks are lessened in size, in many of their tubules the medulla has been replaced by fatty granules, and in some cases every trace of the tubules has disappeared and the nerve been reduced to a fibrous cord. The

symmetry and localization of the paralysis in lead-poisoning would seem to indicate a spinal origin, but, so far as I know, no very distinct spinal lesion has as yet been made out. Dr. E. Lancereaux (*Comptes-Rendus de la Soc. Biolog.*, 1862, tome iv., 3d ser., p. 84), in a very careful study of a case, could only find some of the spinal cells more granular than normal, and apparently atrophied. In some cases of lead-poisoning no decided anatomical alteration can be anywhere detected.

The treatment of chronic lead-poisoning evidently arranges itself under three indications: 1st, to prevent the ingestion of more of the poison; 2d, to aid in the elimination of that in the system; 3d, to relieve symptoms and restore lost functions. In lead-colic both of the last two indications are met by purgatives, to which opium should be added to relieve pain. It is often necessary to use the most powerful drastics, such as croton oil; but senna, salts, and other of the milder cathartics should always be tried first. *Alum* acts in some unknown way as a specific in lead-colic, and from twenty to sixty grains of it may be given four or five times a day. In the more chronic forms of lead-poisoning, to fulfill the second indication baths of sulphuret of potassium should be employed, and iodide of potassium be administered internally. The bath should be given (Dr. A. Eulenburg, *Deutsches Archiv für Klin. Med.*, Bd. iii. p. 506) in a wooden tub, two or three times a week, and should contain six or seven ounces of the salt. The patient, during the half-hour of his continuance in it, should be from time to time well rubbed with a coarse towel. On coming out he is to be thoroughly washed with warm soap-suds. The dose of the iodide should be from fifteen to twenty grains, administered after meals, in dilute solution. When severe cerebral symptoms arise, treatment is of little avail, and should be largely expectant.

The local use of electricity is exceedingly important to restore the lost function of nerve and muscle. When the faradaic current elicits a response it should always be employed, but in some cases (*Meyer's Electricity*, New York, 1869, p. 284) the continued current retains its power after the induced has lost all its influence. The rule is always to apply that current which causes contraction; if both fail, the continued current should be used, the poles being reversed at intervals of four or five seconds. The electrical sances should be tri-weekly, each lasting about fifteen minutes, and they should be persevered in for months. I have seen great improvement in a case which for the first four months yielded no results; indeed, long after voluntary movement had in great measure returned, no form of electricity would cause contraction of the affected muscles.

The following preparations of lead are officinal in the United States Pharmacopœia:

PLUMBI OXIDUM—OXIDE OF LEAD. (PbO.)

Litharge, which is prepared by blowing air through melted lead, occurs in small yellowish or orange-colored scales, which are insoluble in all ordinary

menstrua. It is rarely used as a desiccant astringent powder for ulcers, but its chief employment in medicine is as the basis of the following preparations :

Emplastrum Plumbi.—*Lead Plaster* is made by boiling litharge, olive oil, and water together. Glycerine is set free, and the oleo-margarate of lead is formed. Lead plaster occurs in grayish, cylindrical rolls, which become adhesive at the temperature of the body, and, spread upon kid, is sometimes used as a protective to parts exposed to pressure, or to superficial ulcers or abrasions.

Emplastrum Resinæ.—*Resin Plaster*. *Adhesive or sticking plaster* is made by incorporating resin with lead plaster, and, spread upon linen, is much used in surgery for mechanical purposes.

Emplastrum Saponis.—*Soap Plaster*. This is made by the addition of soap to lead plaster. It is employed chiefly as a protective.

PLUMBI ACETAS—ACETATE OF LEAD. ($\text{PbO}, \text{C}_4\text{H}_3\text{O}_3$ — $\text{Pb}, \text{C}_2\text{H}_3\text{O}_2$.)

Sugar of lead is made by the action of acetic acid upon litharge, or upon sheets of lead exposed to the air. It occurs in transparent, acicular, often aggregated crystals, of a sweet, styptic taste. It is soluble in water, to which it usually imparts a slight milkiness. From its solution it is precipitated black by sulphuretted hydrogen, white by soluble carbonates, chlorides, and sulphates, and bright yellow by the iodide of potassium. It is also incompatible with the mucilage of slippery elm, but scarcely so with that of flaxseed or of pith of sassafras.

THERAPEUTIC APPLICATION.—A solution of the acetate of lead is used very largely in acute external *inflammations* as a sedative and astringent lotion. Although chemically incompatible, it is frequently combined very advantageously in these cases with opium. As a too concentrated solution acts as an irritant, the strength for use on the eye should not exceed one or two grains to the ounce; on the skin, ten grains to the ounce.

Internally, acetate of lead has been employed very largely in *hemorrhage*: indeed, Prof. George B. Wood commends it as the most valuable of all astringents in *hemoptysis* (*Therapeutics*, vol. i. p. 158). I think it is now, however, rarely used for this purpose.

Its chief use at present is in *diarrhœa*. On account of its sedative properties, when the purging is attended with inflammation, it is the most serviceable of all the astringents; and, owing to the promptness of its action, it is also very valuable in cases with profuse serous discharges. In *dysentery* it is very useful whenever the discharges have become copious.

ADMINISTRATION.—Acetate of lead should always be given in pill form. The dose is from two to five grains,—repeated every two, three, or four hours *pro re nata*.

PLUMBI CARBONAS ($2(\text{PbOCO}_2) + \text{PbOHIO} - \text{PbCO}_3$).—The *carbonate of lead* is a heavy, white, tasteless powder, which is insoluble in distilled


water, but slightly so in water containing carbonic acid. It is never used internally, but as an external sedative application. Rubbed up with linseed oil, it constitutes white-lead paint, and in this form, or in that of the ointment (*Unguentum Plumbi Carbonatis*), it is a most efficient dressing for fresh burns. Care must be taken in its use, however, when a large surface is involved, as lead-colic has been caused by its absorption.

PLUMBI NITRAS ($\text{PbO}, \text{NO}_3 - \text{Pb}_2\text{NO}_3$).—The *nitrate of lead* occurs in white, nearly opaque, octahedral, very heavy crystals, which are soluble in alcohol and in seven and a half parts of cold water. This salt is not used except as a disinfectant. Dissolved in water, it forms the so-called *Ledoyen's Disinfectant Solution*. It acts by decomposing the sulphuretted hydrogen, itself being converted into a sulphuret of lead. Its chemical reactions are similar to those of the acetate, from which it may be distinguished by a mixture of it and sulphuric acid striking a red color with morphia.

BISMUTHUM—BISMUTH. (Bi.)

The United States Pharmacopœia includes the metal bismuth in its *Materia Medica* list, and the subnitrate and subcarbonate among its preparations. The metal itself is never used in medicine.

BISMUTHI SUBCARONAS—SUBCARBONATE OF BISMUTH.



A white or yellowish-white powder, tasteless and odorless, totally insoluble in water, soluble with effervescence in dilute nitric acid. The mode of its preparation is too complicated for discussion here, the chief object of the various steps being to get rid of the arsenic, which very generally contaminates the metal. The same assertions may be made in regard to the preparation of the **BISMUTHI SUBNITRAS**, or **SUBNITRATE OF BISMUTH** ($\text{BiO}_3\text{NO}_5\text{HO} - \text{BiONO}_3\text{H}_2\text{O}$), a heavy white powder, odorless, with a faint acid taste, and a decidedly acid reaction when applied to moistened litmus-paper, insoluble in water, soluble without effervescence in nitric acid.

PHYSIOLOGICAL ACTION.—The actions of the subnitrate and of the subcarbonate of bismuth are so exactly similar that they can practically be considered as one thing; and for the sake of brevity I shall speak of these salts simply as bismuth, meaning thereby the officinal preparations of the metal. Orfila and other of the old observers attributed to bismuth violent irritant properties, stating that severe symptoms and even death sometimes followed its ingestion.

There can be no doubt that the results noted by these authorities were due not to the bismuth, but to arsenic with which it was contaminated. Notwithstanding all the care of modern pharmacy, bismuth even yet occasionally contains arsenic, and I have seen it produce from this cause bloody purging. When pure, however, it is free from any irritant properties, and is a feeble

astringent and even a sedative, and can be taken without injury in indefinite quantity.

The officinal salts of bismuth are so exceedingly insoluble that it has been generally believed that they are not absorbed; but the researches of MM. Bergeret and Mayençon (*Journal de l'Anatomie*, 1873, p. 242) strongly indicate the contrary to this. According to this authority, if a piece of paper be wet with a solution of sulpho-cyanide of potassium and dried, it affords a most sensitive and characteristic test for a soluble salt of bismuth,—a beautiful yellow spot appearing at the point of contact. Availing themselves of this test, MM. Bergeret and Mayençon have found that when the subnitrate of bismuth is administered the metal can always be detected after a few hours in the urine. They have also discovered it in the serous exudation of dropsy, and have proven that when a few grains of the salt mentioned are given to rabbits, in from twenty to thirty minutes it can be found in the urine, kidneys, spleen, blood, and muscles, and even eight days after the administration can be detected in all the tissues. Five days after the exhibition of a gramme of the subnitrate to a man, they found traces of the metal in the liver and kidneys; but the analysis of the body of a woman dead sixty-two days after the ingestion of two grammes yielded only negative results.


THERAPEUTICS.—The preparations of bismuth are of great service in various forms of irritation of the alimentary canal. They are very useful to allay *vomiting* dependent upon gastric irritation. In simple *neuralgic gastric pain* following eating, especially when occurring in feeble, badly-nourished subjects, bismuth is often of great service; and even in *carcinoma* it may palliate by alleviating pain and vomiting. In *pyrosis* it is sometimes successful. In simple *diarrhœa* of irritation, and in the *chronic diarrhœa* of camps, the bismuth preparations are often very efficient; and in the *chronic bowel complaints* of children, especially as seen in the summer season, given with pepsin, they are almost invaluable. Topically, these preparations may be used with advantage as desiccant astringent applications in *leucorrhœa*, in *gonorrhœa*, and in *irritable external ulcers*.

ADMINISTRATION.—In order to be efficient, the bismuth preparations must be given in much larger doses than they were formerly used in. To infants five to ten grains may be administered at a dose, and to adults from twenty grains to a drachm.

Under the name of *Liquor Bismuthi et Ammonię Citratis* the British Pharmacopœia recognizes a solution of the citrate of bismuth and ammonia, which was introduced some few years since as an astringent remedy in *chronic diarrhœa*. The preparation has not, however, gained favor in this country, and has the marked disadvantage of being irritant in large doses. Moreover, much of the peculiar value of the ordinary forms of bismuth probably depends upon their insolubility enabling them to act slowly and persistently. The few trials I have made of the liquor bismuthi have convinced me that its action is different from that of the other preparations.

ZINCUM—ZINC.

ZINCI SULPHAS—SULPHATE OF ZINC.



White vitriol occurs in irregular white masses, the *pure* sulphate of zinc in minute, transparent, four-sided, prismatic crystals, which effloresce slightly in dry air, and are soluble in two and a half times their weight of cold, and much less of hot, water; insoluble in alcohol. The taste is styptic and peculiar.

THERAPEUTICS.—The sulphate of zinc is in weak solution a stimulant astringent, in concentrated form an active irritant. Taken in doses of thirty grains it acts as a prompt, efficient mechanical emetic. In smaller doses, of two grains, it is sometimes given in pills as a stimulant astringent in *chronic diarrhoea* with ulceration.

TOXICOLOGY.—Sulphate of zinc in large doses acts as an irritant poison, producing violent vomiting, colicky pains, diarrhoea, prostration, etc. The symptoms which it causes are almost identical with those produced by the corresponding salt of copper. Alkalies and their carbonates are the chemical antidotes to it, producing insoluble precipitates. Eggs and milk should also be exhibited, and the symptoms treated as they arise.

ZINCI OXIDUM VEMALE—*Commercial Oxide of Zinc.* This is a snow-white powder, which is made by burning the metal in the air. It is included in the *Materia Medica* list of the United States Pharmacopœia on account of its use in pharmacy.

ZINCI OXIDUM (ZnO).—The *pure oxide of zinc* is made by heating the carbonate until the water and acid are driven off. It is a yellowish-white powder, insoluble in water, but soluble without effervescence in dilute acids.

THERAPEUTICS.—The oxide of zinc is used externally as a mildly astringent, slightly stimulant, and desiccant application in *skin diseases* and to *ulcers*. When given continuously in small doses it is believed to act as a tonic and alterative upon the nervous system. It is especially useful in convulsive affections, and has been largely used in *epilepsy* and in *chorea*. In the latter disease it is often of service.

Unguentum Zinci Oxidi—*Ointment of the Oxide of Zinc.* (One part to five of ointment of benzoïn.) Especially used in chronic *eczema*.

ZINCI CARBONAS PRÆCIPITATA.—*Precipitated carbonate of zinc* is intended by the Pharmacopœia to replace the impure native carbonate, *calamine*. It is made by precipitating the sulphate of zinc by the carbonate of sodium. It is a white powder, closely resembling in its properties the oxide of zinc.

Ceratum Zinci Carbonatis—*Cerate of the Carbonate of Zinc.* (One to five.)

ZINCI ACETAS.—*Acetate of zinc* ($\text{ZnO}, \text{C}_4\text{H}_3\text{O}_3 - \text{ZnC}_2\text{H}_4\text{O}_2$) is made by the action of acetic acid upon the commercial oxide of zinc. It occurs in white, micaceous crystals, which effloresce in a dry atmosphere and are very soluble in water. The taste is astringent and metallic. The acetate of zinc


resembles in its physiological and therapeutic properties the sulphate, but is probably somewhat less active. It is chiefly used in collyria (one to two grains to one fluidounce), and as an injection (five to twenty grains to one fluidounce) in *gonorrhœa*.

LIQUOR ZINCI CHLORIDI—*Solution of the Chloride of Zinc.* (See *Caustics*.)

CUPRUM—COPPER. (Cu.)

Copper wire.

CUPRI SULPHAS—SULPHATE OF COPPER.



The sulphate of copper occurs in blue, transparent, slightly efflorescent, rhomboidal prisms, or their fragments. It dissolves in four parts of cold and in two of boiling water, but is insoluble in alcohol. With ammonia its solution precipitates a bluish-white hydrated protoxide of copper, which redissolves when an excess of the alkali is added, forming a rich deep-blue solution.

PHYSIOLOGICAL ACTION.—In very dilute solution the sulphate of copper acts locally as a stimulant and mild astringent; in a more concentrated form it is an irritant; in powder it is a very mild caustic, which is scarcely capable of destroying sound tissue. Taken internally in very small amounts and continuously, it is thought to have a corroborant influence upon the nervous system. In doses of five to fifteen grains it acts as an irritating emetic, and in larger amounts is an irritant poison.

THERAPEUTICS.—The chief internal use of sulphate of copper is as a mechanical emetic. As it is more irritating than sulphate of zinc, it therefore acts more rapidly and in smaller dose. For the same reason, however, it is not so safe as the white vitriol, and cannot be repeated so freely when its action fails.

As a stimulant and astringent it is occasionally administered, in pill form, in *chronic diarrhœa* with ulceration. In small repeated doses it has been used in various nervous affections with doubtful advantage.

The chief value of the so-called "blue stone" is as an external application. When applied in solid form to ulcers, it destroys flabby granulations and exerts a powerful excitant influence. Its solution acts more feebly, and is sometimes employed as a dressing for indolent *ulcers*, but more frequently as a stimulant and alterant to mucous membranes, as in *granular conjunctivitis*.

TOXICOLOGY.—The symptoms of acute copper-poisoning generally come on in about a quarter of an hour after the ingestion of the poison, but may be postponed for from one to two hours. They consist of violent vomiting and purging, accompanied by very severe colicky pains. The matters vomited are greenish or bluish, the stools glairy, mucous, and at times bloody. There is a very strong taste of copper in the mouth, and often constant expectoration. Death may occur in a few hours, preceded by convulsions, paralysis, delirium, anæsthesia, and other symptoms of great nervous disturbance. Sometimes a tendency to syncope is very marked. The urine is lessened or suppressed.

If the patient survive for twenty-four hours, jaundice nearly always shows itself. After this, death from exhaustion may occur, but not rarely a favorable issue results, in which case the symptoms of gastro-intestinal inflammation with fever develop themselves.

As the action of the sulphate of copper is exceedingly rapid, any antidote to be of avail must be given at once and act quickly. Milk and eggs are almost always at hand, and are the most efficient antidotes. No time should be lost in attempting to separate the yolk from the white of the egg, but the eggs should be broken into a bowl as quickly as possible, a little water added, and the whole stirred up and exhibited. The dose should be repeated several times, especially when there is vomiting. The yellow *prussiate of potash*, when pure, is harmless, and precipitates instantly an insoluble compound of copper from solutions of its salt. When it is to be had in time, it may therefore be used as an antidote to the sulphate. The treatment of copper-poisoning after the administration of the antidote consists in meeting the indications as they arise; opium should be used freely. When death occurs, the results of gastro-intestinal inflammation are usually found; sometimes the intestine has a decided bluish tint, and occasionally submucous ecchymoses occur. In exceptional cases, it is said, there are no evidences of inflammation in the alimentary canal.

Chronic copper-poisoning, although rare, does occur among those who work in the metal. The chief symptoms "are a coppery taste in the mouth, giddiness, pain in the bowels, vomiting, occasional diarrhoea, and wasting of the body." Dr. Clapton (*Medical Times and Gazette*, June, 1868) has pointed out as characteristic the presence of a *green* line upon the gums; this was also observed by Prof. Taylor, but its constancy is not yet assured.

CUPRI SUBACETAS ($2\text{CuO}, \text{C}_4\text{H}_3\text{O}_3 - \text{Cu}, \text{C}_2\text{H}_3\text{O}_2$).—The officinal impure *subacetate of copper*, or *verdigris*, occurs in masses of a pale-green color, which are often composed of minute silky crystals. It is sometimes used externally instead of the sulphate, which it resembles in its physiological, therapeutic, and toxicological properties.

CUPRUM AMMONIATUM.—*Ammoniated copper* is made by triturating together the sulphate of copper and the carbonate of ammonium. It occurs in deep azure-blue masses, with a strong ammoniacal odor and styptic metallic taste. It is soluble in water. In overdoses it acts as an irritant poison. In small doses (one-half to two grains three times a day) it has been used as a tonic to the nervous system in *epilepsy* and other diseases.

ARGENTUM—SILVER. (Ag.)

The United States Pharmacopœia contains among its preparations the cyanide, the nitrate, and the oxide of silver. The first of these is used solely for making hydrocyanic acid, in connection with which, in the present work, it will be treated of.

ARGENTI NITRAS—NITRATE OF SILVER. (AgONO_2 — NO_3Ag .)

This is officinally prepared by heating together silver, nitric acid, and a small quantity of water. It is a heavy anhydrous salt, crystallizing in translucent, shining, rhombic plates, and having a styptic, exceedingly metallic, corrosive taste. It is soluble in its own weight of cold water. Muriatic acid or a soluble chloride throws down from its solution a white curdy precipitate *wholly* soluble in ammonia. For external use the crystals are melted and run into moulds, where they harden into round, grayish, brittle sticks, about the size of a goose-quill, and having a radiated crystalline fracture. These are the officinal *argenti nitras fusa*. As only the pure salt will make well-formed crystals, the impure products are always manufactured into the preparation just named, which should therefore not be employed internally.

When the nitrate of silver, either in substance or solution, is exposed to the conjoint influence of light and of even a minute portion of organic matter, it turns black, and is converted into an insoluble substance, which has been believed to be metallic silver, but is more probably an oxide. For this reason the white stains which it first makes when applied to living tissues soon become black.

PHYSIOLOGICAL ACTION.—Nitrate of silver coagulates albumen, and, when applied in its pure state to living tissues, acts as a caustic, coating them over with a white almost membranous film. The caustic action is, however, not a deep one, because penetration of the salt into the tissues is soon prevented by the thick and tough skin or stratum which is formed. When applied in a dilute solution it acts as an astringent, constricting the vessels and overcoming relaxation. Its local action, however, is not simply that of an astringent, but is certainly peculiar and apparently alterative to the nutritive processes.

When taken internally in sufficient dose, this salt, by virtue of its corrosive action, is a poison, producing gastro-enteritis; but it also acts directly upon the nervous system. Orfila and other of the earlier observers experimented upon it by injecting it directly into the veins of animals. When exhibited in this way, it must, by coagulating the albumen of the blood, produce thrombi, to which the subsequent symptoms are in greater or less measure to be ascribed. This method of experimentation can therefore throw but little light upon the action of nitrate of silver when taken into the stomach.

It is evident that in the stomach the nitrate of silver cannot long maintain its integrity. Dr. Bogolowsky has found (*Virchow's Archiv*, Bd. xlv. p. 413) that when the nitrate is added to a peptone it is readily dissolved, and that the solution formed does not coagulate albumen. In this or some other analogous form silver is undoubtedly absorbed: that it is absorbed is proven by its having been found in various internal organs and by the discoloration which follows its protracted use. When it is exhibited for a long continuous period, the skin often acquires a peculiar bluish slate color, which may

become very dark, and in decided cases the conjunctiva and even the mucous membrane of the mouth are involved. The dark color is undoubtedly due to the presence of silver in the skin.

Both Heller and Orfila failed to detect silver in the urine of animals taking it; but probably it is eliminated, though slowly and in very small quantities, by the kidneys.

By an elaborate series of experiments, M. Chas. Rouget (*Archives de Physiologie*, July, 1873, p. 356) has shown that upon all animals from a crab to a dog the soluble salts of silver act as a poison, causing in mammals vomiting and purging, and in them and the lower animals violent disturbance of the motor functions, as shown by paralysis and convulsions, and of the respiration, ending finally in death by asphyxia. This is in accord with the observations of other investigators. MM. Rabuteau and Mourier have found that the almost instantaneous death which Chareot and Ball first noted as following the injection of a large dose of the nitrate of silver into the veins is due to a direct paralyzing influence of the drug upon the muscle of the heart. M. Rouget has never seen this form of death follow the hypodermic or internal administration of the poison, the heart always continuing to beat for a greater or less length of time after the cessation of respiration, and also retaining its irritability.

As already stated, both convulsions and paralysis are present in *argyria*, or silver-poisoning. The convulsions are severe, generally tetanic, and seem to resemble very closely those caused by strychnia, since Rouget states that in the frog they are evidently reflex, excited by the least peripheral irritation. A peculiarity which that observer notes is their persistence after the complete abolition of voluntary movements.

The death is due, in *argyria*, to cessation of the respiration; Rouget (*loc. cit.*, p. 351) even states that he has witnessed the suspension of the latter function in the frog whilst the activity of the reflex movements was much beyond normal. In the dog and in the full-grown cat this asphyxia is accompanied by an outpouring of mucus in the lungs, pulmonary congestion and œdema being found on post-mortem examination.

Two theories have been propounded as to the cause of the asphyxia: one, that it is simply due to the choking up of the lungs by the congestion and the excessive secretion whose origin is an altered state of the blood; a second, that both the asphyxia and the lesions in the lungs have their origin in a direct action of the poison upon the nerve-centres.

The first view has been especially supported by Krahmer and by MM. Rabuteau and Mourier. Unfortunately, I have not seen the original papers of these physicians; but, according to Rouget, the basis of argument of Krahmer is simply the ecchymoses which he found in horses dead of the poison, whilst that of Rabuteau and Mourier is the fluidity of the blood after death, and the existence in it of globules which, on account of their solubility in ammonia, were believed to be the chloride of silver. The French observers were, how-

ever, almost certainly mistaken in their belief that these granules were chloride of silver, since ammonia dissolves hæmatin as freely as it does the chloride named.

In 1864 Charcot and Ball (*Gazette Médicale*, 1864) made a series of experiments in which a silver salt that did not coagulate albumen was injected directly into the blood. They noted not only the respiratory embarrassment, but also that the hinder extremities were suddenly paralyzed, and concluded that both the asphyxia and the lung-trouble were due to an affection of the central nervous system.

In 1869 Dr. Bogolowsky, of Moscow, studied (*Virchow's Archiv*, 1869, Bd. xvi.) the action of a peptone of the nitrate when used hypodermically. He found, on examination of the blood of a poisoned animal, that the spectrum analysis (*loc. cit.*, p. 415) betrayed nothing abnormal; that the red corpuscles appeared paler and their outline more delicate than normal; that the white corpuscles were natural. On the other hand, Rouget (*loc. cit.*, p. 361) examined microscopically the blood of animals poisoned with the nitrate of silver, and found it perfectly normal. The only conclusion to be drawn from all this seems to me to be that at present there is no proof whatever that the symptoms of acute argyria are due to alterations in the blood.

That the embarrassment of respiration is not due to local lesions in the lungs is abundantly shown by the experiments of Rouget, who found that whilst in all animals these respiratory symptoms are very prominent, in only a few species are decided pulmonic lesions found after death. From all these facts I think it highly probable, if not altogether certain, that the theory propounded by MM. Charcot and Ball is correct.

That the motor disturbance is centric, not peripheral, in its origin, is shown by the fact noted by Rouget (*loc. cit.*, p. 354), that the muscles and nerves preserve their excitability after the arrest of the respiration.

The various facts which have been thus far brought forward in regard to the physiological action of silver, although interesting to the toxicologist, have very little reference to its therapeutic use, since it is never employed to produce an acute constitutional influence.

The action of the drug when exhibited continuously for a length of time in large doses has been investigated by Dr. Bogolowsky upon dogs and rabbits. He found that it produced loss of appetite, wasting, slight lowering of bodily temperature, diarrhoea, diminution of the quantity of urine passed, with increase of its specific gravity and often with the presence of albumen, and transitory paralyses. How far some of these symptoms were due to the direct constitutional action of the poison, and how far to derangement of the digestion dependent upon its local influence, is perhaps an open question. The local action was avoided, however, as much as possible, by the use of an albuminate or of the double phosphate of silver and soda, which does not coagulate albumen. Comparative examinations of the blood showed that the hæmoglobin was reduced by more than one-third. The blood was also rendered very

aplastic, as was betrayed by the constant tendency to the formation of ecchymoses. As some one has suggested that the silver in these cases replaces the iron of the blood-corpuscles, Dr. Bogolowsky made a chemical examination of the latter, but failed to find any traces of silver in them,—no doubt because it was not there.

The solid tissues were found, after death from chronic argyria, to be in an advanced stage of degeneration, which especially affected epithelial structures. The first change was swelling and opacity of the cells, with obscuration of the nucleus. After this came fatty degeneration, fatty globules in the cell, destruction of nucleus, and finally of the cell itself. The liver and kidneys were profoundly influenced, as was also the muscular structure, especially of the heart.

The summary which has been here offered comprises all our knowledge of the physiological action of the preparations of silver. Unfortunately, it does not throw much light upon their therapeutic use. The results of the chronic poisoning are so closely analogous to those produced under similar circumstances by antimony, arsenic, and probably other metallic poisons, as to indicate that silver given internally acts upon the nutrition of the body,—in other words, that it is an *alterative*.

THERAPEUTICS.—By far the most frequent employment of nitrate of silver in therapeutics is for its local action, either upon the surface of the body or upon those mucous membranes that can be reached directly by the drug.

As a simple *caustic*, the salt may be used whenever only a superficial action is required; for reasons already given (page 42), it is useless whenever it is necessary to produce a deep eschar.

As a *caustic* and an *alterative*, it is applied in solid form to many *ulcerated surfaces*, for the purpose of destroying superficial diseased tissue and of substituting, when the eschar separates, a healthy for an unhealthy action.

As an *antiphlogistic*, nitrate of silver acts not only as an astringent, but also in some way not clearly understood. In the various inflammations of the mucous membranes, such as *conjunctivitis*, *faucitis*, *laryngitis*, *urethritis*, etc., it is employed very frequently, not only in the stage of relaxation, but in the beginning of the attack. Its influence in the former case is more marked.

In *conjunctivitis*, the solution employed should not, under ordinary circumstances, be stronger than one or two grains to the ounce; and it should not be used at all if any corneal ulceration exists, since a deposit of silver is liable to occur and to produce opacity.

In *faucitis*, the strength of the solution may vary from fifteen to thirty grains to the fluidounce. A solution of forty grains to the ounce may be looked upon as caustic to the mucous membranes, and should be only used as such. In ordinary cases of *sore throat*, the application once a day or every alternate day is generally sufficient. It is best made by means of a good-sized camel's-hair brush, each part of the inflamed surface being distinctly touched, and not the whole simply daubed or slopped over by means of a very

large brush or a sponge probang, as is often done. In severe cases it may be necessary to use the solution twice a day. In *laryngitis*, the solution may contain from ten to twenty grains to the ounce, and should be applied with a brush by the aid of the laryngoscopic mirror.

An attack of *urethritis* may sometimes be aborted in its forming stage by the injection of a strong solution (grs. xii to f $\overline{3}$ i) of the salt; but the practice is of doubtful expediency, since when it fails it greatly aggravates the trouble. In the *advanced stages of gonorrhœa*, weak injections (grs. i or ii to f $\overline{3}$ i) are often very serviceable.

Many years ago (1828) Mr. John Higginbottom originated the practice of treating *erysipelas* by the nitrate of silver, and his plan has received a great deal of commendation from authorities, but certainly has not been generally adopted by the profession. One or two cases of bad results, from ulceration of the skin apparently due to the local application, have deterred me from giving the method a fair trial, and I do not feel able to offer any opinion upon the practice. Mr. Higginbottom in a recent essay (*London Practitioner*, vol. ii. p. 34, 1869) reaffirms the value of the treatment, stating that its want of general adoption is due to its being so often imperfectly carried out, and gives the following directions:

“The affected part should be well washed with soap and water, then with water alone to remove every particle of soap, then to be wiped dry with a soft towel. The concentrated solution of four scruples of the nitrate of silver to four drachms of distilled water is then to be applied two or three times on the inflamed surface, and beyond it on the healthy skin to the extent of two or three inches. The solution may be applied with a small piece of clean linen, attached to the end of a short stick, the linen to be renewed at every subsequent application. As the solution of the nitrate of silver is colorless, it is necessary to pass a little linen, just moistened, over every part where it has been used, in order to be equally diffused, so that no part may be left untouched. In about twelve hours it will be seen whether the solution has been well applied. If any inflamed part be unaffected, the solution must be immediately reapplied. Sometimes, even after the most decided application of the nitrate of silver, the inflammation may spread; but it is then generally much less severe, and is eventually checked by repeated applications. It is desirable to visit the patient every twelve hours until the inflammation is subdued.”

In *superficial inflammations* other than erysipelatous I have frequently used nitrate of silver in this way, often with great advantage. Freely applied to the skin of the whole finger, it will sometimes even abort a commencing *felon*.

Internally, the nitrate of silver is exceedingly useful in stomachic and to a less extent in enteric diseases, exerting no doubt a purely local influence. In that form of *dyspepsia* characterized by the vomiting of large quantities of yeasty fluid, it has yielded in my hands better results than any other remedy; and the same may be said of *chronic gastritis* and of *gastric ulcer*.

The rules of administration are identical in these three diseases. In the first place, regulation of the diet is imperative: if the case be a bad one, all eating of meals should be suspended, and the patient receive every two or three hours a cup of sweet milk, with sound toasted bread broken up and thoroughly softened in it. The medicine should be administered in pill form, one-quarter to one-half grain three or four times a day, taken when the stomach is empty. In very serious cases, when all food is rejected by the stomach, it is sometimes advisable to allow absolute rest for two or three days to that viscus, the patient being fed by the rectum, and only a little water and pills of silver with opium being taken by the mouth. Under these circumstances, the return to the usual method of taking food must be very gradual, at first only a tablespoonful each of milk and of lime-water being administered every hour. In *chronic enteritis* or *colitis*, nitrate of silver is sometimes of great service, especially if there be ulceration.

For its constitutional effects nitrate of silver is used solely in diseases of the nervous system. It cannot be denied that occasionally in *epilepsy* it has achieved brilliant results, but the successes are certainly far less numerous than the failures. There is one serious objection to the employment of the salt,—namely, the discoloration of the skin which sometimes follows its continuous use. Again, it is not understood in what class of cases the drug is of especial value, and there is no means of judging as to its applicability to any individual case. When other means have failed, however, the nitrate of silver may be tried in epilepsy, the patient or his friends being informed that although with proper precautions the chances of discoloration are very few, yet it may occur.

In *chronic inflammations* of the spinal cord, whether affecting chiefly the posterior columns and constituting *locomotor ataxia*, or the anterior and giving rise to *paraplegia*, the nitrate of silver is one of the few remedies that are ever of any service; although it most frequently fails, yet it often does good, and in some cases has apparently even permanently arrested the disease.

TOXICOLOGY.—The symptoms produced by the ingestion of large doses of nitrate of silver are partly gastro-intestinal and partly cerebro-spinal. In some cases the one series of phenomena predominate; in others, those of the other class. In a case at the Hôpital St.-Louis in 1839 (*Beck's Medical Jurisprudence*, vol. i. p. 675, Phila., 1863) the symptoms were insensibility, violent convulsions, and dilated pupils, with, on a partial return to consciousness, intense gastric pain; complete restoration of consciousness did not occur until eleven hours after admission into the hospital, and the coma returned at intervals during several days.

Vertigo, coma, convulsions, great muscular weakness, paralysis, with intense disturbance of respiration, are in these cases the manifestations of disturbed innervation; whilst the abdominal symptoms are those of gastro-enteritis. The diagnosis can generally be made by the discolorations of the lips and skin,—at first white, afterwards black,—and by the blackish or brownish vomit; when the customary antidote has been given, both vomit and stools are generally

white and curdy. At post-mortem the stomach and bowels are found corroded, often ecchymosed and with patches of a white or grayish color. Poisoning by nitrate of silver is not common, and I know of but three fatal cases,—one in 1837 (Taylor, *Principles and Practice of Medical Jurisprudence*, second edition, vol. i. p. 319), one in 1861, a woman killed by fifty grains in solution in divided doses, one in 1871, a child destroyed by a piece of the solid stick three-quarters of an inch long, in spite of the immediate and free administration of the antidote (Scattergood, *British Medical Journal*, May, 1871).

The treatment consists in the administration at once of large amounts of common salt,—the chemical antidote,—the constant use of large draughts of milk, and the meeting of symptoms as they arise.

The fatal dose of silver varies very much, according, no doubt, as to the presence of substances capable of decomposing it in the stomach. Thirty grains have killed; and recovery has taken place after the ingestion of an ounce (Poumarède's case, Husemann, *Handbuch der Toxicologie*, Berlin, 1862, p. 868).

Chronic *argyria*, or discoloration of the skin by silver, is unaccompanied by any disturbance of health, although in severe cases the discoloration affects not only the skin, lips, gums, sclerotic, but even the internal organs, such as the liver, spleen, kidneys. It is therefore not due, as has been thought, to the chloride of silver, since the latter only becomes dark under the influence of the light, but to a deposition of silver itself or of its oxide. The minute quantity of the metal present is shown by the analysis of Versmanns (*Virchow's Archiv*, xvii., 1859), who in 14.1 grammes of dried liver found only 0.0068 gramme of metallic silver (0.047 per cent.), and in 8.6 grammes of dried kidney 0.053 gramme (0.061 per cent.). Greater or less success has been claimed for various treatments in argyria, but in general they are equally futile. Rogers states that blistering will lighten the color very much, and Eichmann asserts (Husemann, *Toxicologie*, 871) that he has cured two cases by the use of potash baths and of soap baths, each four times a week. The older authorities commend the use of iodide of potassium internally. Dr. L. P. Yandell has reported (*American Practitioner*, June, 1872) two cases in which large doses of the iodide were given for many months for syphilis, and the mercurial vapor-baths used at the same time for the same purpose, with the result of a complete cure of the argyria. The fading was gradual.

ADMINISTRATION.—The nitrate of silver should always be given in pill, and, when it is desired to obtain its constitutional influence, after meals, during the process of digestion; but when its local action on the alimentary canal is required, it should be administered one or two hours before meals; and if the bowels are to be reached, the pill should have been made some time, so as to be dissolved as slowly as possible. When it is given in epilepsy or other chronic disease, its administration should be suspended for one week at the end of every third week, and its employment should not extend over a longer time than three months without a protracted intermission.

The *oxide of silver* (ARGENTI OXIDUM) is an olive-brown powder, very slightly soluble in water, which the United States Pharmacopœia directs shall be prepared by precipitating the nitrate of silver with solution of potassa. It has been introduced into medicine as a substitute for the nitrate, with the idea that it would accomplish in diseases of the nervous system all that that drug is capable of, and at the same time not discolor the skin. With our present knowledge of the method of absorption of the nitrate, this seems highly improbable, and the reason that no case of discoloration by the oxide has occurred, without doubt, is simply its infrequent use. Oxide of silver is not caustic when locally applied, but probably exerts some astringent action, and has been commended in *pyrosis*. In nervous affections it is probably of equal value with the nitrate. The dose is a grain, given in pill three or four times a day.

CLASS II.—TONICS.

THERE are certain substances in nature which, when taken internally, act upon the nutrition of the various tissues so as to restore lost tone, not by calling into play the vital principle of contractility, but by increasing the power in the part. Such substances are known as tonics. They differ from astringents in that they affect nutrition, and consequently in the slowness and permanence of their action. They differ in a similar manner from simple stimulants, and as they do not call into sudden action forces already existent in the part, but increase power by increasing nutrition, their influence is a permanent one and is not followed by depression.

They are, of course, indicated by debility dependent upon impaired nutrition,—*i.e.*, debility owing to actual loss of power. When the debility is due to a sudden depressing influence, as in snake-bite, they are of no service whatever. They are especially valuable in convalescence from acute disease; during the progress of the disorder they more often do harm than good.

Tonics should, of course, never be used when plethora exists. Sthenic inflammatory action is also a contra-indication to their employment; but when inflammation exists with a general state of debility, these drugs may form an essential part of the treatment.

The tonics are divisible into several sub-classes, which I shall consider separately.

SIMPLE BITTERS.

Probably all bitter vegetable substances possess tonic properties, but in many of them, as in morphia and strychnia, these properties are completely overshadowed by other inherent powers. There are, however, bitter vegetable substances which so act upon the stomach as to invigorate digestion and to affect thereby the general nutrition, without exerting any direct influence upon other portions of the body than the alimentary canal. These are the simple bitters. In overdoses they nauseate, and may act slightly on the bowels. They appear all to act alike,—differing more in strength than in quality,—so that one may be substituted without detriment for another. As they are essentially irritants, inflammation or over-sensitiveness of the alimentary mucous membrane distinctly contra-indicates their administration. They are indicated by loss of appetite and loss of stomachic tone.

QUASSIA.

The wood of *Simaruba excelsa*, a large tree, native of Jamaica. This wood is light both in density and color, somewhat resembling that of the tulip-tree, but distinguished by its intensely bitter taste. It is kept in the shops in billets and in raspings. The active principle of it appears to be *Quassin*, an intensely bitter, neutral, crystalline principle discovered by Winckler (*Repert. für Pharmacie*, Bd. liv. 85, Bd. lv. 85). *Simaruba* (U. S. P.), the bark of *Simaruba officinalis*, also contains quassin, and may be substituted for quassia.

PHYSIOLOGICAL ACTION.—*Quassin* is said in large doses to be actively poisonous to insects and even to mammals (Stillé's *Therapeutics*, i. p. 472; Husemann, *Die Pflanzenstoffe*, p. 718), but no detailed study of its action has as yet been made.

THERAPEUTICS.—Quassia is probably the most active of all the simple bitters, and may be used whenever such remedies are indicated. In cases of *seat-worms* in children, a strong infusion of quassia (℥ii to Oj) affords a most harmless and efficient injection. Its exhibition should be preceded by an enema of simple water, after a stool, so as thoroughly to wash out the rectum and allow access to every fold of the rectal mucous membrane.

Infusum Quassiae—*Infusion of Quassia* (℥ii to Oj). Dose, one to two fluidounces.

Extractum Quassiae—*Extract of Quassia*.

This watery extract is an excellent preparation. Dose, one to two grains.

Tinctura Quassiae—*Tincture of Quassia* (℥i to Oj). Dose, twenty drops to a teaspoonful.

GENTIANA—GENTIAN.

The root of the *Gentiana lutea* or yellow gentian of the Alps. This root occurs in the shops either in pieces of various size and shape but generally several inches in length, or else in transverse slices. The texture is spongy, the odor faint but peculiar, and the taste bitter. It contains *gentisic acid*, which was discovered by Leconte and is tasteless and physiologically inert. The active principle is probably the *gentiopikrin* of Kromayer, a neutral, crystalline substance, of an intensely bitter taste.

THERAPEUTICS.—Gentian is one of the most efficient of the simple bitters, and may be used whenever such a remedy is indicated. It is never given in substance, but in one of the following preparations:

Infusum Gentianæ Compositum—*Compound Infusion of Gentian*. This contains, besides gentian (℥ss to Oj), bitter-orange peel, coriander, and alcohol, and is a very excellent stomachic bitter. Dose, one to two fluidounces.

Tinctura Gentianæ Composita—*Compound Tincture of Gentian*. This contains gentian (℥i to Oj), bitter-orange peel, and cardamom. Dose, one fluidrachm to half a fluidounce.

Extractum Gentianæ—*Extract of Gentian (watery)*. Dose, two to four grains.

Extractum Gentianæ Fluidum—*Fluid Extract of Gentian*. Dose, ten minims to half a fluidrachm.

BERBERIS—BARBERRY.

Under the above title is included in the secondary list of the United States Pharmacopœia the bark of the root of *Berberis vulgaris*, a native of Europe. This drug contains the alkaloid berberina, which, although not itself official, is found in so many recognized drugs as to require notice.

Berberina occurs as a yellow powder, or in fine yellow needles or prisms, of a bitter taste and neutral reaction, moderately soluble in cold, freely in hot, water, freely soluble in alcohol. With acids it forms mostly crystallizable, golden-yellow salts.

Its effects upon animals have been studied by Falck and by Guenste. In dogs it produced, when given in very large doses, restlessness, convulsive tremblings, thirst, and diarrhœa, and finally partial paralysis of the hind legs. From seven to fifteen grains of it killed rabbits in from eight to forty hours. The symptoms were increased frequency of respiration, and tremblings, followed by decrease of the rate of breathing, paresis, paralysis of the hind legs, great dyspnœa, and finally convulsions. In man, as yet, no serious symptoms have been recorded as produced by berberina. Buchner took nearly twenty grains without causing anything more serious than a loose stool. As a bitter tonic it has been used by various physicians in doses of from two to five grains, and its action in this dose appears to be that of a simple bitter. It may be given in pill form or dissolved in alcohol.

CALUMBA—COLUMBO.

The root of *Cocculus palmatus*, a climbing vine of Mozambique. It occurs in the shops in transverse disk-like slices, oval or circular in outline, one or two inches in diameter, of a spongy texture, having a yellowish surface, very bitter taste, and slightly aromatic odor. It contains a great deal of starch, besides berberina, and, it is said, in lesser amount, *columbin*, a bitter neutral principle crystallizing in rhomboid prisms or needles.

THERAPEUTICS.—A bitter, slightly aromatic tonic, useful as a stomachic in cases in which a simple bitter is indicated. It is not used in substance, but in one of the following preparations:

Infusum Calumbæ—*Infusion of Columbo* (℥ss to Oj). Dose, one to two fluidounces.

Tinctura Calumbæ—*Tincture of Columbo* (℥ii to Oj). Dose, one to two fluidrachms.

Extractum Calumbæ Fluidum—*Fluid Extract of Columbo*. Dose, fifteen minims to half a fluidrachm.

EUPATORIUM—BONESET.

The herbal parts, gathered after flowering, of the indigenous *Eupatorium perfoliatum*, a tall, coarse composite, recognizable by its perfoliate leaves,

through whose centre grow the stems and branches. This drug given in cold infusion (℥i to Oj—dose, f ℥i-ii) is an efficient but disagreeable bitter tonic. Its chief employment is as a sudorific. The hot infusion when taken freely (five to six ounces), the patient being well covered in bed, produces free sweating, and has been very largely used in “general colds,” *muscular rheumatism*, etc., for this purpose. The only objection to the remedy is its disagreeable taste. In the dose of a pint the infusion has been employed as an emetic.

The following simple bitters, the products of native plants, are also recognized by the United States Pharmacopœia, and are occasionally used:

COPTIS, or *Goldthread*, the yellow, fine, thread-like roots of the *Coptis trifolia*, an elegant bitter. *SABBATIA*, or *American Centaury*, the herb of *Sabbatia angularis*. *FRASERA*, or *American Columbo*, the root of the *Frasera Walteri*. The bark of the *CORNUS SERICEA*, or *Swamp Dogwood*, of the *CORNUS CIRCINATA*, or *Round-leaved Dogwood*, and of the *CORNUS FLORIDA*, or *Dogwood*, are all included in the materia medica list. They all possess feeble tonic properties, and the latter has been thought to have some antiperiodic power.

The neutral crystalline principle of the willow, *Salicin*, is probably a simple bitter, although antiperiodic properties have been attributed to it. Its influence upon the system must be very slight, since Ranke (Husemann, *Die Pflanzenstoffe*, p. 963) took nearly three ounces of it in as many days without the induction of any notable symptom.

PECULIAR BITTERS.

These are bitter vegetable tonics possessing other properties besides those of the simple bitter.

PRUNUS VIRGINIANA—WILD CHERRY BARK.

Wild cherry bark is the product of the *Prunus* (*Cerasus*) *serotina* or wild cherry tree, not of *Prunus Virginiana* or Choke cherry, whose name it bears. It occurs in pieces of various sizes, usually without epidermis. The color is a reddish cinnamon; the taste slightly astringent, bitter and peculiar, resembling that of peach-leaves. It contains tannic acid, bitter extractive, amygdalin, and emulsin.

Amygdalin is a nitrogenous, crystallizable, odorless glucoside, of a slightly bitter taste. It is soluble in water and alcohol, but not in ether.

Emulsin is an albuminous principle, which is soluble in water, and, like other forms of albumen, is coagulated by heat, alcohol, acids, etc.

When amygdalin in watery solution is brought in contact with emulsin, it is decomposed, forming prussic and formic acids and a colorless, thin, volatile oil, which, when pure, has a peculiar agreeable odor and a burning taste. According to Liebig and Wöhler (*Ann. Chem. Pharm.*, xxii. 1), seventeen

grains of amygdalin yield one of hydrocyanic acid: therefore, if thirty-four grains of amygdalin be mixed with sixty-six grains of emulsion of sweet almonds, a two per cent. solution of hydrocyanic acid will be formed.

PHYSIOLOGICAL ACTION.—Amygdalin administered by itself is nearly, if not quite, without effect upon the organism. Widtmann and Denk (Husemann, *Die Pflanzenstoffe*, p. 688) took as much as sixty grains of it without inducing any effect, and their results have been confirmed by Reil and others. Lehmann, it is true (*ibid.*), found that at times fifteen grains of it by the mouth were sufficient to cause death in the rabbit, but Köl liker and Müller have shown that this was owing to its being converted into prussic acid by the emulsin contained in the green herbage in the stomach of the rabbit.

THERAPEUTICS.—In wild cherry bark properly administered there are three active ingredients: tannic acid, bitter extractive, and prussic acid; and to their combined action the general effect is due.

As the tannic acid is in small quantity, its influence is not marked: but probably some of the reputation which the remedy formerly enjoyed as being useful in the night-sweats of phthisis was due to it. When given in the ordinary dose, I have never been able to detect any sedative effect from wild cherry bark; but the infusion administered with sufficient freedom certainly ought to exert such influence. *Prunus Virginiana* is therefore slightly astringent, sedative, and, owing to its bitter extractive, tonic.

It is frequently useful in *phthisis* when a roborant is needed and at the same time a calmative action on the nervous system is called for. It may be used in *hectic fever* from any source.

The infusion is also useful as an adjuvant to other tonics, especially sulphuric acid, in *debility*, with a tendency to night-sweats during convalescence from acute disease.

On account of its pleasant taste and traditional reputation, the syrup of wild cherry is often employed as the basis of cough-mixtures.

Infusum Pruni Virginianæ—*Infusion of Wild Cherry Bark* (℥ss to Oj). Dose, one to two wineglassfuls.

Extractum Pruni Virginianæ Fluidum—*Fluid Extract of Wild Cherry Bark*. Dose, half a fluidrachm to a fluidrachm.

Syrupus Pruni Virginianæ—*Syrup of Wild Cherry Bark*. Dose, half a fluidounce.

CINCHONA.

Under the general term of *Cinchona* the United States Pharmacopœia recognizes the “bark of all the species of the genus *Cinchona* containing at least two per cent. of the alkaloids which yield crystallizable salts.” Under the name of *CINCHONA FLAVA*, or *YELLOW CINCHONA* (syn. *Calisaya Bark*), the same standard list includes the bark of *Cinchona Calisaya*; under the name of *CINCHONA PALLIDA*, *PALE CINCHONA*, that of *Cinchona Condaminea* and *C. micrantha*; under the title of *CINCHONA RUBRA*, *RED CINCHONA*, that of *C. succirubra*.

All the trees yielding quinia and its congeneric alkaloids belong to the genus *Cinchona*, and inhabit the slopes of the Andes Mountains from the northern extremity of South America to the lower portion of the republic of Bolivia, at an altitude of from five to ten thousand feet. The bark is collected by persons known as *Cascarilleros*, and is exported in large bundles or bales, usually surrounded with raw ox-hide. To these packages the term *seroons* is applied.

The *Pale Bark* occurs in quills or cylindrical rolled pieces, with a grayish or brownish adherent epidermis. It derives its name from the light fawn-color of its powder.

The *Calisaya* or *Royal Yellow Bark* comes both in quills and flat pieces. The epidermis is absent, or, if present, is loosely attached. The powder is of a decidedly yellow color, sometimes almost orange. The fracture is short and fibrous, presenting semi-transparent, sharp spiculæ, which are apt to become detached and penetrate the fingers.

The *Cinchona rubra*, or *Red Cinchona*, occurs in quills and flat pieces. The epidermis is generally present, and is often very warty. The color of the powder is a faint reddish-brown. Beneath the epidermis is in most specimens a distinct, red, resinous layer, which is very apparent in cross-section.

Large quantities of bark under the name of *Carthagena Bark* find their way into commerce through the northern ports of South America. For descriptions of these the reader is referred to works on *Materia Medica*.

Very successful attempts are being made to cultivate the cinchona-trees, especially in the Himalaya Mountains, where many millions of them have been planted and are thriving. As yet the East India bark has not been put into commerce, but in a few years the supply from this source will doubtless be very large.

CHEMICAL CONSTITUTION.—The principal constituents of cinchona are volatile oils, *quinia*, *quinidia*, *cinchonina*, *cinchonidia*, *aricia*, *tannic*, *kinic*, and *kinovic acids*, and *cinchona red*. The alkaloids exist in combination with the acids. Quinia and cinchonina only are officinal.

QUINIA.

This alkaloid was first distinctly separated from the other ingredients of the bark by Pelletier and Caventou in 1820. When quinia is precipitated by an alkali from a solution of its salt, it usually falls as a hydrate, which may be crystalline. By sufficient heat the hydrate is melted and the water is driven off. On cooling, the alkaloid, now free from water, forms a white, opaque, crystalline mass.

QUININÆ SULPHAS—SULPHATE OF QUINIA.

The neutral, officinal sulphate of quinia occurs in light silky crystals, soluble in seven hundred and forty parts of cold or in thirty of boiling water, readily soluble in alcohol, very freely so in acidulated solutions, insoluble in

ether. The aqueous solution, upon the addition of chlorine and afterwards of ammonia, assumes a green color. According to Dr. G. Kerner, if the original solution be colorless and clear and in a clean test-tube, one-thirty-thousandth part of the alkaloid can be recognized by this chlorine-ammonia test. When to one hundred parts of the salt, dissolved in nineteen hundred and twenty parts of cold dilute acetic acid (sp. gr. 1.042), are added successively four hundred and eighty parts of alcohol (sp. gr. 0.837) and sixty parts of a saturated alcoholic solution of iodine, crystals of *Herapathite* separate, in the form of right-angled quadrate rhombic leaves, which when seen by transmitted light are olive-green, but in reflected light are bright metallic green, resembling the elytra of Spanish flies. When heated, the sulphate of quinia becomes phosphorescent, emitting a pale-green light at 155° to 160° C., and at a higher temperature melts with the development of purplish vapors. When the neutral sulphate of quinia is dissolved in water acidulated with sulphuric acid, it is converted into the soluble bisulphate, which may be obtained in orthorhombic prisms by evaporation.

PHYSIOLOGICAL ACTION.—Quinine or its salts in powder or solution are, when applied upon a part denuded of its epidermis, very active irritants. Upon the skin they have little or no influence, but upon the mucous membranes they exert a very perceptible stimulant or irritant action.

A. Eulenburg found that when quinia was brought into contact with a nerve it did not cause contraction in the tributary muscles, but when placed upon the muscles themselves it induced immediate violent action. He therefore concludes that it is not a nerve-irritant, but is a muscle-irritant. When it is administered to dogs in sufficient quantity, it produces restlessness, followed by muscular tremblings, which have been compared to those of paralysis agitans, loss of power deepening into more or less complete paralysis, great dyspnoea, and cerebral symptoms, such as blindness, stupor, or violent delirium, dilated pupils, coma, and convulsions. When the drug is introduced by the stomach, vomiting generally occurs, and at times diarrhoea also.*

The first symptoms of cinchonism, as produced by small therapeutic doses (ten grains) in man, are usually ringing in the ears, slight fullness in the head, and perhaps some deafness. With the use of larger doses these symptoms are intensified: the deafness is very marked, disturbed vision† may exist, and the flushed face, with the sense of distention in the head, may point towards a cerebral congestion, which is in some cases relieved by spontaneous epistaxis. In decided cinchonism, giddiness and staggering in walking are very common. After toxic doses, severe headache, delirium, stupor, complete deafness and blindness, dilated pupils, embarrassment of respiration, great weakness, convulsions, paralysis, and finally collapse, may result.

* See M. F. Melier, *Mémoires de l'Académie*, t. xii, p. 722, 1843; Wm. O. Baldwin, *American Journal of the Medical Sciences*, April, 1847; P. Briquet, *Traité thérapeutique de Quinquina*, Paris, 1855.

† I have seen complete amaurosis produced in a lady by twelve grains of quinine.

The minimum fatal dose of quinine is not known, but it must be large, and probably varies very much. In the famous case of Bazire two ounces caused his death, whilst his wife barely recovered from the effect of one ounce.

A close physiological study of quinia can best be made by investigating its effects upon the different systems of organs *seriatim*, and this I shall now do.

According to the experiments of Briquet, a solution of sulphate of quinia injected into the carotid will in some cases produce meningitis. In doing this, it is evident, the salt acts rather as an irritant to the membranes of the brain than as a nervous stimulant: indeed, experimental evidence proving that quinia is a cerebral stimulant seems to me to be wanting. The chief proof that the alkaloid does act in such manner lies in the fact that persons who have been taking it regularly for some time will occasionally, upon the sudden withdrawal of their daily dose, manifestly be less active without than with it. Briquet may be right in his belief that in small doses it acts as a nervous stimulant, but the proof of his correctness at present is clinical rather than experimental.

In sufficiently large doses quinia without doubt abolishes the functions of the cerebrum.

The action of the alkaloid upon the spinal centres and nerves has been especially studied by A. Eulenburg (*Reichert's Archiv für Anatomie*, 1865). According to his experiments, the drug in poisonous doses abolishes all reflex actions before voluntary motion, since poisoned frogs, in which no movement could be excited by mechanical or chemical irritations of the skin, would immediately turn into their normal position when laid on the back. That these movements were voluntary was shown by the fact that section of the cord high up prevented them. According to the experiments of Dr. T. A. Chaperon, however (*Pflüger's Archiv für Physiologie*, 1869, p. 295), this lessening of reflex action is due not to a direct influence of the quinia upon the cord, but to a stimulation of the inhibitory reflex centres which Setschenow has proven to exist in the cerebrum of the frog. Thus, when the reflex activity in a frog poisoned with quinia had almost disappeared, section of the cord was instantly followed by the return of the normal activity in the parts below the point of division; again, if the cord were divided before the administration of quinia, no depression of the reflex activity below the point of section occurred. This result certainly accords with the experimental observation of Eulenburg that in quinia-poisoning no stage of excitement of reflex activity precedes that of the depression of function. According to Chaperon, quinia exerts but little influence upon the nerve-trunks.

Upon the stomach and intestines quinia acts very much as a simple bitter. In moderate doses it stimulates digestion and increases the appetite; in large doses it not unfrequently causes nausea and vomiting. When there is any morbid irritability of the mucous membrane of the stomach or bowels, its irritant action is often very marked, and its continued use in large doses has been known to cause gastritis.

Many years since, M. Piorry asserted that a large dose of quinia would produce a distinct immediate lessening of the size of the spleen in cases of intermittent; but the testimony of very numerous observers to the contrary is so concurrent as to render the truth of his observation highly improbable. Although observers* are not wanting who assert that the exposed spleen of an animal can be seen to contract when quinia is injected into the veins, yet the evidence seems to me insufficient to prove that the alkaloid really has any such action.

The action of quinia upon the *circulation* has been studied most closely by Briquet, who found that large doses of the alkaloid injected into the stomach, cellular tissue, or veins of dogs produced decided lowering of the arterial pressure. He also found that when a sufficient amount was thrown into the jugular vein the heart was almost instantly arrested in its beating, and, on examination, was found to be relaxed, destitute of contractility, and with its left side full of scarlet blood. He also discovered that if the quinia was thrown into the exposed heart in such a way as to be carried into the coronary arteries, similar phenomena were induced even more rapidly. These experiments made upon the dog have been confirmed upon the frog by Schlockow (*De Chini sulfurici Vi physiologica nonnulla Experimenta*, Vratisl., 1860), and by A. Eulenburg (*Reichert's Archiv für Anatomie*, 1865). The first of these observers found that after section of the vagi quinine induces slowing and enfeeblement of the heart's action; and Eulenburg not only confirmed this, but proved that if the frog's heart be immersed in a solution of quinine its contractions become at once slow and irregular, and soon cease altogether. Many observers agree in asserting that in very large doses (thirty to ninety grains) quinia produces in man lowering of the frequency and force of the pulse. The rate of pulsation has been recorded as falling to below forty per minute, and after poisonous doses the pulse has been imperceptible at the wrist. When the pulse becomes very feeble it may become very rapid; but of course this increased rapidity is indicative not of cardiac strength, but of cardiac weakness. The proof is evidently sufficient to show that in large doses quinia acts as a sedative to the circulation by influencing directly the heart, and probably also in other ways.

I have never, however, been able to perceive any such action in man after ordinary therapeutic doses (three to five grains), and have met with no experiments on animals indicative of its occurrence. I believe, therefore, that in tonic doses quinia produces no perceptible sedation of the circulation.

According to Bonorn and Arvedi, to Magendie, to Monneret, to Melier, and to Baldwin, in animals killed with quinia the blood is found to be dark, defibrinated, fluid, and incapable of forming a clot. Briquet, however, denies that this alteration of the blood is constant or even common in quinia-poison-

* M. Pagès, *Gazette Médicale*, 1846; also Dr. Küchenmeister, *Archiv für Physiol. Heilkunde*, Bd. x.

ing, as he found it in only four out of twenty-three dogs so sacrificed; and he believes that it is merely an accident dependent upon the method of death. In a series of analyses, apparently carefully made, he found that the continued use of quinine augments the proportion of fibrin, but lowers that of the red corpuscles.

In 1867* Prof. Binz announced the fact that quinia added to human blood in the proportion of one part to four thousand immediately checks and in a short time arrests the amœboid movements of the white blood-cells. Confirmation of this has been furnished by Scharrenbroich (*Das Chinin als Antiphlogisticum*, Inaug. Dissert., Bonn, 1867), by Kerner (quoted in *London Practitioner*, vol. vii. p. 321), and by Geltowsky (*London Practitioner*, vol. vii.). The minimum effective strength of the solution has been found to vary in different species of animals, and even in different individuals of the same species.

It is a matter of great interest to determine whether quinia acts in the living organism as on the stage of the microscope; and, to settle this point, Prof. Binz (*Virchow's Archiv*, Bd. xlv., 1869, p. 138) has experimented according to the method of Cohnheim. He found that when the mesentery of curarized frogs to whom quinia had been given was exposed upon the stage of the microscope, no accumulation of white blood-cells in the small vessels, no passage of them out into the tissues, occurred upon irritation; or, if after a time these phenomena commenced, they were at once checked by a small hypodermic injection of the alkaloid. When the inflammatory process had already commenced in a "Cohnheim frog," an injection of quinia would cause the wandering out of the corpuscles to cease, and would bring about a gradual clearing of the white cells from the choked-up vessels. Prof. Binz further took two young cats, and, after poisoning one of them with quinia, examined their blood. In the blood of the unpoisoned animal the white cells were far more abundant than in that of the poisoned cat. From these facts Prof. Binz deduces the conclusion that quinia acts destructively in the system upon the white blood-corpuscles, in the same way as when they are out of the body. A. Martin (*Das Chinin als Antiphlogisticum*, Inaug. Dissert., Giessen, 1868)† has experimentally confirmed the effect of quinia upon the extrusion of white cells of the frog's mesentery, and has found that it also occurs in the centre of parenchymatous organs, such as the liver.

On the other hand, Schwalbe‡ could detect no difference in the blood of a cat before and after poisoning by quinia; and the experiments of Geltowsky (*loc. cit.*) upon frogs and guinea-pigs have yielded similar results: in all

* *Archiv für Mikroskop. Anatomie*, iii., 1867. Consult, also, *Experimentelle Untersuchungen über das Wesen der Chininwirkung*, Berlin, 1868; *Virchow's Archiv*, Bd. xlv., 1869, p. 137. *Berlin. Klinische Wochenschrift*, Nov. 1871.

† Quoted by Binz, *Virchow's Archiv*, Bd. xlv., p. 137.

‡ Quoted by Kerner, *Pflüger's Archiv*, Bd. i. p. 203.

cases after fatal poisoning by the alkaloid, the movements of the corpuscles were found to be very active.

These results are in opposition not so much to the experiments as to the deductions of Prof. Binz,—deductions which seem to me scarcely warranted by his own investigations. The experiments which he performed in the Cohnheim method at most prove only that poisonous doses of quinia prevent or arrest inflammation. The local accumulation and the out-wandering of the blood-corpuscles are the results of a local inflammation or irritation, and if the quinia should in any way check this it would of course put an end to the phenomena mentioned. Again, quinia might, in the case of the cat, have lessened the proportion of blood-cells by checking their formation. Moreover, the experiment was performed under the influence of a very strong dominant idea, and therefore, without confirmation, can hardly be accepted as conclusive.

The exact nature of the action of the alkaloid upon the white blood-corpuscles in the body must, therefore, be considered undetermined, even when poisonous doses are used; when therapeutic doses are employed, the doubt is of course still stronger.

It would seem that quinia acts also upon other portions of the blood than the white corpuscles. Manassein (*Ueber die Dimensionen der rothen Blutkörperchen unter verschiedenen Verhältnissen*, Berlin, 1872) has found that in fever occurring in the lower animals the red corpuscles are diminished in size. If in this condition a decided dose of an antipyretic, such as quinia or alcohol, be given, and the temperature falls, the globules resume their normal size. That the change is due to the fall of the temperature rather than to a direct action of the drug is, I think, demonstrated by the fact of its occurrence whenever the fever heat is lowered by the application of external cold. The experiments of Manassein, therefore, do not prove that quinia exerts any direct action on the red corpuscles. The investigations of Binz (*Archiv für Experimentelle Pathologie und Pharmacologie*, Bd. i., erstes Heft, 1873), however, appear to show that the alkaloid lessens the ozonizing power of the blood; for he found that in young cats, to which he had given a very large but not fatal dose of quinia, the freshly-drawn blood affected the tincture of guaiac much less than it normally should.

When blood is drawn from the body and allowed to stand, acid is developed in it. Zunst (*Beiträge zur Physiologie des Blutes*, Inaug. Dissert., Bonn, 1868), who has studied this subject most closely, divides the investigation into—study of the production of acid in the time from the escape of the blood from the vein to its coagulation, and study of the slow changes which increase its acidity when coagulated until putrefaction has fairly set in. Prof. Binz believes that this development of acid is due to oxidation, and by an elaborate series of experiments has determined that quinia (also sulphate of bebeeria and picrate of soda in almost as great degree) inhibits these changes very greatly in both their varieties. These experiments are in accord with the previous

ones of A. Schulte (*Centralblatt für die Medicin. Wissenschaften*, Nov. 1871); the facts may, therefore, be considered proven.

If ozonized oil of turpentine be dropped into an alcoholic solution of guaiac resin, no alteration of color occurs; but if a drop of blood be added, the blue appears at once: *i.e.*, the blood acts as a carrier of ozone from the turpentine to the resin. Prof. Binz has found that quinia, even in so small an amount as one part in twenty thousand, has a perceptible influence in preventing this. Similarly, when into a dilute watery solution of the sulphate of indigo carbonate of soda is thrown until the reaction is decidedly alkaline, and a little blood, and subsequently ten drops of ozonized turpentine, are added, a green color begins at once to develop, and in a little while passes into the clear yellow of isatin. In this case also the blood acts as a carrier of ozone, and Binz and his pupil Ransoné (*Ueber einige Beziehungen des Chinin zum Blut*, Inaug. Dissert., Bonn, 1871) have found that quinia also inhibits this action, one part of it added to a thousand of the mixture delaying the change of color for an hour. In these experiments Binz used a large number of different salts of quinia, and found that they acted identically. That the action of the alkaloid was on the blood, not on the indigo and guaiac solutions, was shown by the fact that when similar solutions without the blood were shaken in the air and absorbed ozone, the characteristic colorations of its action were produced just as readily when quinia was absent as when it was present. Binz also proved that the red corpuscles were the portions of the blood affected. On adding crystallized hæmoglobin from horses' blood to the guaiac solution he found that it acted as an ozone-bearer between the turpentine and the guaiac, and further demonstrated that quinia had the power of preventing this action.

If it be hereafter established, as seems probable, that quinia exerts a decided antipyretic action (see p. 67) in ordinary fever, it is an exceedingly plausible theory that the lowering of temperature is due to a checking of the ozonizing power of the blood. To attribute, however, the general medical virtues of quinia to an action on the white corpuscles seems to me unreasonable; for from the experiments of Prof. Binz himself upon the lower organisms, it would appear that quinia acts upon all animal germinal matter; and it is probable that the protoplasm of the nervous system, being more specialized than that of the white corpuscles, would be more susceptible of the influence of the alkaloid. Further, according to the experiments of Binz, both conia and camphor act more forcibly upon the white corpuscles out of the body than does quinia. Yet their influence upon the organism is entirely unlike that of quinia. I think these facts are sufficient to show that any theory as to the action of the drug built upon the prime fact discovered by Binz would be at present premature.

As long ago as 1765, Dr. Pringle (*Observations on Diseases of the Army*, London, 1765) called attention to the fact that cinchona bark, in decoction or powder, has the power of preventing for a time putrefaction in flesh; and more recently the subject has been studied by Mayer, by Pavisi, by Hallier

(*Das Cholera-Contagium*, Leipsic, 1867), by Herbst, by Polli, and especially by Binz (*Virchow's Archiv*, Bd. xlv., 1869, p. 68; and *Untersuchungen über das Wesen der Chininwirkung*, p. 20), to whose elaborate articles I must refer my readers for details and references. The experiments of these authorities have demonstrated that quinia in the proportion of one part to three hundred will preserve for a long time flesh, meal, milk, butter, urine, albumen, etc., and will check very markedly the alcoholic fermentation in honey or in syrup. Prof. Binz has demonstrated that this antiseptic action is due to a poisonous influence exerted by the quinia upon the fungi which are the immediate cause of the changes. According to his experiments, the larger infusoria, such as *Paramecia* and *Colpoda*, are killed by a solution of quinia of the strength of one in eight hundred immediately, of one in one thousand after some minutes, of one in twenty thousand after some hours. Upon the ordinary mould *Penicillium*, upon *Vibrios* and *Bacteria*, as well as upon the higher infusoria, quinia acts with a similar fatality. In the case of the *Vibrios* and *Bacteria* a decidedly stronger solution than the one mentioned is required to quiet movement. Bochefontaine (*Archives de Physiologie*, July, 1873) found that a solution of one per cent. was needed for a vigorous rapid action, and that some active granules could even be found in it after three days. According to Binz, the singularity of the influence of quinia is shown by the fact that on *Paramecia* and *Colpoda* a solution of salicin, in the proportion of one part to forty, has no influence. Indeed, these infusoria were not even affected by this strong solution of salicin, and they endured a solution of morphia of one part to one hundred and twenty for an hour, and a five per cent. solution of strychnia for some minutes. Although fungi will appear after a time in ordinary solution of the sulphate of quinia, I think it must be considered well established that this and other salts of the alkaloid are extremely poisonous to the fungi of putrefaction and of other ordinary fermentations.

In 1871, Dr. Monteverdi announced (*Annales et Bulletin de la Société de Médecine de Gand*, May, 1871) that quinia is a uterine stimulant, causing at times in the gravid womb contractions sufficiently violent to induce abortion, and, when given during labor, intensifying greatly the uterine pains, and after labor causing rapid expulsion of the placenta and arresting uterine hemorrhage; affirming, further, that in amenorrhœa or in menorrhagia from uterine inertia its action is no less marked. Although this has been received as new the world over, so long ago as 1855 Dr. John S. Wilson (*Southern Medical and Surgical Journal*, p. 341, 1855) called attention to the uterine action of quinine, and in 1860 reasserted his belief (*Southern Journal of Medicine*, Sept. 1860), which in the mean while had been confirmed by Dr. J. H. Rich in the *Charleston Medical Journal and Review*; and in 1858 Dr. Jos. J. West (*Savannah Journal of Medicine*, vol. i. p. 19) wrote, "Many regard the use of quinine as dangerous and even criminal in any diseases in pregnant women. The belief of these persons is that this substance exercises a direct influence upon the uterus, causing powerful con-

tractions and expulsion of the fetus. And to support this notion they are ready to bring forward innumerable instances of abortion after its use,—of cases of sudden suppression relieved by a prompt use of the same remedy." He then goes on to say that these abortions, etc., were due to the intermittent fever and not to the drug. Surely this is enough to show that the oxytocic action of quinia was believed in many years ago by numbers of our Southern practitioners. The question now is whether the drug has any such action. It is evident that the answer to this should be made out in three different ways. First. Is there any evidence of quinia producing abortion in healthy women or in females of other animals? Second, How strong is the evidence of its producing abortion in women suffering from ague? Third. What is the evidence in regard to the action of quinia during labor?

In regard to the first of these sub-questions, the only affirmative evidence I have met with is in the experiments of M. Rancillia (*L'Union Médicale*, 1873), who saw abortion in two bitches follow the administration of from six to nine grains of quinia: as the pups in one case were already dead before the administration of the drug, it would seem that this investigation was not on such a scale as to be at all conclusive. On the other hand, I have met with no evidence that quinia is capable of inducing abortion in healthy pregnant women. Dr. Sayre's case (*American Practitioner*, 1871, p. 260) is certainly no proof whatever that quinia will originate labor, as labor had commenced under the influence of the hot and cold douche and other measures employed *before* the quinine was given. Prof. Chiara, of Milan, has furnished (*L'Union Médicale*, Nov. 20, 1873) very strong evidence that quinia is incapable of originating uterine contractions in healthy pregnant women. In his public service, two doses of a gramme (15.34 grains) each were given without effect daily for two successive days to eight women all in the eighth month of pregnancy. It being necessary to cause abortion, one gramme was given daily to one woman for seven days, to another for three days, without in either instance any effect, so that the labor had to be brought on in the usual manner.

In answer to the second sub-question, some evidence has already been adduced to show that abortion may be so caused. To it may be added the assertion of Dr. Walraven (*Boston Medical and Surgical Journal*, 1873) that he has frequently seen the exhibition of quinia followed by abortion, the record of two cases of such character by Dr. Burt (*Medical and Surgical Reporter*, 1870), and no doubt the affirmations of others which I have not seen. Opposed to this, however, is the overwhelming fact that the great body of the profession have for centuries been giving quinia in one form or other to pregnant women indiscriminately, and if abortion had been produced it must have been noted long ago. Further direct testimony is not wanting. Malaria often induces abortion, and Dr. Erwin (*St. Louis Medical and Surgical Journal*, March, 1872). Dr. Jas. C. Harris (*American Practitioner*, April, 1872), and Dr. A. Russwurm (*American Practitioner*, 1871, No. 4, p. 127)

testify from personal experience that quinia will arrest abortion from such cause. Dr. J. A. Ashford (*National Medical Journal*, Oct. 1871), Dr. Beauchamp (*American Practitioner*, 1870), Dr. Rooker (*ibid.*), Dr. J. S. May (*ibid.*), Dr. A. d'Arcourd (*Medical News and Library*, May, 1873), have given the alkaloid to hundreds of pregnant women in large doses without result. Other testimony might be adduced; but it seems to me incredible, in the face of daily experience, that even the largest therapeutic doses of quinia are abortifacient in malarial fevers or in health.

The evidence is much more doubtful as regards a negative answer to the third sub-question. It is evident that, as during natural labor quinia has not been used by the profession, there is no general experience upon the subject. Drs. Bouqué (*L'Union Médicale*, t. xii. p. 544, 1871), Deneffe, and Doubuë (*Boston Medical and Surgical Journal*, p. 160, 1873) have all used it in uterine inertia and in post-partum hemorrhage, with the best results. Dr. John Lewis has employed it since 1853 for such purposes (*Medical and Surgical Reporter*, 1870), and Dr. L. A. Hamilton (*Cincinnati Lancet and Observer*, 1861, p. 141) has long been in the habit of relying upon it in these cases. In the report of the Committee on Obstetrics to the Minnesota State Medical Society (*Northwestern Medical and Surgical Journal*, vol. iii. p. 425) the testimony of various physicians is given in favor of this action of the alkaloid. In the *Philadelphia Medical Times* (vol. iii. p. 276) Dr. H. G. Landis details six cases in which quinia appeared to exert during labor a very decided oxytocic effect, but also states that he has used it in other cases (number not given) with negative results. His doses do not appear to have been large enough to test the matter. In two cases of very great uterine inertia in my own practice, very powerful contractions came on shortly after the administration of ten grains of the sulphate of quinia. On the other hand, the experience of Prof. Chiara (*loc. cit.*) is strongly opposed to the idea that quinia has any oxytocic effect whatever. In ten cases of uterine inertia he administered it without effect; in seven women it was given during natural labor without intensifying the already active pains; in three cases of post-partum hemorrhage, quinia was exhibited without checking the flow.

From the evidence that has been adduced, it may, I think, be concluded that quinia has no power to originate uterine contractions in the pregnant woman,—that although there is some reason for believing that in labor full doses of it (ten to fifteen grains) do act as a stimulant to the pains, yet the question must be considered still *sub judice*.

The manner in which quinia finds entrance into the blood has been especially studied by Dr. Kerner (*loc. cit.*). As the gastric juice is very acid, it is evident that the alkaloid will be rapidly dissolved in the stomach, and be put into the conditions most favorable for its absorption; if, however, the salt of quinia escape from the stomach into the intestines, it will be liable to be precipitated by alkaline juices as well as by the bile, whose acids form very insoluble salts with it. The presumption is therefore strong that, when gastric

absorption fails to take place, at least a portion of the quinia will pass out with the fæces. That this actually does occur has been proven by Kerner and others, who have found the alkaloid in the excrement of persons taking it. As the blood is alkaline, it would appear probable that the quinia salt so soon as entering it would be precipitated. That this does not occur, according to the researches of Kerner, is due to the solvent power of the gases contained in normal blood.

The authority mentioned found that one thousand parts of blood which was defibrinated and deprived of its gases, at a temperature of 36° C. dissolved in an hour only 0.398 parts of pure quinia. Water saturated with carbonic acid gas dissolves the sulphate of quinia pretty freely; and Kerner also experimentally determined that when a neutral solution of a salt of quinia is added to a very dilute solution of carbonate of soda no precipitate occurs. It would appear, then, that the quinia is held in solution in the blood by reason of the loosely-combined carbonic acid gas in that fluid.

It has been proven by the analyses of Landerer (*Repertorium für Pharmacie*, Bd. xxv., 1836), of Dietl (*Wiener Med. Wochenschrift*, 1852), of Briquet (*loc. cit.*), and of Binz, that quinine escapes from the body through the kidneys. According to Briquet, it generally may be found in the urine half an hour after the administration of a large dose. Its removal, according to the researches of Binz, goes on slowly, for it is stated (*loc. cit.*, p. 167) that in six experiments only a little more than two-thirds of the ingested quantity was excreted in the first forty-eight hours. Dr. L. Thau, however, in three experiments, out of the 4.4586 grammes of the alkaloid which were given recovered from the urine passed during the forty-eight hours 4.3 grammes, so that only 0.1586 grammes remained unaccounted for. A portion of this residue was perhaps lost in the chemical operations; but it is probable that some of the quinia is eliminated through other channels than the kidneys, since Prof. Binz has found it in the saliva of a poisoned dog, and Landerer (*Buchner's Repertorium*, 1839 and 1842) states that he has detected it in the urine, sweat, tears, milk of nursing women, and in the serum of dropsical effusions. In regard to the rate of its elimination, Dr. Thau determined that from a third to somewhat less than half of the ingested quantity escaped in the first six hours, and that in the first twelve hours about three-fourths were excreted. Dr. G. Kerner (*Pflüger's Archiv für Physiologie*, 1870) has found that the quinia as excreted is in an amorphous, uncrystallizable form. He also has discovered in the urine of persons taking quinia a peculiar substance, sometimes amorphous, sometimes in acicular prismatic crystals, free from bitter taste, possessing the quinia inflorescence, which he believes to be a derivative formed in the body from the ingested alkaloid. He has not been able to get this substance in such quantity as to analyze it or further examine it, but has produced a principle (*dihydroxyle quinia*) which he believes to be identical with it by acting on quinia with the permanganate of potash. An elaborate series of experiments have shown that the dihydroxyle quinia is physiologically inert.

Ranke was, I believe, the first to notice that quinia produced a great decrease in the elimination of uric acid. This fact has been confirmed by Dr. G. Kerner, who has made a very elaborate study of the action of the alkaloid upon renal elimination, with the following results. When about nine grains of quinia were taken in divided doses during the course of the day, the urea was decreased not quite one-eighth, the uric acid to a little less than one-half, the kreatinine was slightly increased, and the nitrogenous material decreased about one-ninth. When a very large dose (thirty-eight grains) was taken in the morning, the urea and the kreatinine were each decreased about one-fourth, as was also the collective nitrogenous material; the phosphoric acid about one-fifth; and the uric acid was lessened about four-fifths. The experiments of Rabuteau (*Bulletin Thérapeutique*, t. lxxv. p. 475) upon the dog, so far as they go, confirm this decrease of the elimination of urea.

THERAPEUTICS.—At present our estimate of the value of quinia in disease, and our knowledge of its therapeutic use, rest solely upon clinical observation. We are not as yet able to apply what information we have of its physiological action, nor, indeed, are we able to explain its known clinical action by its observed effects on the healthy organism.

On account of its power of arresting or preventing putrefactive fermentation by killing the microscopic entities which produce such changes, Prof. Binz has recommended it in the so-called *septic diseases*. The chief evidence which he produces is in some ten experiments made upon dogs and rabbits. In each of these experiments two similar animals were poisoned with putrescent liquids, and to one of the pair quinine was freely administered. In two cases the cinchonized animal recovered, whilst its fellow perished; in three experiments neither of the animals died; and in the other five trials the cinchonized animal lived from two to twenty-four hours longer than the other. These experiments are certainly too few and indecisive to prove in any degree Prof. Binz's view. To my mind they indicate very strongly that quinia has no such influence over the disease as he claims for it. If living germs in the blood were really the cause of the septic symptoms, and quinia killed such germs, its action would be as manifest and unmistakable as it is in intermittent fever. The results of Prof. Binz's experiments indicate no such specific action, but rather that the quinia in such cases does good by sustaining the nervous system, or in some other unknown manner. In *pyæmia* in man, quinia has been frequently employed, but exerts no specific action. In *puerperal* and other *low fevers* it is sometimes given, in large doses, often with apparently great advantage. In Germany a powerful antipyretic action has been claimed for it, but the matter is certainly still *sub judice*. It has not, that I know of, been proven that therapeutic doses of it lower, to any marked extent, animal temperature in the healthy man. Indeed, even Dr. C. Liebermeister (*Deutsches Archiv für Klinische Medicin*, Bd. iii., 1867) acknowledges that numerous experiments have shown him that it has no constant action of such sort, and details a case in which forty grains admin-

istered within seven hours caused no depression of temperature. The same authority claims, however, that by a very large number of experiments he has demonstrated its power of lessening fever-heat. In one hundred and seventy-eight observations in *typhoid fever*, twenty grains of the quinine having been given during the night, the morning temperature was lower than that of the previous evening by, on the average, 1.63° C. On one hundred and seventy-six different occasions a scruple of quinine was given during the day: sixty-nine times the temperature was lower in the evening than in the morning, ten times the same as in the morning, and ninety-seven times higher than in the morning.

Dr. G. Kerner and Dr. Jürgensen have each noticed that full doses of quinia appear in a healthy man not to affect sensibly the temperature, but to prevent the rise which normally occurs from exercise. Thus, in Kerner's experiment, certain gymnastic exercises, which when performed in his ordinary state elevated his bodily temperature 2° C., only affected the latter to the extent of 0.2° to 0.35° C. when quinia was freely exhibited.

In the fever following section of the cord in animals (*London Practitioner*, p. 4, 1870) Binz has noticed that quinia decreases the temperature somewhat: he says, however, if the conditions of the fever are too favorably constituted the effect of the quinine fails thoroughly. The action of the quinia must, therefore, be a feeble one. The drift of our present evidence seems to indicate that quinia exerts in febrile disease a feeble but decided antipyretic action. My own experience is that when twenty grains are given in the twenty-four hours, the drug, in cases of low fever, does more harm by its over-action on the cerebrum than it does good by its antipyretic influence, but that in doses of from ten to fifteen grains a day it is often of service, apparently supporting nerve-power and at the same time tending to lower temperature.

As a simple *tonic*, quinia is used by every one; but I have never been able to see that the ordinary combination of it with iron is of much more value in simple cases of debility than iron with quassia or other simple bitter.

When given in very large doses, quinia, as has already been shown, acts as a powerful depressant, and as such has been used by Briquet and other French physicians in *rheumatism*. As much as sixty or seventy grains a day have been given, and it is beyond dispute that under the influence of these heroic doses the symptoms of inflammatory rheumatism have often rapidly abated; but the method has found little favor out of France, and is less efficient and more dangerous than other plans of treatment now in vogue.

In *inflammatory rheumatism*, after the acute symptoms have abated, when the patient shows evident signs of weakness, especially if there be profuse sweating during sleep, twelve grains of quinine daily are often of great service.

The great value of quinia as a therapeutic agent consists in the wonderful power which it has over all forms of disease dependent upon *malarial poisoning*. Various theories have been brought forward to account for the antiperiodic action of quinia; but, as we know nothing of the nature of

malaria or of the method of its action, and as we are very partially informed as to the effects of quinia upon the healthy organism, it seems to me perfectly futile to endeavor to explain why it averts a paroxysm of intermittent. I shall not, therefore, occupy space with such discussion. The present seems, however, the fitting place to notice the discovery of Dr. H. Bence Jones in regard to the existence of a body similar to quinia in the animal organism.

Quinia and its salts have the remarkable property of converting rays of heat into light, or, in other words, of rendering visible the ordinary invisible rays of the solar or other spectrum. Connected with this fact is probably the phenomenon known as the fluorescence of quinia. When a colorless watery solution of one of its salts is examined, a pale-blue line upon the surface is very noticeable, and Prof. Stokes has shown that solution of quinia has the power of entirely stopping certain of the rays of light, so that when a beam is transmitted through it to light up a second vessel of the solution, this latter displays no fluorescence. Dr. H. Bence Jones (*Lectures on Pathology and Therapeutics*, London, 1867) has found that when the electric light is used this test is so delicate that one grain of the alkaloid may be detected in 1,450,000 grains of water. He has also discovered that man and animals are pervaded by a substance which, in its action on light and in many chemical reactions, very closely resembles, if it be not identical with, quinia. Believing this substance to be probably an alkaloid, he has given it the name of *animal quinoidine*. Drs. Edward Rhoads and William Pepper, Jr. (*Pennsylvania Hospital Reports*, vol. i., 1868), have made observations upon ten cases of malarial fevers in which no quinia had been used, and have found the fluorescence of the blood to be from 0 to $1\frac{1}{2}$ instead of from 3 to 6, which is said to be normal. The significance of these results is at once apparent; but before sufficient grounds shall have been obtained to build a theory upon, far more extended researches on the variance of the fluorescence in health and in diseases other than malarial are necessary. Especially is this the case since Dr. Chalvet (*Schmidt's Jahrbücher*, Bd. cxli, p. 152, from *Gazette Hebdomadaire*, 2d series, t. v., 1868) has found that this fluorescent body exists in various foods and even in wine, and is therefore probably not of animal but of vegetable origin.

Quinia in its relations to *malarial fever* may be considered, first, as a prophylactic, secondly, as a curative agent.

The value of the daily use of quinia to persons exposed to a malarial atmosphere has now been thoroughly tested in all portions of the world. In North and South America, in Europe, in Africa, in India, the prophylactic powers of quinia have been tried on the largest scale in connection with the military and naval services, and the testimony is unanimous in favor of the drug. A single citation will serve to illustrate this fact. Dr. J. B. Hamilton (*Indian Medical Gazette*, Nov. 1, 1873) reports the case of a battery of one hundred and thirty-five men, quartered at Jubbulpore, East Indies, in the same barracks with an infantry regiment. Each of the artillerists received

three grains of quinine every other day; to the infantry none was given. The result was that whilst three hundred out of the five hundred men of the regiment were sick at one time with malarial disease, at no period was more than four per cent. of the battery affected. The dose of quinia as a prophylactic may be considered as three grains a day, or possibly, in very deadly climates, three grains in the morning and two in the evening.

In *intermittent fever*, when there is sufficient time, it may be well to precede the quinia by a mercurial or other purge. If the expected paroxysm be so near that there is not sufficient time for the action of the purgative, the antiperiodic should be administered without previous preparation of the patient. The value of purgatives in obstinate intermittents, as an adjuvant to quinia, is often overlooked, although in some cases the employment of purgatives, and of such diuretics as cream of tartar, seems to be almost essential for the successful use of quinia.

When there is necessity for prompt action, the antiperiodic may be given in a single dose, or in any other method that the circumstances of the case will allow; but ordinarily the best plan is to commence the exhibition of the drug about eight to ten hours before the expected paroxysm, and to continue in hourly doses until from three to four hours before the attack is due.

In *pernicious fever*, or *malignant malarial poisoning*, no time should be lost after the first paroxysm in getting the patient cinchonized, as it may be uncertain whether the attack be of the quotidian or tertian type. At least thirty-five grains of the alkaloidal salt should be administered during the first twenty-four hours of intermission, and twenty-five grains during the second; in very severe types of the disease much larger doses even than these are necessary, less than fifty grains of the drug sometimes appearing to do but little good.

In *remittent* or *bilious fever* it may often be advisable to give purgatives and febrifuges, but it is not proper to delay the exhibition of the antiperiodic on their account. As soon as the remission has appeared, the exhibition of quinia should be begun. Local inflammations or even severe cerebral symptoms occurring during a remittent fever are no contra-indications to the use of the specific. When gastritis exists, on account of the local irritant action of quinia, other channels of entrance to the body than the stomach should be employed.

When the symptoms in remittent fever are severe and seemingly continuous, it may be not only proper, but necessary for the saving of life, to exhibit quinia freely during the period of fever. In large doses the alkaloid is probably antipyretic as well as antiperiodic, and I do not know of any theoretic or clinical objection to its use during the period of fever.

In *malarial intermittent neuralgia*, as in all other forms of abnormal manifestations of malarial disease, quinia is efficient, although it may be necessary to use it in large doses.

In *neuralgia* which, although not dependent upon malaria, assumes the

intermittent type, quinia will often temporarily set aside the paroxysmal attacks, and sometimes effect a cure. The same fact may be stated in broad terms as true of *all non-malarial intermittent* affections. In the great majority of such cases, unfortunately, the action of the quinia is only temporary, and any controlling power is soon lost.

ADMINISTRATION.—Owing to its bitter taste, quinine is generally given in pill, which may be made with gum, or simply by adding a little sulphuric acid to the alkaloidal salt and quickly rubbing up the pasty mass into pills before it hardens. Whenever a rapid action is desired, the quinine should be given in powder, or in solution made by adding a drop of dilute sulphuric acid for every grain of the salt. In the use of pills of quinine care should be exercised to see that they are soft and fresh, for when old and hard they not unfrequently pass through the bowels unchanged. The taste of the powder is best covered by chocolate or by liquorice.

CINCHONINÆ SULPHAS—SULPHATE OF CINCHONIA.

The pure alkaloid cinchonia crystallizes in prisms and needles. The official *sulphate* of cinchonia is in short oblique prisms of a very bitter taste, soluble in fifty-four parts of water, more freely in boiling water, readily soluble in alcohol. From its solution in chlorine-water it is precipitated white by ammonia. Bill's test for it consists in adding the ferrocyanide of potassium in slight excess to its solution. A yellowish-white curdy precipitate is the result. On gently heating, this redissolves, but is again deposited, when the liquid cools, as abundant golden-yellow crystals.

PHYSIOLOGICAL ACTION.—The physiological action of cinchonia is similar to but less powerful than that of quinia. Thus, Conzen (quoted by Husemann) has found that its action on infusoria and on fermentation is similar to but weaker than that of its sister alkaloid, and that on the movements of the white blood-corpuscles its influence seems transient. Upon dogs, according to Bernatzik's experiments, the lethal dose of cinchonia is to that of quinia as 5 is to 4. The history of cinchonia in the organism appears, therefore, to be parallel with that of quinia.

THERAPEUTICS.—As an antiperiodic, cinchonia exerts a similar influence to quinia, but is probably about one-third weaker than that alkaloid, and must be used in correspondingly larger dose. Dr. J. B. Hamilton (*loc. cit.*) affirms as the result of experiment that cinchonia as a prophylactic against malaria is even superior to quinia.

As a tonic I have never been able to perceive that cinchonia acts differently from quinia.

Under the name of *chinoidine* is sold by the manufacturers the black, intensely bitter residue left upon the evaporation of the mother-liquor after the crystallization of the alkaloids out of it. This substance contains probably amorphous quinia and cinchonia, besides quinidia and cinchonidia. It

is an excellent tonic and antiperiodic, and on account of its cheapness is largely used in some of the eleemosynary institutions of this city. It should be administered in about double the dose of quinine, and is most efficient in solution, but on account of its taste is often given in pills. Its solution should be made with acetic acid and water, aromatics being freely added, as it is apt to cause nausea.

The following preparations of *cinchona* are officinal, and may be used as tonics. As such I think they are superior to the simple alkaloids:

Infusum Cinchonæ Flavæ—*Infusion of Yellow Bark* (℥i to Oj). Dose, a wineglassful.

Infusum Cinchonæ Rubræ—*Infusion of Red Bark* (℥i to Oj). Dose, a wineglassful.

Tinctura Cinchonæ—*Tincture of Yellow Bark* (℥iii to Oj). Dose, one to two teaspoonfuls.

Tinctura Cinchonæ Composita—*Compound Tincture of Bark*. *Huxham's Tincture*. As a tonic, I think, the best preparation of the bark; a very elegant remedy in convalescence. Dose, one teaspoonful to a tablespoonful.

Extractum Cinchonæ Fluidum—*Fluid Extract of Bark*. Dose, as a tonic, five to twenty drops.

Extractum Cinchonæ—*Extract of Bark*. Dose, as a tonic, five to fifteen grains.

AROMATICS.

There are certain remedies usually spoken of in treatises on *Materia Medica* under the general head of tonics, which might more properly be considered local stimulants, acting, as employed in medicine, upon the alimentary canal. These are the so-called aromatics, substances dependent for their virtues upon the presence of a volatile oil.

The volatile oils are essentially local irritants, causing when taken into the mouth intense burning pain; when confined upon the skin, rubefaction, blistering, and finally, if the contact be very prolonged, more destructive changes. Internally, taken in very large doses, they cause burning pain in the stomach, increased activity of the circulation, and a species of intoxication. In sufficiently large quantities they are irritant narcotic poisons. When administered in therapeutic doses they act almost exclusively upon the alimentary canal. As compared with that of the simple bitters, their influence is more powerful and more transient. They do not permanently increase the digestive power, but simply increase action for the time being. They are employed chiefly—to give pungency to bitter tonics; as *carminatives*, to stimulate the intestines to contract upon and expel flatus; to prevent the griping of purgatives; to disguise the taste of medicines, and to render nauseating drugs acceptable to the stomach; to act as condiments, and aid in the digestion of the food.

Some of the tonic drugs containing a volatile oil also have in them a bitter principle which modifies their action. Such drugs may be known as *aromatic*

bitters; as bitters they are less powerful than such drugs as quassia, and are especially indicated where the stomach is delicate and easily nauseated.

Inflammation of the stomach or bowels is the chief contra-indication to the use of aromatics. Unlike the simple bitters, they are often very useful in *diarrhœa* of nervous irritability or of relaxation, when no decided inflammation exists.

AROMATIC BITTERS.

ANTHEMIS—CHAMOMILE.

Roman or true Chamomile is the dried flowers of *Anthemis nobilis*, a composite of Europe. They are sometimes single, sometimes double. The single are more aromatic than the double florets. Chamomile contains a bluish or sometimes greenish volatile oil, a bitter principle, and a small amount of tannin.

Matricaria or German Chamomile is the flowers of *Matricaria Chamomilla*, which are decidedly smaller than those of the ordinary chamomile, and have a stronger, less agreeable odor and taste. Their volatile oil is very similar to that of chamomile.

THERAPEUTICS.—An excellent stimulant tonic. Especially useful in convalescence.

ADMINISTRATION.—*Infusum Anthemidis*—*Infusion of Chamomile* (℥ss to Oj). Dose, one to two wineglassfuls before meals.

SERPENTARIA—VIRGINIA SNAKE ROOT.

The root of *Aristolochia serpentaria* and of *A. reticulata*, small herbal plants of the United States. It occurs as fine brittle rootlets attached to a small head, of a camphorous odor and taste, and contains a volatile oil, a yellowish-green resin, and a bitter principle.

THERAPEUTICS.—An elegant stimulant tonic, especially useful as an adjuvant to more powerful bitters. In overdose it is said to cause vomiting, and even purging.

ADMINISTRATION.—*Infusum Serpentariæ*—*Infusion of Serpentaria* (℥ss to Oj). Dose, a wineglassful.

Tinctura Serpentariæ—*Tincture of Serpentaria* (℥ii to Oj). Dose, one to two fluidrachms. An elegant addition to tonic mixtures.

Extractum Serpentariæ Fluidum—*Fluid Extract of Serpentaria*. Dose, twenty drops.

CASCARILLA is the bark of *Croton Eleuteria*, a shrub growing in the West Indies. This bark occurs in quills or rolled pieces, and is to be distinguished by its outer grayish and inner deep-chocolate surface, by its spicy bitter taste, and the pleasant musk-like odor which it gives whilst burning. It contains tannin, volatile oil, and cascarrillin, a neutral, bitter, crystallizable principle. Its therapeutic action is very similar to that of *serpentaria*.

Infusum Cascariillæ—*Infusion of Cascarilla* (℥i to Oj). Dose, a wineglassful.

ANGUSTURA is the bark of *Galipea officinalis*, a small tree of Northern South America. It occurs chiefly in flat fragile pieces, whose surface usually exhibits under the microscope numerous white points or lines. It contains a bitter principle and a volatile oil, and may be used as a substitute for cascarilla or for Virginia snake root.

Infusum Angusturæ—*Infusion of Angustura* (℥ss to Oj). Dose, a wineglassful.

TRUE AROMATICS.

CINNAMOMUM.—*Cinnamon* is the bark of *Cinnamomum Zeylanicum*, a native of Ceylon, and of *C. aromaticum*, growing in China. The finest variety of cinnamon, that from Ceylon, occurs in long, closely-packed quills of a thin, very aromatic bark. *Cassia Bark*, or *Chinese Cinnamon*, is coarser, more broken, and less aromatic. Both varieties contain tannic acid and a volatile oil, which on account of its great fragrance and very pleasant taste is largely used as an adjuvant, or to disguise the flavor of less agreeable drugs.

Oleum Cinnamomi—*Oil of Cinnamon*. A yellowish, or, when old, brownish oil. Dose, one to three drops.

Aqua Cinnamomi—*Cinnamon Water* (f℥ss to Oij). Dose, half a fluidounce to a fluidounce.

Spiritus Cinnamomi—*Spirit of Cinnamon* (oil ℥i, alcohol ℥xv). Dose, half a fluidrachm.

Tinctura Cinnamomi—*Tincture of Cinnamon* (℥jss to Oj). Dose, half a teaspoonful to a teaspoonful.

Pulvis Aromaticus—*Aromatic Powder* (cinnamon, ginger, cardamom, nutmegs). Dose, ten to twenty grains.

CARYOPHYLLUS.—*Cloves* are the unexpanded flowers of *Caryophyllus aromaticus*, a tree growing in the Molucca Islands. This aromatic, largely used as a spice, contains an exceedingly pungent volatile oil, officinal as *Oleum Caryophylli*. The latter is a yellowish oil, becoming dark by age, which, besides being used as a carminative and aromatic, is frequently employed to benumb sensitive dentine, or even exposed pulp, in caries of the teeth. Dropped upon a piece of cotton and placed in the cavity, it will very frequently cure toothache. Dose, one to two drops.

Infusum Caryophylli—*Clove Tea* (℥ii to Oj). This is made with boiling water; the dose is a wineglassful.

MYRISTICA.—*Nutmeg* is the kernel of the fruit of *Myristica moschata*, a tree inhabiting the Molucca Islands. The nutmeg contains both a fixed and a volatile oil.

The mace of commerce (*Macis*, U. S. P.) is the *arillus* or outer imperfect

supernumerary coating of the seed. It also contains a volatile oil, which is said to be identical with that of the nutmeg. The nutmeg is thought to be possessed of a slight narcotic power, and is said when a large dose is taken (one or two) to produce a dreamy, half-unconscious condition. The volatile oil (*Oleum Myristicæ*, U. S. P.) is officinal. Dose, two to five drops.

PIMENTA.—*Pimento*, or *Allspice*, is the unripe berries of the *Eugenia Pimenta*, a tree, native of the West Indies. It contains a green fixed oil and a volatile oil (*Oleum Pimentæ*), which is officinal. Dose, two to five drops.

CARDAMOMUM.—*Cardamoms* are the fruit of the *Elettaria Cardamomum*, which grows in the East Indies. They consist of tough, seemingly fibrous, generally more or less triangular capsules, containing a number of small, hard, very aromatic seeds. The capsule is itself dry and tasteless. In commerce cardamoms are divided into three varieties, according to their length.

According to Trommsdorf, the seeds contain, besides 10.4 per cent. of fixed oil, 4.6 per cent. of a colorless, highly aromatic, volatile oil. Cardamom is a very grateful aromatic, much less stimulating and heating than most of the other substances of its class.

Tinctura Cardamomi—*Tincture of Cardamom* (℞ii to Oj). Dose, half a fluidrachm.

Tinctura Cardamomi Composita—*Compound Tincture of Cardamom*. A very elegant addition to, or vehicle for, tonic medicines. Dose, one to two fluidrachms.

ZINGIBER.—*Ginger* is the dried rhizome or root-stock of the *Zingiber officinale*, growing in the East and West Indies. *Green Ginger* is the *fresh* rhizome. The *Black Ginger* is the root-stock dried with the epidermis on; the *White* or *Jamaica Ginger* is the same, deprived of its epidermis. The fresher ginger is, the greater is its power, and by time and exposure it becomes completely inert. Its active principles are a soft, acrid, aromatic resin, and a yellow volatile oil. Ginger is much used in domestic medicine as a stimulant carminative in *colic*; given in hot water, it is also used as a sudorific and stimulant in the pain due to *suddenly-suppressed menstruation*. It is often added with advantage to other remedies in *dyspepsia*.

Infusum Zingiberis—*Infusion of Ginger* (℥ss to Oj). Dose, a wine-glassful.

Syrupus Zingiberis—*Syrup of Ginger*. Used as a cordial drink or vehicle. Dose, half a fluidounce to a fluidounce.

Tinctura Zingiberis—*Tincture of Ginger* (℞iv to Oj). Dose, half a teaspoonful.

Extractum Zingiberis Fluidum—*Fluid Extract of Ginger*. Dose, ten drops.

Oleoresina Zingiberis—*Oleoresin of Ginger*. Dose, one to three minims, well diluted.

PIPER.—*Black Pepper* is the unripe berries of the *Piper nigrum*, a woody vine-like plant growing in the East Indies. *White Pepper* is the ripe berries, stripped of their skin and dried. It is much inferior to the ordinary variety. The active principles of black pepper are a soft acrid resin and a pungent, fiery, volatile oil.

In 1819 Oersted discovered in black pepper *Piperin*, which crystallizes in colorless, glistening, four-sided, truncated prisms, of a neutral reaction, but capable of combining with acids to form salts. When pure it is tasteless; but very commonly it has a burning taste, due to the presence of some of the volatile oil of pepper. The possession of very active antiperiodic properties* has been claimed for piperin, and it was for a time employed in *intermittent fever*, but has fallen into complete disuse. The dose as an antiperiodic is four grains repeated once or twice during the interval between the paroxysms. Pepper is very largely used as a condiment; but, as its taste is more hot than aromatic, it is rarely used internally in medicine except as an addition to simple bitters or to antiperiodics, generally in the form of the officinal *oleoresin*, the dose of which is a minim. In atonic *dyspepsia* the latter preparation is a most excellent adjuvant to tonic pills.

CAPSICUM.—Under the general term of *Capsicum* the U. S. Pharmacopœia recognizes both the ordinary West India *Cayenne* pepper, the fruit of *Capsicum annum*, and the product of Liberia, the fruit of the *C. fastigiatum*, the African pepper. The berries of the first-named variety are large, bright red, and conical or ovate; those of the second are less than an inch long, and much more fiery than their compeers. *Capsicum* contains as its active principle an exceedingly acrid oleoresin. The name of *Capsicin* has been applied by different observers to the oil, to the resin, and to their combination, but should, I think, be dropped, as having no definite meaning.

Capsicum is a very powerful local irritant, its oleoresin when applied to the skin producing in a very few minutes intense pain and redness, and finally destroying the cuticle. In the alimentary canal it acts in a similar manner: thus, moderate doses produce merely a pleasant feeling of warmth in the stomach, whilst overdoses may cause gastro-intestinal inflammation, with severe pain as well as vomiting and purging. The chief use of *Cayenne Pepper* is as a condiment; yet it is often added with advantage to tonic pills to increase their immediate action on the stomach. When there is habitual feeble digestion, with flatulence, its free use on food sometimes seems to do good.

In adynamic disease, especially as occurring among drunkards, capsicum is often very useful by stimulating the stomach up to the point of digesting food.

Locally, either as the diluted tincture in a gargle, or applied in powder or

* For a discussion of the subject, see *Die Pflanzenstoffe*, p. 492.

tincture by means of a swab, it is useful in *severe tonsillitis*, especially in that accompanying scarlet fever.

ADMINISTRATION.—The dose of capsicum is four to five grains in pill form. Of the officinal *Oleoresina Capsici*, which is to be preferred on account of its lesser bulk, from half a grain to one grain.

Infusum Capsici—*Infusion of Capsicum* (℥ss to Oj). Used as a gargle.

Tinctura Capsici—*Tincture of Capsicum* (℥ss to Oj). Used locally, or sometimes in half-drachm doses in cases of drunkards.

OLEUM CAJUPUTI.—The *Oil of Cajeput* is obtained from the leaves of *Melaleuca Cajuputi*, a tree growing in the Molucca Islands. This volatile oil is of a green color, peculiar fragrant odor, and burning, camphoraceous taste. It is not very irritating to the skin, and is exceedingly destructive to low forms of life, and consequently has been used as a *parasiticide* externally, and even internally against the *Ascarides*. I have never used it except as a carminative and aromatic stimulant in cases of *intestinal pain* and *spasm*, and in *serous diarrhœa*. When employed in these affections it is very efficient. As a counter-irritant it has been used in *rheumatism*; as a stimulant to the skin in *psoriasis*, *acne rosacea*, and *pityriasis*. The dose internally is from ten to fifteen drops.

CANELLA is the bark of the *Canella alba*, a tree growing in the West Indies. This bark contains a bitter extractive, as well as a fragrant very acrid volatile oil. It is never used except in combination. *Pulvis Aloës et Cancellæ*—*Powder of Aloës and Canella*. *Ilia Pica*. A popular remedy in cases of *amenorrhœa* to which aloes is adapted. Dose, ten to twenty grains.

AURANTII AMARI CORTEX—BITTER ORANGE PEEL. AURANTII DULCIS CORTEX—SWEET ORANGE PEEL. *Confectio Aurantii Corticis*—*Confection of Orange Peel*. The orange peels are themselves scarcely medicinal, but are officinal as affording preparations much used as vehicles.

AURANTII FLORES.—The flowers of the orange are officinal for the preparation of the *Orange Flower Water*—*Aqua Aurantii Florum*—which is used as an elegant vehicle, free from medicinal properties.

The fruits of the following umbelliferous plants: *Fœniculum vulgare*. *Carum Carui*, *Coriandrum sativum*, *Pimpinella Anisum*, are officinal under the respective names of *Fœniculum* (*Fennel*), *Carum* (*Caraway*), *Coriandrum* (*Coriander*), *Anisum* (*Anise*). They all depend for their virtues upon volatile oils which are officinal. The *Oil of Anise of commerce* is chiefly the product of a Chinese tree, the *Illicium Anisatum*, or *Star Anise*, from whose five-to ten-rayed capsular fruit it is obtained by distillation. The *Aqua Fœniculi* and *Spiritus Anisi* are officinal. All of these fruits and their preparations may be used as carminatives and stomachics.

The herbal portions of the following mints are officinal: *Lavandula vera*, *Rosmarinus officinalis*, *Salvia officinalis*, *Mentha piperita*, *Mentha viridis*,

Melissa officinalis. They are respectively known as *Lavender* (*Lavandula*), *Rosemary* (*Rosmarinus*), *Sage* (*Salvia*), *Peppermint* (*Mentha piperita*), *Spearmint* (*Mentha viridis*), and *Balm* (*Melissa*). The most important preparations of them are as follows: *Spiritus Lavandulæ Compositus*—*Compound Spirit of Lavender*, a very elegant and agreeable stomachic and cordial. Dose, a fluidrachm to half a fluidounce. *Aqua Menthæ Piperitæ*—*Peppermint Water*, and *Aqua Menthæ Viridis*—*Spearmint Water*, both very frequently used as vehicles. *Spiritus Menthæ Viridis* and *Spiritus Menthæ Piperitæ*—*Essence of Spearmint* and *Essence of Peppermint*, used as carminatives, in doses of from ten to twenty drops. The oils of lavender, rosemary, peppermint, and spearmint are also officinal, and may be used in doses of from three to ten drops as carminatives. Sage contains quite largely of tannin, and its officinal infusion is sometimes used as a gargle.

The *Volatile Oil of Gaultheria* (*Oleum Gaultheriæ*) is frequently used to flavor preparations, on account of its very pleasant taste. *Calamus*, the rhizome of the *Acorus Calamus* or sweet flag, also contains a volatile oil, and is included in the secondary list of the United States Pharmacopœia.

MINERAL TONICS.

FERRUM—IRON. (Fe.)

As iron constitutes a necessary integrant portion of the red blood-corpuscles, it is a food rather than a medicine. A very large proportion of the various articles of ordinary diet contain a trace of it; and as, according to Quevenne, it is very rarely to be found in normal urine, it must accumulate in the system. When, however, the supply, from any cause, is insufficient, or the power of digesting and assimilating it is impaired, or an abnormal excretion of it occurs, or by direct discharge, as in hemorrhage, it is drawn off from the blood, a less than normal amount of it exists in the body. When this is the case, the condition ordinarily known as *anæmia* existing, the circulating fluid contains comparatively few red disks. It is this state which constitutes the great indication for iron. Under its use, if the cause of the *anæmia* be prevented from operating continuously, the color returns to the cheeks, because there is an absolute increase of the hæmato-globulin in the blood. Thus, Prof. Simon (*Animal Chemistry*, London, 1845, Sydenham Society edition) reports a case of chlorosis in which, under the steady use of iron for sixty-four days, the globulin increased from 30.86 parts to 90.80 parts per thousand, and the hæmatin from 1.431 parts to 4.598 parts per thousand. This increase is certainly due in part simply to the supply of the peculiar food of the red corpuscles; but it seems probable that the iron acts not merely as a food, but also as a veritable stimulant to the organs which produce these bodies. Since, however, we have no definite knowledge as to the early history of the red disks, it is evident that any theory as to the method in which iron

increases their number must be a mere conjecture. When given to healthy subjects, iron increases to some extent the number of the corpuscles, and produces a degree of plethora; but after a time the blood appears, as it were, to become saturated with it and ceases to assimilate it. Thus, Brücke found that, when administered continuously to rabbits, after a certain period the ingested iron began to escape from the kidney so rapidly that it could be recovered with but little loss.

It appears to be a well-established fact that one of the functions of the red blood-corpuscles is to convert oxygen into ozone, which is the efficient form of the element in the system (see paper by A. Sasse, *Vierteljahresschrift für Prakt. Heilkunde*, 1866, zweiter Band). The oxide of iron outside of the body certainly possesses an ozonizing power similar to that of the red disk. Thus, a spot of iron mould, *i.e.*, iron oxide, on linen will in time destroy the fabric. The reason of this is the corroding action of the ozone which is slowly generated by the oxide of iron. From a similar cause a fleck of rust on a bright surface of steel will steadily enlarge and deepen. It would seem *a priori* probable that in the blood iron acts as it does out of the body. If this be so, by increasing oxidation an increase of the iron in the blood should cause elevation of temperature and increased elimination of urea. The studies of W. Pokrowsky (*Virchow's Archiv*, Bd. xxii.) have shown that, in cases of anæmia, after the exhibition of iron the temperature does rise, even when in the beginning it was not below normal, and that simultaneously there is an increase in the daily elimination of urea; and the experiments of Botkin, as quoted by Sasse (I have not seen the original), establish the same fact in regard to healthy men. The increased oxidation cannot be due simply to an increase in the number of the red corpuscles, for whilst the latter accrues slowly, Pokrowsky found that the temperature sometimes rose within five hours after the exhibition of the first dose. It would seem, therefore, that iron acts directly in the blood as an ozonizing agent.

Iron in very minute quantity (Quevenne, *Mémoire sur l'Action physiologique et thérapeutique des Ferrugineux*, *Archives de Physiologie, de Thérapeutique et d'Hygiène*, Oct. 1854, p. 93) is at times present in normal urine, and the exhibition of it as a medicine increases the percentage. According to the researches of Quevenne (p. 95), the elimination of the ordinary medicinal salts of iron is always very slight; but the experiments of Becquerel (*Simon's Chemistry*, vol. ii. p. 264) are not in accord with this. The amount of iron in the urine was found by him to vary, in those taking ferruginous preparations, very greatly and unaccountably from day to day, but the elimination commenced as soon as the administration. According to Quevenne, the haloid salts escape in the urine much more freely than the other preparations.

Almost all of the preparations of iron are more or less astringent, and when in the blood very probably they exert a direct influence upon the tissues, contracting them not merely by increasing their tone, but also by acting on their vital contractility.

The preparations of iron may be divided into those which are soluble and those which are insoluble in water. At first sight it would appear that the former class of preparations would be those most readily absorbed. The experiments of Quevenne (*loc. cit.*) have, however, demonstrated that these soluble preparations are largely precipitated by the gastric juice even when it is strongly acid. This precipitate is probably an albuminate, mixed, when the gastric juice is alkaline, with the oxide of the metal. It has been further demonstrated by Quevenne that these precipitates do not yield so large a percentage of material to the acid gastric juice as do some of the insoluble salts. According to Mialhe (*Chimie appliquée*, Paris, 1856), after the entrance of an iron salt into the blood its power of acting as a ferruginous tonic depends upon its capability of being decomposed by the alkalies of the blood in such a way as to give origin to albuminates. From these considerations it would appear that the question of solubility in water is of very little importance in the choice of a chalybeate. Quevenne has demonstrated that the reduced iron yields the largest percentage, and, as it is nearly free from astringency, it is probably the best chalybeate. If a soluble preparation be desired, the ammonia or potassa tartrates are very unirritating. If Mitscherlich be correct in his deductions, the protosalts are more readily absorbed than the sesquisalts, and consequently should be preferred as a general rule.

When iron is exhibited in the usual doses, but a small proportion is absorbed, the remainder escaping with the fæces, to which it imparts a black color. This black color is due to a conversion of the iron into tannates and sulphurets; the tannic acid of the first compound being derived from the food, the sulphur of the second from the intestinal gases.


THERAPEUTICS.—The chief indication for the use of iron is the existence of *anæmia*; the contra-indication, a state of *plethora*: on these points sufficient has already been said, and the peculiar actions of certain salts will be considered under their respective preparations. The preparations of iron which are recognized by the Pharmacopœia are very numerous, much more so than there is any reason for.

FERRUM REDACTUM—REDUCED IRON.

This preparation is also known by the names of *Ferri Pulvis*, *Iron by Hydrogen*, and *Quevenne's Iron*, and is as a chalybeate the best of all the various medicinal forms of iron. It is made from the subcarbonate (sesquioxide) by exposing it at a white heat to the action of hydrogen, which takes away the oxygen and leaves the pure metallic iron. It occurs as a light, iron-gray, tasteless powder, which should be completely dissolved by diluted sulphuric acid without yielding the odor of sulphuretted hydrogen, and when touched with a lighted taper should ignite and burn to the brown oxide of iron. If it be black, or if it fail to answer the tests given above, it is impure, and indeed, as offered in the shops, not rarely it is entirely spurious. Of all the efficient preparations of iron it is the freest from astringency. The dose is

from two to five grains, taken in pill form, after meals. As it is entirely tasteless, it is frequently given to children in chocolate-drops or lozenges, which they take without suspecting that they contain medicine.

FERRI OXIDUM HYDRATUM—HYDRATED OXIDE OF IRON.



The *Hydrated Sesquioxide of Iron*, made by precipitating the tersulphate with ammonia, is a reddish-brown powder, which is used solely as an antidote to arsenic. For this purpose it should be freshly prepared, and should be so moist as to constitute a magma. Its virtues are deteriorated by age, even when it is kept under water, and are entirely destroyed by drying. If the solution of the tersulphate of iron be not at hand in an emergency, the perchloride will yield just as useful a product, and carbonate of sodium may be substituted, if circumstances necessitate it, for the ammonia. The precipitate falls at once, and may be washed by putting it in a piece of muslin, squeezing out the original fluid, and then pouring on some fresh water. As the sesquioxide of iron is perfectly innocuous, it should be very freely administered when used as an antidote, especially since it only acts when in excess. A teaspoonful to a tablespoonful of it should be stirred up in water and taken at once, the dose being repeated several times if necessary.


FERRI SUBCARBONAS—SUBCARBONATE OF IRON.

A reddish-brown tasteless powder, which is made by precipitating the sulphate of iron with the carbonate of sodium. The greenish carbonate of the protoxide is first thrown down, but in drying this absorbs oxygen, gives off almost all of its carbonic acid, and is converted into the anhydrous sesquioxide. This preparation is a feeble chalybeate, rarely used. It has been recommended in very large doses (twenty grains) as a sort of specific in *neuralgia*.

PILULA FERRI CARBONATIS—PILL OF THE CARBONATE OF IRON.

This blackish-green pilular mass is made by precipitating a syrupy solution of the sulphate of iron by carbonate of sodium. Throughout the subsequent washings, and during drying, sugar is kept constantly present in large amount, so as to prevent the absorption of oxygen and consequent conversion of the protocarbonate into the sesquioxide. This is a very good chalybeate, nearly free from astringency, and may be given in doses of from three to five grains, in pill form. *Pilulæ Ferri Compositæ* contain the protocarbonate of iron and myrrh. They are sometimes used in anæmia with amenorrhœa; one to three pills three times a day. The *Mistura Ferri Composita* is a liquid preparation containing substantially the same ingredients as the compound pills. Dose, one to two tablespoonfuls.

FERRI SULPHAS—SULPHATE OF IRON.



The pure protosulphate of iron is made by dissolving iron in dilute sulphuric acid. It occurs in transparent efflorescent rhombic prisms of a pale bluish-green color and a metallic styptic taste. The sulphate of iron is a very decided astringent, and in a concentrated form and sufficient amount acts as an irritant poison, producing vomiting, purging, and gastro-intestinal inflammation. Externally its solution (five to twenty-five grains to the fluidounce) is used as an astringent lotion. It has been especially recommended in this form (5i to Oj) in *erysipelas*. As a simple chalybeate the sulphate of iron should never be used. In *chronic diarrhœa* it is sometimes employed as a tonic astringent. Dose, five grains; in the form of the dried sulphate (*Ferri Sulphas Exsiccata*), three grains.

LIQUOR FERRI SUBSULPHATIS—MONSELL'S SOLUTION.

The solution of the subsulphate of the sesquioxide of iron is a most powerful astringent and styptic, and is used solely as such. It is but slightly irritant, and is the best of all the astringents for staunching hemorrhage when it can be applied directly to the part, as in external wounds, or in *hæmatemesis*. In hemorrhage from the stomach, ten drops of it may be given in a table-spoonful of water, the dose being repeated as necessary. Applied by means of the atomizer, it is often very efficient in *hæmoptysis*. In such a case the fluid to be atomized should have the strength of from five to twenty drops to the ounce. The inhalation should last from five to fifteen minutes, and be repeated at intervals of an hour or longer. It is very essential that the liquid should be finely pulverized.

The *Solution of the Tersulphate of the Sesquioxide of Iron* is also officinal, under the title of LIQUOR FERRI TERSULPHATIS. Owing to its irritant action, it is used only in pharmacy to make the sesquioxide preparations.

TINCTURA FERRI CHLORIDI—TINCTURE OF THE CHLORIDE OF IRON.

This preparation contains the sesquichloride of iron, muriatic acid, and alcohol, and, from the reactions of the last two ingredients, muriatic ether. It is prepared by adding alcohol to the officinal *Liquor Ferri Chloridi*, and is a reddish-yellow, astringent, irritating, somewhat corrosive liquid. It is an excellent chalybeate, possessed of peculiar properties, probably in some measure due to the ether which it contains. It is a diuretic, increasing often very decidedly the daily urinary secretion. It appears also to exert an astringent influence upon the urino-genital mucous membrane, and is frequently employed with tincture of cantharides in *gleet*. In *chronic Bright's disease* it is often of very great service as a chalybeate diuretic. In *erysipelas* it is constantly employed with remarkable results, controlling the disease in a

manner not yet understood. Analogy has suggested its employment in other adynamic affections, such as *diphtheria* and *pyæmia*, but its value in these diseases is much more doubtful.

Locally it is used as an astringent in sore throat; for this purpose its strength should be reduced at least one-half. As it is very destructive to the teeth, care ought to be exercised in its use about the mouth, and also in its administration. The dose is from fifteen to thirty drops, given as a chalybeate three times a day; in erysipelas every two or three hours. The orange-yellow crystalline deliquescent *Sesquichloride of Iron* (FERRI CHLORIDUM) ($\text{Fe}_2\text{Cl}_3 - \text{Fe}_2\text{Cl}_6$) is officinal, but is very rarely used.

There are four officinal *Citrates of Iron*, all soluble in water. Two of these (FERRI CITRAS and FERRI ET AMMONII CITRAS) occur in garnety-red scales, and are simply mild chalybeates. Dose, five grains. The *Citrate of Iron and Quinia* (FERRI ET QUINIE CITRAS) is in transparent scales, varying from reddish brown to yellowish brown in color, with a tint of green. The *Citrate of Iron and Strychnia* (FERRI ET STRYCHNIE CITRAS) contains one per cent. of strychnia.

There are two officinal *Tartrates of Iron* (FERRI ET AMMONII TARTRAS and FERRI ET POTASSII TARTRAS), both occurring in garnety scales, and soluble in water. Dose, five grains.

The *Lactate of Iron* (FERRI LACTAS) occurs in greenish-white crystalline crusts or grains, soluble in forty-eight parts of water. It is a good chalybeate. Dose, five grains.

There are two officinal *Phosphates of Iron*, which may be used in five-grain doses. FERRI PHOSPHAS is a bright slate-colored insoluble powder. FERRI PYROPHOSPHAS is an excellent preparation, occurring in apple-green scales, somewhat soluble in water, and nearly free from ferruginous taste.

The officinal *Oxalate of Iron* (FERRI OXALAS), a yellow powder, and *Prussian Blue* (FERRI FERROCYANIDUM), are neither of them eligible preparations.

The *Ammonio-ferric Alum* (FERRI ET AMMONII SULPHAS) occurs in octahedral crystals, of a pale violet color. It is freely soluble in water, is very astringent, and is only used in atonic *leucorrhæa*, in which affection marked benefit is often gained by the exhibition of it in five-grain doses three times a day.

ACIDUM SULPHURICUM—SULPHURIC ACID.

Oil of Vitriol ($\text{SO}_3\text{H}_2\text{O} - \text{H}_2\text{SO}_4$) is when pure a colorless, heavy liquid (sp. gr. 1.843). On exposure to the air it rapidly absorbs moisture and becomes less dense. When its specific gravity is 1.78 it deposits crystals of the bihydrated acid at 28°F. and may burst the glass in which it is kept. Owing to its great affinity for water, it is used in chemistry as a desiccant.

PHYSIOLOGICAL ACTION.—Concentrated sulphuric acid is powerfully corrosive both of animal and vegetable tissues, abstracting the elements of

water and leaving the carbon untouched. It consequently *blackens* organic matter at the same time that it destroys its texture. When administered in therapeutic doses and absorbed into the blood, it is converted into sulphates, and as such, so far as is known, is eliminated.

It escapes by the kidneys, as has been proven by Dr. H. Bence Jones (*Lectures on Pathology and Therapeutics*, London, 1867) in regard to large therapeutic doses, and is attested by Maukopff as occurring after poisoning. As, however, the amount of elimination by the kidneys seems to be slight, it is very probable, as Dr. Headland (*Action of Medicines*, London, 1852) conjectures, that it is excreted both by the lower bowel and by the skin. Locally applied, dilute sulphuric acid is an astringent, and clinical experience proves that it exerts a similar action when taken into the system. Under such circumstances its astringent influence is most marked upon the skin and intestine,—parts which are believed to excrete it: it is therefore possible that this action is in a measure local, and dependent upon the presence of the excreted acid.

THERAPEUTICS.—Concentrated sulphuric acid is not rarely used as an escharotic, for which purpose it is mixed with finely-powdered charcoal so as to form a paste. Appropriately diluted, it has been used as a stimulant and astringent lotion in *venereal* and other indolent *ulcers*.

Internally, sulphuric acid is very useful as an astringent in *colliquative sweats* (*night-sweats*) and in profuse *serous diarrhœas*. I have used it with great advantage in the sudden serous vomiting and purging of infants known as *cholera infantum*.

It has been employed with advantage in *cholera*, and a remarkable series of observations by Dr. R. G. Curtin (*Philadelphia Medical Times*, vol. iii. p. 649) at least furnish good reason for further testing its powers as a *prophylactic* against this disease. The facts recorded by Dr. Curtin are as follows. A very severe epidemic of the disease ceased in the Insane Department of the Philadelphia Almshouse within twelve hours after the lunatics were all put upon the free use of sulphuric acid lemonade; the only new case after this being in a man who refused to use the prophylactic. Two days after the use of the sulphuric acid was stopped, two new cases occurred, the epidemic being again arrested by the use of the acid. In the surgical wards of the Hospital Department the acid was used from the beginning of the epidemic, and these wards, although in no way isolated, were the only parts of the institution unvisited by the disease.

Sulphuric acid was formerly used in *hemorrhages*, but is now rarely employed. It is, I think, much less efficacious than some other remedies.

In *acute lead-poisoning* dilute sulphuric acid is an efficient antidote, and it has been proposed by M. Gendrin (Dr. Bennett, *London Lancet*, 1856) as a remedy in *chronic lead-poisoning*. As, however, he combined its internal exhibition with the daily use of warm sulphur-baths, it is doubtful how much of the successful result was due to its action. It is difficult to perceive how

it can do good in these cases, and I do not think the clinical proof that it does do good has as yet been brought forward.

TOXICOLOGY.—When swallowed in concentrated form, sulphuric acid acts as a corrosive. Death from consequent collapse has resulted in two hours and a half (Case *Medical Times and Gazette*, vol. i., 1863), but usually the course of sulphuric acid poisoning is much more protracted, the fatal result having been delayed in some cases for several months. The usual symptoms are pain in the mouth, throat, and epigastrium, violent vomiting, often of tarry matters, with symptoms of collapse, such as cold extremities, feeble pulse, suppressed voice, and clammy skin. The mind is usually clear to the last. Profuse, and sometimes bloody, salivation is commonly present. The parotids sometimes swell as early as the fourth day, and Maukopff (*Sydenham Year Book*, 1863) has seen suppurative parotitis apparently induced by closure of the duct of Steno, retention of secretion, and consequent irritation of the gland. The later symptoms are those of ulceration of the œsophagus and stomach, and need not be dwelt upon here. The larynx is often profoundly affected. There is a very marked increase in the amount of sulphates in the urine, which may be albuminous and contain granular casts. Desquamative nephritis may be developed several days after the subsidence of the first symptoms. Thus, in one of the cases reported by Maukopff, urine which had ceased to be albuminous on the third day became so again on the twentieth, with a simultaneous development of casts containing blood-corpuscles; after death tubular nephritis was found. Another symptom noted by Maukopff was intercostal neuralgia.

On post-mortem, greater or less destruction of the œsophagus and stomach, or of the air-passages, is found. The black color of the slough is a diagnostic sign which an examination of the mouth will sometimes render available before death.

The most important part of the treatment of sulphuric acid poisoning consists in the immediate and free exhibition of the antidotes, which should be given in milk or in water. The best antidotes are chalk, magnesia, white-wash, and soap. Christison condemns the use of the alkaline carbonates, because they are themselves irritating. As, however, in these cases time is a matter of vital importance, if the alkaline carbonates be the only antidotes at hand they should be used unhesitatingly.

ADMINISTRATION.—Sulphuric acid should of course be given properly diluted, and with the requisite precautions to prevent its injuring the teeth. It is best administered in one of the following preparations :

Acidum Sulphuricum Dilutum—*Dilute Sulphuric Acid* (℥ii to Oj). Sp. gr. 1.082. Dose, ten to thirty drops.

Acidum Sulphuricum Aromaticum—*Aromatic Sulphuric Acid* (℥iii to ℥xix). A dark reddish-brown liquid, containing alcohol and aromatics. Dose, ten to twenty drops.

ACIDUM MURIATICUM—MURIATIC ACID. (HCl.)

Muriatic Acid is a colorless aqueous solution of hydrochloric acid gas, having the specific gravity of 1.160, and containing rather more than 33.9 per cent. of the gas. The muriatic acid of commerce commonly has a yellowish tint, produced by sesquichloride of iron or sometimes by organic matter.

PHYSIOLOGICAL ACTION.—In its concentrated form muriatic acid is highly corrosive, but less so than either sulphuric or nitric acid. Its astringent properties are not all decided, if indeed it really possess any. As it probably is one of the natural acids of the stomach, it would seem as though it ought to be capable of aiding digestion. It also appears to have influence over the glandular system of the alimentary canal, increasing by its action their normal secretions.

THERAPEUTICS.—In *stomachic dyspepsia*, muriatic acid, with or without pepsin, is often useful by aiding in the digestion of the food. In other cases, where the *dyspepsia* is *intestinal* with a tendency to diarrhoea and loss of appetite, muriatic acid combined with strychnia and some aromatic, such as compound tincture of cardamom, is often very advantageous.

In *low fevers* the use of mineral acids has been highly extolled by various authorities. I have seen a number of cases treated upon this so-called "Swedish plan," and have never been able to perceive that the acids do any real good.

Locally, diluted muriatic acid has been recommended by Bretonneau in *diphtheria*. He employed a mixture of one part of the acid to two of honey, but bolder practitioners have used it of full strength, with, it is claimed, good effect. It should be applied by means of a little mop, scrupulous care being exercised to prevent any of the acid coming in contact with parts not protected by false membrane.

TOXICOLOGY.—Muriatic acid, as a poison, is similar in its effects to, but less powerful than, sulphuric acid, recovery having occurred after the ingestion of an ounce of the officinal acid (*Boston Medical and Surgical Journal*, vol. xv.). The treatment is similar to that of poisoning from other mineral acids.

ADMINISTRATION.—The acid is best given in the form of the officinal *Acidum Muriaticum Dilutum* (3iv-Oj). Sp. gr. 1.038. Dose, ten to thirty drops properly diluted.

ACIDUM NITRICUM. (NO₅,HO—HNO₃.)

A liquid of the specific gravity 1.420, which as first made is colorless, but by exposure to the light acquires a yellow tint. It oxidizes all of the common metals except gold, and is exceedingly corrosive to living tissue, which it stains an indelible yellow. When diluted it converts most animal and vegetable substances into oxalic, malic, or carbonic acid.

PHYSIOLOGICAL ACTION.—When applied to any portion of the living

organism, nitric acid acts as a most powerful chemical caustic. When very greatly diluted, it is a simple local stimulant, with very slight astringent powers. Owing to its chemical activity, its vapor was at one time used as a disinfectant; but it has been superseded by other substances. Taken internally in small amount, it acts as a stimulant upon the glandular system of the alimentary canal, and in serous diarrhœa appears to exert an astringent influence. It seems to me very probable that these effects are local rather than constitutional, due to a direct action of the acid upon the mucous membrane of the bowel. On entering the blood, nitric acid must be converted into a nitrate. In regard to its elimination we have but little definite information, but it probably escapes through the same channels as does sulphuric acid.

THERAPEUTICS.—Nitric acid is used quite frequently as an escharotic, especially in cases of *chancres* and *venercal* or other *warts*. In its employment care should be taken to protect the sound tissue by oil, or, still better, by a layer of soap. It may be applied by means of a splinter of wood, or, if it is to be used more freely, by a little mop. When it has penetrated as deeply as is desirable, washing the part with warm soapsuds will prevent further action. Very much diluted (five to twenty drops to the ounce), it forms a good stimulant lotion for indolent *ulcers*. It should not be employed as a mouth-wash, on account of its destructive action on the teeth. Of course this does not apply to its use as a caustic in *canerum oris*, in which, as in other forms of *acute gangrene*, such as *phagedenic chancres* and *hospital gangrene*, it is very efficient. In these cases, to do good it must be applied thoroughly.

Internally, nitric acid has been used in *low fevers*, but with, I think, doubtful advantage.

In *dyspepsia*, in *chronic hepatic congestion*, in the *oxalic acid diathesis*, in the *dyscrasia* of *constitutional syphilis*, nitric acid has been employed with advantage, but is, I think, much inferior to the nitro-muriatic acid.

In 1826 Dr. Hope claimed for the *Acidum Nitrosum* a specific action in *serous diarrhœa*, including the sudden acute diarrhœas of hot climates, and in the chronic *dysenteries* originating under similar circumstances. The formula he employed is as follows:

R Acidī nitrosi, fʒi;
Misturæ camphoræ, fʒviii;
Misce, et adde Tinct. opii, gtt. xl.

S.—A fourth part to be taken every three or four hours.

Under the name of *Hope's Camphor Mixture* a preparation similar to this has been much used, but has gradually lost the confidence of the profession, chiefly, I believe, because on theoretical grounds the original formula has been departed from. *Nitrous Acid* of the shops (*Acidum Nitrosum*, Edinburgh Pharmacopœia) is an orange-red liquid, which may be looked upon as a solution of nitric oxide in nitric acid. When it is diluted with water it is after a short time converted into simple nitric acid. For this reason it has

been customary to substitute nitric acid for the *Acidum Nitrosum* of Hope's original formula. It should be noted, however, that the latter only provided sufficient of the remedy to last a few hours, and, as the reaction which has been spoken of requires some time for its performance, I do not think that theory in truth warrants the change. Practically I have failed absolutely with the new formula, when immediate relief was afterwards obtained by the use of the medicine prepared according to the old plan. Made in this way and used whilst fresh, Hope's Camphor Mixture is a very efficient though disagreeable remedy in diarrhoeas connected with disordered secretion of the liver and other glands of the alimentary canal.

ADMINISTRATION.—The dose of the strong acid is from five to twenty drops; of the diluted (*Acidum Nitricum Dilutum*, ℥iii to Oj, sp. gr. 1.068) is from fifteen to sixty drops.

TOXICOLOGY.—When taken in a concentrated form, nitric acid is a corrosive poison, exceeding even sulphuric acid in violence. The symptoms so closely resemble those induced by the latter substance that it is unnecessary to detail them here, as is also true of the treatment. The distinguishing character is to be found in the *color* of the affected tissues, which in nitric acid poisoning are stained of a deep yellow.

ACIDUM NITROMURIATICUM—NITRO-MURIATIC ACID.

This preparation is made by mixing three parts of nitric with five parts of muriatic acid. If the acid be sufficiently strong, an orange-colored liquid is formed, with the evolution of intensely-irritating vapors. After standing for a length of time, the red color changes to a golden-yellow. It is in this state that the United States Pharmacopœia directs the acid to be used. By longer standing the *golden-yellow* becomes *lemon-yellow*, and the odor of chlorine is almost entirely lost. These changes are hastened by light, but will occur in the dark and in well-stoppered bottles. Although the golden-yellow acid is directed by the Pharmacopœia, yet careful clinical studies have convinced me that the acid acts much more efficiently when freshly prepared and of a deep-red color. In some cases it has seemed to me useful only when in the latter form. The lemon-yellow acid is nearly valueless.

PHYSIOLOGICAL ACTION.—In concentrated form nitro-muriatic acid is exceedingly corrosive. Our knowledge of its action in small doses is purely clinical, and will be spoken of under the head of Therapeutics. The chemistry of the acid is so complex and uncertain that no reasonable conjecture can be made as to the form in which it is absorbed or the method of its escape from the body. That it is absorbed in some form is proven by its occasionally producing salivation, even when applied to the skin by means of baths.

THERAPEUTICS.—The remedial value of nitro-muriatic acid depends chiefly upon the power which it possesses, to a much greater degree than any other of the mineral acids, of influencing the action of the liver and other glandular organs of the alimentary canal. Originally proposed by Dr. N. Scott, of

Bombay, in the *chronic hepatitis* of hot climates, it has been used with great success by Annesley, Martin, and other famous India surgeons. The remedy would seem not to be indicated in hepatitis with high fever and a tendency to rapid suppuration so much as in the slower form of the affection, which normally ends in chronic enlargement and induration of the viscus. In the habitual *congestion* of the *liver* occasionally seen in this climate I have used it with the most marked benefit.

In the still milder affection known as "*biliousness*," whose pathology is probably a torpid condition of the small glands of the alimentary mucous membrane as well as of the liver, nitro-muriatic acid has yielded in my hands most excellent results. That the remedy does act upon the liver is proven by the fact that in these cases it sometimes produces violent bilious diarrhœa. When *jaundice* depends upon obstruction or upon any of the severer organic diseases of the liver, the acid is of little if any use; when, however, the jaundice depends upon torpor of the liver, or even when it is catarrhal in origin, the remedy may be of great service. Even in the early stages of *cirrhosis*, whilst the liver is still enlarged, nitro-muriatic acid should be tried, as in some cases apparently of this character great benefit has been derived from its use.

In those forms of *chronic diarrhœa* in which the disease is really an intestinal dyspepsia, nitro-muriatic acid may be of the utmost service, benefiting and even curing cases which have resisted all other treatment. As the effect of the acid is not a sudden one, it is evident that it acts in these cases not as an astringent, but by restoring the normal digestive power.

There is a morbid condition, probably dependent upon defective primary assimilation, in which the chief symptoms are general malaise, a feeling of weakness, a lack of elasticity, a very great depression of spirits, in which crystals of oxalate of lime are generally to be found in the urine, and in which nitro-muriatic acid produces in a few days a surprising revolution.

As a "blood-purifier" the acid has been employed in *constitutional syphilis*, and in various ulcerative *skin-affections*. In these diseases it no doubt does good by improving digestion and increasing glandular action, but there is no reason to believe that it is a direct alterative.

ADMINISTRATION.—For reasons which have already been given, when nitro-muriatic acid is administered internally it should be freshly prepared; and as the changes which have been spoken of take place more rapidly when the acid is mixed with water, the officinal *dilute* nitro-muriatic acid is an ineligible preparation. As light hastens its deterioration, the strong acid should always be kept in a dark bottle with a glass stopper. Directly after mixing the acids the evolution of gas may be so great as to necessitate its being allowed to escape. After six or eight hours, however, the bottle should be closely stoppered. The dose of the strong acid is from five to eight drops, properly diluted, and taken through a tube after meals.

In *chronic hepatic diseases* the external application of the acid seems to

give even better results than its internal use. In India, according to Sir Ranald Martin, the bath is used as follows: 1. Take Hydrochloric Acid $f\text{℥}iii$, Nitric Acid $f\text{℥}ii$, Water $f\text{℥}v$. Mix. Two gallons of water with six fluidounces of the above mixture suffices for a bath, which will keep fit for use during three days, provided half a fluidounce of acid and a pint of water are added morning and evening. The bath must of course be given in wooden or earthenware vessels, and if it becomes necessary to warm it only a portion should be heated, and the rest then added. In urgent cases the whole body may be immersed in the bath, but generally a foot-bath is preferable, the inside of the thighs and arms and the hepatic region being at the same time sponged. The bath should be repeated twice daily, lasting each time for ten or fifteen minutes.

I have had no experience in this method of using nitro-muriatic acid; but have derived great benefit from the application of the acid over the hepatic region. My plan has been to wring out a large piece (eight by ten inches) of spongio-piline, or of canton-flannel (several layers), in a lotion of a strength varying, according to the irritability of the patient's skin, from one to three fluidrachms to the pint, and to apply this over the right hypochondrium, covering it with a piece of oiled silk supported by a bandage. The application sometimes causes a prickling sensation, and after a time may produce a profuse local sweating. The dressing may be left on from half an hour to an hour, and be repeated three or four times a day; some patients can wear it almost continuously.

TOXICOLOGY.—When taken in poisonous doses, nitro-muriatic acid produces symptoms and results similar to those following the ingestion of nitric acid. The color of the stain produced by it is yellow, and its antidotes are the same as those of other mineral acids.

PHOSPHORUS. (P.)

Phosphorus is a translucent, when pure nearly colorless, but usually slightly yellowish, highly inflammable elementary body, which is tasteless, but possessed of a peculiar alliaceous odor. It is obtained from the phosphate of lime of calcined bones, by taking away the lime with sulphuric acid and deoxidizing the residuum by heating with charcoal. It is insoluble in water, sparingly soluble in ether, absolute alcohol, and the oils, freely so in chloroform. It takes fire at 100° F., and melts at 108° F. In the shops it is in cylindrical sticks, covered with a whitish layer, and having when cut a waxy consistence and lustre. When properly treated, it is converted into several allotropic forms, the red phosphorus, black phosphorus, and the crystallized metallic phosphorus of Hittorf. The first of these is the most important; it is brittle, does not take fire at ordinary temperatures, and is said not to be poisonous.

PHYSIOLOGICAL ACTION.—The physiological action of phosphorus in therapeutic doses is probably entirely different from that which it exerts when in larger amounts. It is a constituent of most of the more important tissues,

and is especially abundant in the nerve-centres. Like iron, cod-liver oil, etc., it appears to act when given in minute quantity as a stimulant to the nutrition of the tissues, into whose composition it enters. So far as the nervous system is concerned, this assertion rests exclusively upon clinical observation; but Dr. Wegner (*Virchow's Archiv*, June 22, 1872) has experimentally demonstrated such an action upon the bony tissues. He found that when adult animals were fed upon minute doses of phosphorus the spongy tissue in the long and short bones was thickened, and the compact tissue rendered more dense. After a time new tissue was deposited upon the inside of the shafts of the long bones, and this went on in some instances until the marrow-cavity was obliterated. The action upon the bones of growing animals was even more marked.

Phosphorus was at one time believed to be a diffusible stimulant, and it possibly may exert such an influence upon the nervous centres, especially those of sexual life. I have, however, never seen any evidences of such action.

For reasons to be hereafter adduced, it is certain that in poisonous doses phosphorus acts as phosphorus, and when it is administered therapeutically it certainly enters the blood in its elemental form, and, I believe, acts as such. Dr. Wegner advances the following reasons for believing that it does not act as phosphoric acid so far as the bony tissues are concerned: First, no similar action can be obtained from phosphoric acid unless from eight hundred to one thousand times the proportional dose be given. Second, the newly-formed tissue is at first gelatinous. Third, there is no excess of phosphates in the bone. Fourth, when the food is deprived of lime the same new tissue arises, but remains in a soft gelatinous state.

THERAPEUTICS.—The chief use of phosphorus in medicine is as a nutrient tonic to the nervous system. In all cases of *nervous exhaustion*, whether involving the cerebral or spinal centres, it is of great value. I have seen marked benefit from its use when the symptoms were not severe enough to indicate organic lesion, but the most remarkable results have been in the cases in which the structure of the centres was apparently deeply implicated. In threatening *cerebral softening*, in *myelitic paraplegia* from excessive venery, it is the only drug which appears really to affect the nerve-centres.

In *neuralgia* attention has recently been drawn by several writers to its virtues; and as neuralgia is often simply an expression of exhausted nerve-power, the use of phosphorus is commended by reason as well as by experience.

It is probable that it may be of value in some cases of impaired vitality, although the nervous system be not obviously implicated. Dr. H. Eames (*Dublin Journal of Medical Sciences*, Jan. 1872) states that he has obtained great benefit from its use in obstinate skin-affections, such as *lupus*, *acne*, and *psoriasis*. Attention has been recently called abroad to its use in *cataract*.*

* Dr. Tavignot, *Revue de Thérapeutique Médico-Chirurgicale*, August and September, 1871; Prof. Gioppi, *Giornale d'Oftalmologia*, abstract in *N. Y. Medical Record*, 1872.

On account of its marked influence on the development of bone, Dr. Wegner suggests its use in *osteo-malacia* and in *rickets*, but as yet there has been no clinical experience with the drug in these disorders.

Prof. Samuel R. Percy has used a preparation of phosphorus (see Administration) with great advantage in cases of repeated *furuncular* eruptions (*Prize Essay, Transactions of the American Medical Association*, 1872, p. 659).

TOXICOLOGY.—The ingestion of a fatal dose of phosphorus is not followed by any sensible effects for some time. After, however, from three to twelve hours a sense of weakness and of general wretchedness manifests itself, and in a large proportion of the cases (according to Lewin eighty-eight per cent.) is accompanied, or soon followed, by vomiting. With the emesis there is nausea, and in most cases the patient complains of abdominal pain, the severity of which, however, never equals that of corrosive poisoning. The matters vomited consist of food, mucus, and bile. During the first eight or ten hours they often smell strongly of phosphorus, and are luminous in the dark. The vomiting may persist during the whole attack, but generally ceases on the second or third day, to reappear with the subsequent jaundice, when coffee-colored vomit from exuded blood is ejected. The pain, which in most cases abates with the vomiting, often spreads from the epigastrium over the whole abdomen, and in rare instances is paroxysmal. If it reappear in the latter stages, it is apt to affect especially the right hypochondrium, and is associated with decided tenderness in the region named and in the epigastrium.

The tongue is whitish or abnormally red, sometimes furred. There is generally fever, loss of appetite, and thirst. Maukopff has noted a morning and evening temperature of from 37° C. to 39° C. and from 37.4° C. to 39.8° C. respectively. During the latter part of the case there is very often a remarkable fall in the temperature, which is generally, but not always, a precursor of death. The lowest point I have seen noted was 31.2° C. (88.2° F.) some hours before death.* In some cases fever is altogether absent or comes on just before death.†

The stools are at times normal in character and frequency, but there is generally diarrhoea or constipation, with flatulence. Late in the attack the passages are in most cases very light clay-colored, or even whitish, and exceptionally they are bloody. In some cases they are phosphorescent.

Jaundice comes on in from thirty-six hours (cases reported by Maukopff, *Wiener Medicin. Wochenschrift*, 1863, and by Tüngel, *Klinische Mittheilungen von der Medicin. Abtheilung des Allgemein. Krankenhauses in Hamburg*, 1861) to five days (Lebert and Wyss, *Archives Générales*, 6th series, xii., September, 1868) after the ingestion of the poison. In most cases it appears first

* See case of Dr. Battmann, *Archiv der Heilkunde*, 1871, p. 257.

† See case of Concato, *Sydenham Soc. Year Book*, 1869-70, p. 454.

in the conjunctiva, but sometimes the urine gives previous warning of its approach. In some cases there is with it a decided and palpable increase in the size of the liver, which may pass, if the patient live long enough, into an equally apparent lessening of the bulk of the viscus. The severe nervous symptoms are rarely if ever developed until after the jaundice, although early in the attack there is not unfrequently anxiety, headache, giddiness, and dreamy unquiet sleep, or even sleeplessness. The more pronounced nervous symptoms consist of delirium, which may be wild and is very frequently erotic, with somnolence ending in coma and death, occasionally preceded by convulsions. According to Taylor, the latter are a certain sign of approaching dissolution. Very generally, partial spasms and fibrillar contractions of the voluntary muscles occur, although there is always, in not too rapid cases, progressive paresis of the voluntary muscles. Death sometimes takes place suddenly from collapse and cardiac paralysis, but more commonly the patient dies comatose from a gradual failure of the respiration and circulation.

If recovery occur, it is by a gradual amelioration of the symptoms, and the health of the patient is apt to be impaired for some time. Apparently desperate cases will sometimes convalesce unexpectedly, and Tüنگel states that a favorable issue may take place even after violent delirium.

The urine is almost always much affected by the poison. Very commonly it is scanty, is albuminous, and sometimes it contains sugar. As was first pointed out by Munk and Leyden (*Die acute Phosphorvergiftung*, Berlin, 1865), after jaundice has set in, bile-acids, as well as biliary coloring-matter, are always to be found in the urine. Not unfrequently a cloudy sediment, consisting in part of epithelial cells, often tinged with bile, is deposited. Dr. Oswald Kohts (*Pflüger's Archiv*, Bd. iii. p. 1) and other observers have found leucin and tyrosin in the urine of dogs poisoned with phosphorus, and undoubtedly these substances are sometimes to be met with in the human excretion: indeed, Lebert and Wyss (*loc. cit.*, p. 269) stated that tyrosin has so been found in one case, and Dr. Ossikovsky (*Wiener Med. Presse*, 1872) asserts that in another case he has found both of these substances abundant in the urine. The albuminuria generally follows, but may precede, the icterus. A very remarkable and apparently constant constituent of the urine is the sarco-lactic acid.

Phosphorus induces in animals symptoms parallel with those commonly seen in man; although Kohts states that he has seldom seen albuminuria in animals, even when the structure of the kidneys was profoundly altered. Orfila, Magendie, Munk, and Leyden found that the only effects following the injection of phosphuretted oil into the jugular vein of an animal were the exhalation of phosphoric acid and broncho-pulmonary inflammation: on examination (Munk and Leyden), oil, but no phosphorus, was found in the lungs. Ludimar Hermann and Alfred Brunner (*Deutsch. Archiv für Klin. Med.*, p. 198) have shown that in these experiments there was embolic arrest of the oil in the pulmonary capillaries, and consequent inflammation with subsequent oxidation

of the phosphorus. When the phosphuretted oil is injected in the form of a fine emulsion, the characteristic symptoms and post-mortem appearances of phosphorus-poisoning result.

In some instances phosphorus-poisoning presents symptoms quite different from the typical array. Death may take place in a few hours, and in such cases jaundice is not generally present. Zeidler reports a case in which death occurred in about forty-two hours, from suppression of urine followed by collapse and erotic delirium. In a case of Bollinger's (*Deutsches Archiv für Klin. Medicin*, Bd. vi., 1870) the chief symptoms were vomiting, pain and tenderness over the abdomen, great weakness of pulse, gradually-developed paralysis of the legs, and death, without jaundice, in four and a half days. The autopsy revealed hemorrhagic effusion between the membranes and the spinal cord, and also into the sheaths of the proximal portions of the spinal nerves.

In women, fatal doses of phosphorus very commonly produce a bloody pseudo-menstrual discharge, and when pregnancy exists almost invariably induce abortion or miscarriage.

The older toxicologists spoke of an erosive gastritis as a common result of phosphorus-poisoning, but it is now well established that such affection is very rarely induced by the drug.* As was first pointed out by Virchow (*Virchow's Archiv*, Bd. xxxi., 1864), there is universally a gastro-adenitis, which causes the gastric mucous membrane to be thickened, opaque, whitish, grayish, or yellowish-white. Under the microscope the epithelial cells appear swollen and filled with granules and oil-globules, and in very advanced degeneration the cells completely break down. This gastro-adenitis is not due to a local action of the phosphorus, because it occurs when the poison is introduced through other channels than the mouth. The duodenum and intestines suffer similar changes. The liver is generally very much enlarged, friable, and light-colored; sometimes it is mottled, and sometimes portions of it are deeply stained with bile. The cells are gorged with fat-globules, and in some cases there are small-celled interstitial thickenings due to hyperplasia of the trabecular tissue. The gall-bladder may be full or empty. In protracted cases the liver undergoes atrophy, with destruction of its secreting cells. The kidneys, especially in their cortical portion, suffer a degeneration similar to that of the liver, the epithelium becoming enlarged, granular, fatty, and finally undergoing destruction. The voluntary and cardiac muscles, the spleen, the lungs, and probably all the tissues, partake of the universal fatty degeneration† which Wegner has shown to involve even the minute arterioles.

* It would appear probable that if it occurred at all it would be in rapidly fatal cases: yet Tüنگel did not find it in a patient dead in nine hours of phosphorus-poisoning.—*Virchow's Archiv*, Bd. xxx., 1864.

† The scope of the present work does not allow of a full discussion of the pathology of phosphorus-poisoning. The reader is referred to the following papers:

Études cliniques et expérimentales sur l'Empoisonnement aigu par le Phosphore. Par

The blood is often profoundly affected, becoming very dark, losing its power of coagulation, and apparently suffering also in its corpuscular elements: for ecchymoses are almost universal, and hæmatin crystals are occasionally found in the viscera. In the case of Concato (*loc. cit.*) the white corpuscles were observed during life to be increased in number, and the red to be diminished in size and altered in form. The ecchymoses occur in all parts of the body, but are apt to be especially pronounced in the mediastinum and the serous membranes.

Dr. Mayer states that when very large doses of the poison have been taken, the blood and even the urine (?) may be phosphorescent (*Constatt's Jahresbericht*, Bd. v., 1862, p. 123).

The elimination of bile-acids in the urine shows that the jaundice of phosphorus is caused not by an arrest of secretion, but by an occlusion of the biliary passages and consequent resorption of the bile. Dr. O. Kohts (*loc. cit.*) has apparently demonstrated that the occlusion is most frequently due to the duodenitis involving the common duct, so as to obliterate its lumen by the swelling of the mucous membrane. In some cases, however, it is probable, as believed by Wyss, Alter, and Ebstein, that a catarrhal inflammation of the minute gall-ducts is the cause of the jaundice, and also that the result is in part effected through pressure upon those ducts by the swelling of the glandular and trabecular tissue.* It is proper to state that Demarbaix and Willmart (*Presse Médicale*, xxi., 1869, and *Schmidt's Jahrbücher*, p. 152, Bd. cxliv.) insist that the icterus is not really heptogenous, but hæmic in origin, chiefly because they have found hæmatoidin in the urine. This fact, however, proves only that the blood is altered by the poison: it does not disprove the liver-origin of the jaundice.

Acute phosphorus-poisoning so closely resembles yellow atrophy of the liver that their clinical distinction is sometimes difficult, nay, impossible. Distinct phosphorescence in the breath, vomit, or stools would of course be direct evidence of poisoning. This phosphorescence, however, very often cannot be detected: according to Vetter (*Virchow's Archiv*, Bd. liii. p. 186) it can be rendered more evident in the vomit, stools, etc., by acidifying with sulphuric acid and warming in a shallow dish. When death ensues during the first week of phosphorus-poisoning, the enlarged liver affords a distinctive proof of poisoning; but when the case is more protracted, the atrophied liver of phosphorus cannot be distinguished from that of the natural disease. The symptoms during life rarely, if ever, afford sufficient ground for a positive

Prof. H. Lebert and Dr. O. Wyss.—*Archives Générales*, Sept. 1868 (an elaborate essay on the whole subject).

Zur pathologischen Anatomie der acuten Leberatrophie und der Phosphorvergiftung, von Dr. Otto Bollinger.—*Deutsches Archiv für Klin. Med.*, Bd. v., 1869.

Ueber Icterus bei Phosphorvergiftung, von Dr. O. Kohts.—*Ibid.*

* For an elaborate discussion of the cause of jaundice, see Kohts's Memoir, *Deutsches Archiv*, Bd. v. p. 168; consult also that of Dr. Bollinger, *Centralblatt für die Med. Wiss.*, 1869, and *Deutsches Archiv*, Bd. v., 1869.

diagnosis. The lull in the symptoms after the first onset of the disease happens more generally in phosphorus-poisoning than in yellow atrophy. Yet the clinical differences between various cases of either affection are greater than those which have been relied upon as separating the two affections. Köhler has asserted that oxymandelic acid in atrophy of the liver replaces the sarco-lactic acid of phosphorus-poisoning, and stress has been laid upon the asserted facts that in the natural disease leucin and tyrosin are present in abundance in the urine, whilst in the poisoning they are absent. In yellow atrophy, however, tyrosin is not unfrequently absent from the urine, and leucin present in very small amount, whilst the contrary has already been stated to occur in some cases of phosphorus-poisoning. In regard to the acids in the urine, very careful chemical analysis would in any case be necessary to determine the presence of either of them, and sufficient evidence is certainly not yet forthcoming to show that either of them is really characteristic. Chemical examination is then absolutely necessary in all medico-legal cases. (For discussions of the diagnosis between yellow atrophy and phosphorus-poisoning, see Köhler, *Sydenham Year Book*, 1870, p. 455; Schultzen and Ries, *Annalen des Berlin. Krankenhauses*, Bd. xv., 1869; and especially Dr. I. Ossikovsky, *Wien. Medicin. Presse*, xiii., 1872, abstracted in *Schmidt's Jahrbücher*, Bd. cliv. p. 15. For cases in which the question was legally raised, investigated, and discussed, see *Schmidt's Jahrbücher*, Bd. cxli. p. 167; *Sydenham Year Book*, 1832, p. 430; *Annales d'Hygiène*, Jan. 1869.) According to M. Poulet (*Gazette Médicale de Paris*, Aug. 1872), phosphorus is always eliminated in the urine as hypophosphoric acid, and consequently the poisoning can always be recognized by adding nitric acid to the urine and heating to calcination. If hypophosphoric acid be present, as dryness is reached the mixture suddenly bursts into a flame like a packet of matches.

It has of late years been demonstrated that phosphorus passes into the blood as phosphorus, and not in the form of phosphoric acid or other compound. In poisoning-cases in men the breath is said sometimes to be distinctly phosphorescent; and in animals Bamberger has found phosphorus in the blood, and Husemann and Marne in the liver, two or three hours after its ingestion; W. Dybkowsky (*Hoppe-Seyler's Medicinisch-Chemische Untersuchungen*, Heft i. p. 54) has detected it in the blood and liver ten hours after its ingestion; and other observers have demonstrated its presence in almost all of the tissues. It seems probable that to some extent it finds entrance into the circulation by being dissolved in the various fatty matters contained in the alimentary canal. At the temperature of the body, however, it yields abundant vapors, and Bamberger has demonstrated that these readily and rapidly pass through animal membranes. He has found that defibrinated blood, when separated from the fumes of phosphorus only by an animal membrane, rapidly becomes saturated with the poison. Dybkowsky (*loc. cit.*) has confirmed this, and it cannot be doubted that in a similar manner living blood absorbs the poison from the alimentary canal.

W. Dybkowsky (*Hoppe-Seyler's Medicinisch-Chemische Untersuchungen*, Heft i.) renders it probable that the theory of Schuchardt (*Henle und Pfeufer's Archiv*, N. F., Bd. viii.) that the phosphorus to some extent in the alimentary canal, but much more largely in the veins, is converted into phosphuretted hydrogen, and that some of this compound and some of the phosphorus itself is oxidized in the venous blood, so that phosphoric acid, besides phosphorus and phosphuretted hydrogen, is emptied into the arterial blood; further, that the last two compounds are oxidized at the expense of the arterial blood and the tissues it feeds, and that the poisoning is due to this deprivation of oxygen. For the details of the experiments upon which these conclusions rest I must refer the reader to the original memoir.*

The indications for treatment in phosphorus-poisoning are very evident. It is plain that no medication can influence the terrible organic lesions induced, and that the primary object must be to prevent the absorption of the poison. Emetics and purgatives are, therefore, of prime importance. As phosphorus is soluble in oils, *no fatty matters* should be allowed either in the food or medicines. As an emetic, *sulphate of copper should always be chosen*.

The minute particles of phosphorus adhere so closely to the alimentary canal that they cannot be dislodged by mechanical means, and an antidote is urgently demanded. For the purpose of oxidizing the poison, Duflos suggested *magnesia usta* and *liquor chlorini*, and Scherer the chloride of lime; but in practice these substances have been found of no value, on account of the slowness of their action.

The oil of turpentine, originally proposed by Andant (*Journal de Médecine de Bruxelles*, 1868-69) as an antidote to phosphorus, has been largely used by experimenters, with apparently contradictory results, which, as is now known, were due to the employment of different varieties of the oil. There are in European commerce three varieties of turpentine,—the rectified, the German, and the French. Jonas (*Liebig und Wöhler's Annalen der Chemie*, Bd. xxxiv.) found that whilst the pure oil has no effect upon phosphorus, with the acid French oil it forms a crystalline, spermaceti-like mass. This is soluble in ether, alcohol, and alkaline solutions, and has received the name of turpentine-phosphoric acid. It is said to be eliminated by the kidneys unchanged, and to exert no deleterious influence. The elaborate experiments of Vetter on dogs and rabbits gave results in accord with these facts, for he found the rectified and German oils to be of no value in phosphorus-poisoning, whilst the crude acid French oil was distinctly antidotal. Unfortunately, the French oil is not kept in American drug-stores, and I have found that neither the ordinary American oil of turpentine nor the Canada Balsam has any action on phosphorus.

* M. Lecorché (*Archives de Physiologie normale et pathologique*, tome i., 1868, tome ii., 1869) believes that phosphorus acts in the blood as phosphoric acid, but does not establish his opinion. For a discussion of this, see Dybkowsky's paper.

As was pointed out by MM. Eulenberg and Guttman (*Aertz. Literaturblatt*, 1868, No. 12, quoted in *Sydenham Year Book*, 1868, p. 450), and subsequently by Prof. Bamberger (*Wiener Medicinische Presse*, Jan. 1872; *Virchow's Archiv*, June, 1872), phosphorus in a solution of a soluble salt of copper becomes immediately black, owing to the formation of a phosphide of the metal. Prof. Bamberger (*loc. cit.*) also asserts that, whilst this change is very rapid, that induced by turpentine is a slow one, and, from an elaborate series of experiments upon animals, concludes that copper is much the more valuable and certain antidote. In human poisoning, then, sulphate of copper should be given in dilute solution, three grains every five minutes until vomiting is induced. After this, if the French oil be accessible, it may be given freely in emulsion. Otherwise, sulphate of copper, with opium to restrain the emesis, should be administered in such doses as the stomach will retain. Sulphate or citrate of magnesia should be used as a quickly-acting purge, and symptoms as they arise should be judiciously treated.

Matchmakers and other artisans who are exposed by their occupations to the fumes of phosphorus suffer from chronic poisoning, which, whilst in many cases it profoundly affects the vitality of the sufferer, is especially distinguished by the occurrence of necrosis of the upper or lower jaw. It has long been known that those artisans who have bad teeth are especially liable to be seriously affected, and the experiments of Wegner have demonstrated that the necrosis of the jaw is due to the local action of the vapor of phosphorus upon the part. He found that when rabbits were kept in an atmosphere full of the fumes of the poison no necrosis ever occurred, unless, by means of an unsound tooth or an artificial wound, the atmosphere had access to the bone. If such access were, on the other hand, allowed to any bone of the body, periostitis and subsequent necrosis resulted. Further, when rabbits received continuously small doses of the phosphorus by the mouth, no necrosis occurred even after wounds which laid bare the bones. As phosphorus-necrosis belongs to the province of the surgeon rather than of the physician, I will not discuss it further here.

ADMINISTRATION.—Phosphorus may be given in pill or in solution, but the liquid form is preferable. The following methods of preparation yield very elegant products, and afford, I think, the best methods of exhibiting phosphorus. Into five fluidrachms of pure almond or olive oil, contained in a glass flask or bottle, drop three grains of transparent phosphorus in small fragments. Place the whole in a water-bath at 175° F., and agitate until dissolved. One minim of this oil represents $\frac{1}{100}$ grain of phosphorus, and may be administered in capsules or in accordance with the following formula: R Olei phosphorati, fʒss; Ol. gaultheriæ, fʒss; Mucil. acaciæ, fʒiiss. M. S.—Dose, one teaspoonful after meals, to be increased to a dessert-spoonful as the stomach will bear it.

Prof. S. R. Percy advises the following formula for solid phosphorus. Take of purified butter of cacao a weighed quantity, melt it over a water-bath, and

for each one hundred grains add one grain of transparent phosphorus in small pieces ; keep it on the water-bath until the phosphorus is all dissolved. When almost cold, add one or two drops of naphtha. It may be poured, while warm, into gelatine capsules or into shallow dishes ; when cold, small pieces may be worked in a porcelain mortar into a pill mass, and quickly rolled out into pills of the desired size. These pills may be coated with collodion, gelatine, or white shellac dissolved in alcohol.

CLASS III.—CARDIAC STIMULANTS.

THE term cardiac stimulants is here used to designate a number of medicines which, when given internally, increase the power and force of the circulation, and are used by the physician for such purposes. There are some substances which are heart-stimulants in reality, but which possess other properties in so great a degree as to overshadow their cardiac relations, and are not used by the physician to affect the circulation. Such medicines are considered in connection with those powers which give to them their clinical value, and are consequently not included in the present class.

Some of the members of this class are slow in their operation, some more rapid. Some produce increase in the pulse-rate, some lower it. It is evident, then, that no general indications can be laid down for their use, but that medicines so diverse in their relations to disease must be studied individually.

AMMONIA. (NH_3 .)

Ammonia is a colorless, irrespirable, highly irritant gas, of a strong alkaline reaction, extremely soluble in water. It is obtained upon a large scale as a waste product in the manufacture of coal gas, and is officinal in the *Materia Medica* list of the U. S. Pharmacopœia in the form of the stronger water of ammonia, and the muriate, carbonate, nitrate, and sulphate of ammonium.

PHYSIOLOGICAL ACTION.—Locally applied, ammonia is a very powerful irritant. When inhaled, it causes intense irritation, and finally inflammation of the mucous membrane of the air-passages, and its solution, if kept in contact with the skin, reddens, blisters, and at last produces even sloughing of the parts. When taken into the system in moderate doses, it acts as a stimulant; but beyond this we have no definite knowledge of what it does. Large doses of it cause in a healthy person a transient feeling of giddiness and exhilaration, and probably in most instances an increase in the force and frequency of the pulse; although in Wibmer's experiments sometimes the arterial system seemed to be unaffected. Indeed, I have been unable to discover that the reputation which ammonia enjoys, and probably justly, as a powerful arterial stimulant, has other than a clinical basis. Its volatility and the great fugaciousness of its effects when administered indicate that it is eliminated in the lungs; and probably this really happens, although the researches of Dr. H. Bence Jones seem to demonstrate that at least a portion of

it is oxidized in the system (*Philosophical Transactions*, London, 1851). The observer mentioned was surprised to find that even large doses, far from increasing the alkalinity of the urine, seem at times to heighten its acidity. It occurred to him that the ammonia might be oxidized; and he accordingly found that the natural product of its oxidation, nitric acid, appears in the urine after the exhibition either of the ammonia itself or of its tartrate, carbonate, or muriate.

When taken in large amounts, ammonia acts as a violent corrosive poison, producing generally abdominal pain, vomiting, bloody purging, and other symptoms of gastro-enteritis, with collapse and finally death. In some cases symptoms of impending suffocation, resulting in death from asphyxia, have occurred, and at the autopsy intense redness and congestion of the bronchial mucous membrane have been present, due no doubt to the irritant having found its way into the bronchi.

The intellect may be clear almost to the very moment of death, or stupor, and finally coma, may be developed.

In the rare instances in which death has taken place within five minutes from the ingestion of the poison, the fatal result has probably been brought about by œdema of the larynx. When the sufferer survives the first few hours, recovery may occur; but death sometimes happens long afterwards from the organic lesions which have been produced.

The treatment of poisoning by ammonia consists in its neutralization as soon as possible by vinegar or other dilute acid, and the meeting of indications as they arise. If the œdema of the glottis be threatening, tracheotomy should at once be performed.

THERAPEUTICS.—Externally, ammonia is much used as a constituent of irritating liniments, and, on account of its efficiency and cheapness, is very valuable. By inverting a watch-glass full of the stronger water of ammonia upon the skin, a blister may be raised in a very few minutes; but, as the effects of the application are very apt to be severe, the use of it is justifiable only under rare circumstances.

Internally the chief indication for the use of ammonia is *failure of the heart's action*. The more sudden and purely functional this is, the more efficacious is the remedy, which should in such cases be not only administered by the stomach, but should also be inhaled through the nostrils, as the local action of the irritant vapor upon the mucous membrane has a very arousing influence. When the failure of the circulation depends upon a slow and persistent cause, as in *adynamic fevers*, ammonia is not generally useful, but may be employed as an adjuvant to alcohol in the crisis of the disorder.

In *poisoning by venomous serpents*, ammonia has been largely used, but certainly is in no sense antidotal, since, according to the experiments of Dr. Fayrer (*Indian Annals of Medical Science*, 1872), mixing it with the poison before injecting the latter into an animal does not in any way delay the fatal result. As an adjuvant to other more powerful stimulants, and especially to

alcohol, ammonia may be useful in these cases. Dr. G. B. Halford, of Melbourne, Australia, has claimed (*Melbourne Argus*, 1872) that when injected into the veins its effects in poisoning from snake-bite are very extraordinary, and several cases of recovery after its use in this manner have been reported. It is far from certain, however, that these cases would have died had no medication been practiced; and Dr. Fayrer states that in an extended series of experiments upon animals he has not found its injection to be of any use. The real value of the remedy must therefore be considered as yet undetermined, but, as the injection can do no harm, it should be practiced (although ammonia certainly is not a specific), yet never to the exclusion of other measures.

In failure of the heart during *anæsthesia*, and in *poisoning* other than from snake-bite, hypodermic injections of ammonia* are worthy of trial, as in some reported cases they have seemed to be of very great service. The same may be said of *sudden collapse* in disease, as sometimes is seen in the *exanthemata*, and not rarely in *pernicious malarial fever*.† Prof. Stillé and other authorities claim for ammonia an antidotal influence in alcoholic intoxication; but that it can relieve absolute drunkenness is, I think, very doubtful.

Ammonia appears to have a tendency to act upon the mucous membrane of the lungs, and may be used as a stimulant expectorant in adynamic pleural inflammations, as in typhoid *pneumonia*.

As a stimulant antacid, ammonia is frequently of service in cases of *head-ache* from gastric acidity.

ADMINISTRATION.—There are four official preparations of uncombined ammonia itself,—namely, AQUA AMMONIÆ FORTIOR (*Stronger Water of Ammonia*), sp. gr. 0.900, AQUA AMMONIÆ (*Water of Ammonia*), sp. gr. 0.960, SPIRITUS AMMONIÆ (*Spirit of Ammonia*), and SPIRITUS AMMONIÆ AROMATICUS, or *Aromatic Spirit of Hartshorn* as it is usually called.

To reduce the strength of the first of these preparations to that of the second requires the addition of one and a half measures of water. The spirit is of varying strength, but is somewhat weaker than the simple water. The aromatic spirit contains both ammonia and its carbonate. For hypodermic use the waters of ammonia are to be preferred; from twenty minims to a drachm of the weaker preparation, diluted with an equal part of water, may be thrown at once into a vein. The spirits, especially the aromatic, are best suited for internal use. The dose of the simple spirit is from twenty-five drops to a teaspoonful, properly diluted.

The *Carbonate of Ammonium* (AMMONII CARBONAS, $\text{NH}_4\text{O}, \text{CO}_2 - 2\text{NH}_4\text{CO}_3$) is the best preparation for continuous use and in typhoid *pneumonia*. It occurs in white, translucent, fibrous masses, which on exposure

* See *Indian Medical Gazette*, June 1, 1872; also, for a case in which the ammonia appears to have done great good, *Chicago Medical Journal*, 1872.

† See Dr. Zuelzer, *Revue de Thérap.-Méd.-Chirur.*, July 1, 1872.

become opaque and efflorescent, parting with ammonia and passing from a sesqui- into a bi-carbonate. It is soluble in four and a half times its weight of water, and may be given in solution in doses of from five to ten grains, repeated *pro re nata*.

The *Nitrate of Ammonium* (AMMONII NITRAS, $\text{NH}_4\text{O.NO}_3$ — NH_5NO_3) is officinal for the preparation of nitrous oxide; the *Sulphate* (AMMONII SULPHAS, $\text{NH}_4\text{O.SO}_3$ — $2\text{NH}_4\text{SO}_4$), for the preparation of *Ammonio-ferric Alum*. The *Muriate of Ammonia* will be considered elsewhere.

ALCOHOL.

Absolute alcohol, *i.e.*, alcohol free from water, is a colorless, volatile liquid, boiling at 172°F. , not congealed by a cold of -166°F. , having the specific gravity of 0.796. It is not officinal, and is never used except for chemical purposes.

The only forms in which pure alcohol is recognized by the Pharmacopœia are ALCOHOL FORTIS, STRONGER ALCOHOL, sp. gr. 0.817, containing 8 per cent. of water; ALCOHOL, sp. gr. 0.835, with 15 per cent. of water; ALCOHOL DILUTUM, DILUTE ALCOHOL, sp. gr. 0.941, with 61 per cent. of water.

Alcohol also exists in the officinal SPIRITUS FRUMENTI, or WHISKY, and SPIRITUS VINI GALlici, or BRANDY, which are obtained respectively by the distillation of fermented grain and of fermented grapes, and should contain from 48 to 56 per cent. of absolute alcohol; and in the officinal wines, VINUM PORTENSE, or PORT, and VINUM XERICUM, or SHERRY.

For medicinal use, brandy should be at least four and whisky at least two years old.

Alcohol is formed out of sugar by fermentation; but as a discussion of the natural history and chemistry of this process, to be of value, would occupy much space, the reader is referred for it to works especially devoted to chemistry and to *materia medica*.

PHYSIOLOGICAL ACTION.—The phenomena induced by the ingestion of alcohol are, unfortunately, so well known as to make any description of them here unnecessary. I have not met with a close experimental study of the order in which the nervous centres are affected, but it is scarcely doubtful that alcohol acts upon them as does ether, excepting that the latter substance, being much more volatile than the alcohol, is consequently absorbed and eliminated much more rapidly, so that its influence is more evanescent. I know by experiment that the vapor of alcohol is capable of producing the stupor known as anaesthesia, and, further, that this anaesthesia may be deepened into death, accompanied by all the phenomena of fatal ether-narcosis.

It is a well-known clinical fact that alcohol given to healthy men increases the frequency, and, to the fingers, the force of the pulse. The very careful and elaborate experiments of Parkes and Wollowicz upon man, with the sphygmograph, indicate that there is produced by brandy an increased rapidity

of the ventricular contraction, with shortening of the period of rest or diastole. The tracings furnished, however, no distinct indications of increased arterial pressure. It is strange that no thorough study of the action of alcohol on the circulation has as yet been made. Dr. H. Zimmerberg, however, asserts that in experiments upon cats with the kymographion he found that alcohol caused a decided reduction of the pulse-rate and of the arterial pressure. If the vagi were cut during this depression, a sudden rise beyond the normal point in both pulse-rate and arterial pressure occurred. When the alcohol was injected towards the heart into the jugular vein, the fall in the arterial pressure was almost instantaneous and very marked. After section of the vagi the alcohol lessened the arterial pressure without affecting the pulse-rate. These observations would seem to prove that alcohol slows the pulse by stimulating the vagi inhibitory centres, and lessens the blood-pressure by weakening the heart.

This result is, however, so directly opposed to daily experience that there must be some fallacy underlying it. I think this will be found to be in the enormous doses of alcohol used by Dr. Zimmerberg.* Alcohol in moderate amounts appears to be a powerful stimulant of the heart and circulation, but in poisonous doses very probably does diminish both the force and frequency of the pulse.

Owing no doubt to the sensations of warmth induced by its local action on the stomach and by the increased activity of the circulation in the extremities, alcohol has been looked upon as a promoter of animal heat. As long ago as 1848, however, Duméril and Demarquay asserted that after the administration of large doses there is a fall of temperature. Of late years much attention has been given to the subject and positive results have been achieved. As almost all experimenters are in accord, it does not seem worth while to occupy space with a discussion of the subject. References are given to the principal original memoirs.†

* The equivalent amount of brandy would be, in a man, three or four pints.

† For those desirous of reading up the literature of the subject, the following references are given:

- N. S. Davis, *Transactions American Medical Association*, 1855, p. 577.
 C. Bouvier, *Pflüger's Archiv für Physiologie*, p. 370, 1869; Obernier, *Ibid.*, p. 499, 1869.
 A. Godfrin, *De l'Alcool, son Action physiologique, ses Applications thérapeutiques*, Paris, 1869.
 C. Bouvier, *Wirkung der Alcohol auf die Körpertemperatur*, Bonn, 1869.
 Manassein, *Centralblatt für die Med. Wissenschaften*, 1869.
 P. Ruge, *Virchow's Archiv*, Bd. xlix. p. 265.
 C. Binz, *Virchow's Archiv*, Bd. li. p. 153; *London Practitioner*, vol. iii., 1869, vol. v., 1870.
 Brown-Séquard, *Journal de la Physiologie*, 1859, p. 467.
 Jacobi, *Deutsches Klinik*, 1857.
 Tscheschichin, *Reichert's Archiv für Anatomie*, 1866.
 Ringer and Rickards, *London Lancet*, 1866, p. 208.
 Richardson, *Medical Times and Gazette*, vol. ii. p. 704, 1869.
 Parkes and Wollowicz, *Transactions of the Royal Society*, 1870.

It is certainly demonstrated that lethal doses of alcohol produce in animals a fall of temperature which often amounts to 5° C., and that even when intoxication or alcoholic narcosis is alone induced the depression of temperature may amount to 3° C. The proportionate dose necessary to produce distinct nervous symptoms is enormous in the lower animals as compared with man; yet, after the ingestion of amounts of alcohol which are not enough to cause intoxication in the animal, the fall of temperature is slight.—rarely, if ever, more than 1° C., and according to Ruge (*loc. cit.*, p. 265) usually from $\frac{1}{2}^{\circ}$ to $\frac{3}{5}^{\circ}$ C.

The experiments of Richardson show that in some cases very minute doses of alcohol increase slightly the temperature ($\frac{1}{2}^{\circ}$ F. in mammals, 1° F. in birds). As regards animal heat, alcohol acts upon man as upon the lower animals. After doses only sufficient to increase the activity of the circulation, probably as a result of this increase, there is sometimes a very trifling exaltation of temperature (Parkes and Wollowicz). After larger doses there is a slight fall of temperature, and when full intoxication is induced this fall may amount to 3° F. (Ringer and Rickards).

Upon animals suffering from pyæmic fever Bouvier and subsequent observers have found that alcohol exerts a decided antipyretic action, very large doses of it lowering the temperature as much as $8\frac{1}{2}^{\circ}$ C., and altogether preventing the occurrence of fever if narcosis be produced before the development of the latter. In fever in man alcohol exerts a similar influence, but in order to make its antipyretic action decidedly manifest, doses so large as to be toxic must be given (Ringer and Rickards). It has been noticed both in man and in the lower animals (Bouvier) that when the individual is accustomed to the free habitual use of alcohol the temperature is scarcely affected even by large doses.

Very recently, Dr. Franz Riegel (*Deutsches Archiv für Klin. Medicin*, 1873) has made a very elaborate original investigation upon men, comprising as many as eighty-six experiments, and has arrived at the following conclusions, which are in exact accord with what was already in type when his memoir came to hand. As they embody the whole subject in a few words, and may be received as absolutely demonstrated, I offer no apology for interpolating them here:

“1. Alcohol, even in moderate doses, in many cases causes a lowering of the temperature of the body. The amount of this diminution averages as a rule

Anstie, *Stimulants and Narcotics*, London.

Mainzer, *Ueber die Wirkung des Alkohols auf die Temperatur des Gesunden*. Inaugural Dissertation, abstracted by Binz in *Virchow's Archiv*, Sept. 1871.

C. Bouvier, *Centralblatt für die Med. Wissenschaften*, Dec. 1871.

I. S. Lombard, *New York Medical Journal*, June, 1865.

Sulzynski, *Ueber die des Alkohols, Chloroforms und Ethers Einfluss auf den thierischen Organismus*. Inaug. Dissertation, Dorpat, 1865.

S. Rabou, *Berlin. Klinische Wochenschrift*, 1871.

Radziejewski, *Centralblatt für die Med. Wissenschaften*, 1871.

only some tenths of one degree. 2. Only exceptionally is there noticed an elevation of the temperature consequent upon the administration of alcohol; not unfrequently, at least after minute doses, there is no noticeable change. 3. The diminution of temperature in convalescents is, as a rule, less than in healthy subjects, or it may be altogether wanting. 4. In those who habitually drink alcoholic stimulants, the depressing influence of alcohol upon the temperature is almost always wanting. 5. The frequent repetition of the doses of alcohol diminishes their lowering effect upon the temperature. 6. The amount of diminution of temperature is directly proportional to the dose of alcohol given. 7. The depression of temperature caused by alcohol is for the most part of but short duration, and the temperature soon returns to its previous grade."

That the antipyretic action of alcohol is not exerted through the nervous system was proved by Binz, who found that the drug acted powerfully upon the fever of animals after cervical section of the spinal cord, and even prevented the post-mortem rise of temperature. It would appear to be a necessary corollary to this that *alcohol in very large doses lowers temperature by directly checking tissue-metamorphosis.*

The effect of alcohol upon the elimination of carbonic acid by the lungs has been investigated by several observers, with different results. According to the researches of Böcker (*Beiträge zur Heilkunde insbesondere zur Krankheitsgenussmittel und Arzneiwirkungs Lehre*, abstracted by Claude Bernard in *Journal de Pharmacie*, tom. xv., 3d series, 1849), of N. S. Davis (*Transactions of the American Medical Association*, 1855), of Hammond (*Physiological Memoirs*, Philadelphia, 1863), and of M. Perrin (*Archives Générales*, 6th series, tome iv.), there is a decided lessening in the amount of carbonic acid gas exhaled. Dr. E. Smith, however, found that small doses of alcohol increased the elimination of the gas, although brandy, whisky, and gin always lessen the production. It is probable that the action of the small doses of spirits taken may vary, as it is readily conceivable that by checking or aiding digestion, by influencing the circulation, or in some other way, they may at times exert an indirect action which shall overcome any direct influence they may possess. Be this as it may, the weight of evidence appears to be at present in favor of their usually diminishing the elimination of carbonic acid, although the matter cannot be considered as entirely settled.

One of the most important facts to be determined about alcohol is its influence upon the excretion of nitrogenous material. Böcker is commonly believed to have experimentally determined that it lessens the excretion of urea. I have seen only an abstract of the original paper. In it this is not positively asserted, but seems to be inferred. Dr. Hammond has performed a very elaborate series of experiments upon himself: first, when just sufficient food was taken to maintain the weight of the body; second, when more than enough for that purpose was ingested; third, when not enough was taken. Under all these circumstances, urea, chlorine, and phosphoric acid were less-

ened in amount by the ingestion of alcohol. Parkes and Wollowicz (*loc. cit.*) affirm that their experiments gave a contrary result. In examining the reports of their experiments I find, however, that on one of the days the man taking the alcohol had a chill followed by fever. If this day be omitted, the average daily excretion of urea during the alcoholic period was 34.35 grammes; during the time when brandy was taken, 34.8 grammes; and during the water period, 35.02 grammes. The ingestion of alcohol seems, therefore, to have reduced the elimination of urea by about ten grains a day. Although the matter cannot be considered absolutely settled, yet the present conclusion fairly is that alcohol lessens the excretion of urea.

From the time of Liebig's celebrated classification of food until the appearance of the memoir of MM. Lallemand, Duroy, and Perrin, ingested alcohol was almost universally believed to be burnt up in the body. These latter observers asserted, however, that alcohol escapes unchanged from the body, not only because they were unable to detect in the blood or tissues any of the results of its oxidation, such as aldehyde or acetic acid, but also because they found it unchanged in the expired air, the sweat, and especially in the urine. The results obtained by the French investigators were, however, seriously questioned by E. Baudot (*L'Union Médicale*, 1863), who demonstrated that the chromic acid test which Duroy and Perrin had relied on for detecting alcohol in the excretions is so delicate as to reveal .165 grain of alcohol in a quart of water; and Baulot further affirmed as the result of twenty experiments that except after immense doses the amount of alcohol eliminated by the kidneys is so small as practically to amount to nothing. In 1866 Dr. Schulinus (*Archiv der Heilkunde*, 1866), by a very elaborate and laborious investigation, confirmed the results of Baudot, showing that alcohol taken into the blood finds its way by exosmose into all organs in similar proportion, and does not escape through the kidneys unless in very trifling amounts. In several experiments he found that one-fourth of the ingested alcohol had disappeared from the body after from two to three and one-fourth hours, and, as but a fractional portion of the lost amount was eliminated, he concluded that it must have been burnt up.

In 1865 a number of experiments yielded to Anstie, of London (*Stimulants and Narcotics* (reprint), Philadelphia, 1868), similar results, and also demonstrated that the elimination from the lungs is exceedingly trifling. Thudichum investigated the matter on a large scale in 1864, and again with the assistance of Dupré in 1866 (*Tenth Report of the Medical Officer of the Privy Council*, London, 1868). In order to avoid the fallacies of the chromic acid test, the alcohol was obtained from the urine by repeated distillations. In the first instance forty-four bottles of wine, containing four thousand grammes of alcohol, were drunk by thirty-three men, out of whose urine, collected during the next six hours, ten grammes, or only 0.25 per cent., of the injected alcohol were recovered. In the experiments of 1866 the process was substantially the same, but, greater care being taken to get absolute

accuracy and to avoid loss during distillation, 0.82 per cent. of the amount administered was found in the urine. Quite recently, Subbotin (*Zeitschrift für Biologie*, vii., 1871; *Schmidt's Jahrbücher*, 1872, Bd. cliv. p. 261) has made an apparently close experimental study of the subject upon six rabbits, and has shown that elimination continues for a longer time than had been generally believed, and that twice as much of the alcohol escapes by the skin and lungs as by the kidneys. In one experiment he found that 16 per cent. of the alcohol escaped unchanged in the first twenty-four hours; elimination after this time, although perceptible, amounted to very little. As he, like Lallemand and his colleagues, experimented with poisonous doses, his results confirm rather than contradict those of Baudot, Schulinus, Anstie, Thudichum, and Dupré; for it is manifestly evident that after such doses elimination would be proportionately greater than after smaller quantities, as there naturally must be a limit to the powers of the system to oxidize alcohol.

The concurrent work of the investigators* whom I have mentioned certainly demonstrates that but a small proportion of ingested alcohol is eliminated. If it does not escape from the body it must be oxidized in the body, and must partake of the nature of a food. It has been objected to this that no one has as yet been able to detect† in the blood any of the ordinary products of its oxidation; the probable reason of this is, however, that the oxidation is carried as it were at one bound to its ultimate end, the production of water and of carbonic acid. A strong corroborative proof that alcohol is largely consumed in the body is even furnished by the experiments of Lallemand, Duroy, and Perrin themselves, for they proved that elimination ceased, or at least could not be detected by the most delicate tests, *before* the alcohol had all escaped from the body.

A very strong confirmation of the theory that asserts the oxidation of alcohol in the body is found in the researches of our countryman Dr. H. Ford, whose experiments have, however, not been repeated, and therefore lack the absolute authority of complete confirmation. If they shall prove accurate, their importance and originality render them worthy of high praise. Dr. Ford (*New York Medical Journal*, Jan. 1872) has worked upon the supposition that the hepatic sugar must be converted into alcohol in the body before its final destruction, and, using large quantities of blood of animals, has sought by repeated distillations to obtain alcohol from it. Space is wanting to describe in detail the very elaborate methods employed by Dr. Ford. The tests which he relied on to prove that the liquid obtained was alcohol were the chromic acid test, the peculiar inflammability, and the optical appearance of the alcohol in the conducting tubes at the time the distillate commenced to

* I believe these results have also been confirmed by Wöhler, *Journal des Progrès*, xi., but I have not seen the original paper or any abstract of it.

† Duchek (*Vierteljahrsschrift für die Prakt. Heilkunde*, Bd. iii., 1853) thought that he had demonstrated the presence of aldehyde in the blood of animals poisoned with alcohol; but his experiments were not really carried far enough to prove it.

boil. In order to prevent any possible oxidation of the alcohol during the process, Dr. Ford sometimes added sulphuretted hydrogen.

The results of ten experiments are shown in the following table :

No.	Weight of Blood.	Interval from Death to 212°.	Temperature when distilled.	Weight of First Distillation.	Weight of Final Distillate.	Weight of Alcohol obtained.	Weight of Alcohol for 10,000 parts of Blood.	With or without HS.
15	6970	60 m.				0.0350	0.0932	without.
16	9734	56	101°	1602	0.8416	0.0198	0.0203	without.
17	9137	70	100.3	1636	1.6218	0.0605	0.0662	with.
18	9236	77	99.5	1623	3.6130	0.0444	0.0480	with.
19	8988	60	99	1555	2.6092	0.1337	0.1509	with.
20	8854	60	98	1555	1.7320	0.0760	0.0858	with.
21	9423	45	96	1560	1.8722	0.0708	0.0751	without.
22	9112	61	98	1550	0.9552	0.0350	0.0384	without.
23	27330	48	98	14050	10.6883	0.2928	0.1071	with.
24	36300	51	99	17600	14.0606	0.5652	0.1556	with.

Pushing his researches still further, Dr. Ford used various tissues as the substances to be distilled. He also made elaborate calculations, based on the carbon ingested and on the carbon exhaled, as to the amount of alcohol which ought to be found in the capillary blood of the lungs. The results are expressed in the following table :

Alcohol in the capillary blood of the lungs :

{ calculation based on "carbon ingested"		0.5403
{ " " " " "carbon exhaled"		0.5794
"	putrescent lung-tissue (mean of exp. 8, 9, and 11)	0.3819
"	fresh " (" " 12, 13, and 14)	0.3076
"	putrescent thoracic blood (mean of exp. 1, 2, 3, 4, and 5)	0.7625
"	fresh " " (mean of table)	0.0841
"	putrescent liver-tissue (exp. 6)	4.3138
"	fresh " (mean of exp. 25, 26, and 27)	0.0190

The important facts seemingly established in the above table are : the correspondence between the amount of alcohol in the thoracic blood as obtained by calculation and by experiment ; that the smallest quantity of alcohol is to be obtained from fresh liver-tissue, the greatest from putrescent liver-tissue, in which the glycogen must have undergone fermentation.

The fresh thoracic blood was blood which had not traversed the lungs ; the putrescent thoracic blood of course represented the same blood with all its sugar fermented.

These researches of Dr. Ford are certainly corroborated by the discovery of Dupré (*The Doctor*, Feb. 1, 1873) that a substance exactly resembling alcohol is found in very minute quantity in the urine even of teetotalers,*—a dis-

* It is asserted that the substance discovered by Dupré " is not alcohol, although it presents the characters by which we generally recognize alcohol. This body passes over among the earliest products of distillation, gives acetic acid on being oxidized, reduces the

covery made nearly simultaneously by M. Béchamp (*London Lancet*, 1873), who obtained, from the urine of persons who had not taken any alcoholic beverage for a long time, alcohol in sufficient quantity to burn it.

From what has been said, it is certainly deducible that alcohol in small amount is an *arterial* and *cerebral stimulant*, increasing functional activity in the nervous and circulatory apparatus; is a *food*, in the sense that it is destroyed in the system and there performs a physiological office; is, when in sufficient quantity, a retarder of tissue-changes, because it lowers animal heat independently of any action on the nervous system, and also probably checks the excretion of carbonic acid and of nitrogen. Dr. Hammond's experiments indicate that it is a real food, for he gained weight when taking it upon a diet which he had previously proven insufficient by itself to maintain his bodily weight.

If the alcohol be oxidized in the body, and be a food, as it seems to me is clearly proven, it must of course generate force, measurable by the modern standard of the heat-unit. A little calculation will show the importance, or rather the great amount, of the generated force. According to Dupré (*London Practitioner*, vol. ix., 1872, p. 33), one gramme of alcohol oxidized in the body evolves 7184 units of heat, whilst the same weight of lean beef gives off only 1482 units of heat. It has been estimated that 9.3 ounces of lean beef, equal to about two ounces of alcohol, will supply the necessary force to maintain the circulation and respiration of an average man for one day. That is, four ounces of strong spirit will suffice for this purpose. Since to the ability of furnishing material whose consumption shall give power is added the ability to restrict waste and to stimulate the functions of circulation and of the nervous system, it is evident that in alcohol we have a most important means of sustaining the system during the strain of an acute exhausting disease.

THERAPEUTICS.—Our knowledge of the physiological properties of alcohol shows that its chief therapeutic value in acute disease is as a stimulant, a temporary imparter of power which shall enable the system to stand some strain of like duration—to bridge over some period of weakness.

The cases to which it is especially adapted may be divided into three classes.

First. Those in which there is a temporary loss of heart-power, as in fainting from exhaustion, loss of blood, or other cause. In these cases the alcoholic stimulant should, if possible, be given hot, and not much diluted; with it should also be exhibited some more rapidly-acting diffusible stimulant, such as ammonia.

Second. Those acute diseases in which the powers of the system are in danger of being used up; to aid in the digestion of food and the maintenance of power. Alcohol, as has already been stated, is to a certain extent a food, but it will

bichromate of potash when dilute sulphuric acid is present, and its aqueous solution has a lower density than water. It furnishes iodoform, and exists in the urine in a very small quantity." Since it appears to possess all the physical and chemical characters of alcohol, to ordinary minds it is alcohol.

not of itself sustain life for a long time, and should in adynamic disease always, unless for special reasons, be combined with milk, or occasionally with eggs. One great source of its value in these diseases is the power which it imparts of assimilating food, and in milk-punch are furnished the stimulant to digestion and the most perfect food known for digestion. This use of alcohol is apart from its office in the lowest stage of fever as a heart- and nerve-stimulant. Employed for this purpose it is useful in *all* stages of the *adynamic fevers*, such as *typhus* and *typhoid*. By the exhibition of three or four ounces of milk every two hours, with one or two drachms of brandy or whisky, from the beginning of the attack, in many cases the development of the severe adynamic symptoms may be prevented.

In the advanced stages of diseases, when the *typhoid state* is well developed, then alcohol should be given boldly,—to quiet by stimulation the nervous and circulatory systems,—to afford a food which shall in a measure replace the natural pabulum,—to aid in the digestion of milk and other simple nourishment,—to aid in lowering temperature by checking the tissue-waste of fever,—in a word, to enable the system to stand the drain upon its vital powers, and at the same time to check such drain.

Properly administered, it always promotes, not arrests, secretion in these cases. The guide to the amount given should be the effects produced: so long as it lowers temperature and pulse-rate, moistens the dry tongue and skin, and quiets the nervous disturbance, it does good; if, however, the tongue grows drier, the pulse puts on an angry, bounding character, and the patient becomes restless and uneasy, stimulation is being pushed too far, and the amount exhibited should be lessened. The antipyretic action of alcohol has suggested its use in all cases of high temperature; as, however, this is only one of its actions, and as it is not decided unless very large doses be given, alcohol cannot be employed as a general febrifuge. True arterial excitement and sthenic inflammation certainly contra-indicate its use. The rule may be laid down as follows: high temperature is an indication for the use of alcohol only when other symptoms also demand it; in itself high temperature is never a contra-indication to alcohol. In *acute sthenic diseases*, after the progressive stage has passed and the results of the disease simply remain to be overcome, alcohol and milk will often save life. Thus, in *acute pneumonia*, when so much consolidation has occurred as to render it doubtful whether the exuded matter can be removed, or in *abscess*, when large amounts of pus have formed, the demand may be very great for alcohol as a food and as an aider of digestion, and sometimes as a stimulant.

Third. Those in which there is a depressing agent. In many forms of *poisoning*, alcohol may be used with signal advantage simply as an arterial and nervous stimulant, to overcome the influence of a depressing agent. Thus, in *snake-bite* the free use of it affords, with the hypodermic use of ammonia (see Ammonia), the best method of treatment. In *poisoning by aconite, veratrum viride*, or other similar substance, where death is threatened through

failure of the heart-power, alcohol in some form is imperatively needed. In these cases it should be given freely, not much diluted, and, if convenient, hot. Not too large amounts should be given at a draught, but the dose should be repeated every five, ten, or fifteen minutes, until the desired effect has been obtained or death has resulted.

What has been said up to this point in regard to the therapeutic action of alcohol has had reference to acute disease. The value of the drug in some chronic diseases cannot be doubted; but in prescribing it the physician should never lose sight of the possible danger of producing a habit far worse in its fruits than death itself.

In *chronic neuralgia*, in *hypochondriasis*, in *melancholia*, temporary relief may sometimes be obtained by the use of stimulants; but the very relief afforded doubles the temptation to the frequent use of the alcohol, and, as the system becomes habituated to its action and the dose has to be more and more increased, the habit of frequent stimulation grows almost of necessity into drunkenness. For this reason I do not think the physician is ever justified in prescribing alcohol for its narcotic stimulant effect in these cases. The chief legitimate uses of alcohol in chronic diseases are to aid in digestion; to furnish a food which, without any digestive effort upon the part of the system, shall be absorbed and shall take the place of more ordinary food; and to check excessive tissue-waste. Of course these indications exist only in such diseases as are either dependent upon or closely associated with a condition of system in which the general nutrition is depraved. In purely local affections the use of alcohol is rarely called for except in the last moments of life, when it may always be employed to afford relief and to protract for a short time the struggle.

In *chronic dyspepsia*, alcohol administered with the food often aids very materially in the assimilation of the latter; but care has to be exercised in prescribing it, for the same reasons as were given when speaking of the use of stimulants in melancholia a moment since. In many cases of *chronic neuralgia*, not as a narcotic stimulant, but as a food and a stimulant to nutrition, alcohol is often of the greatest service. The danger of establishing a fatal habit in this disease is, however, excessive. In almost all cases in which alcohol is called for in neuralgia, cod-liver oil is also indicated, and it is generally best to exhibit the two remedies together, so as to obtain the easy assimilation of the oil and to guard against any evil moral results.

In *phthisis* and its congener *scrofulosis*, there can be no doubt as to the great value of alcohol; and in the latter stages of consumption its judicious use as an antipyretic narcotic stimulant to lessen the sufferings of the patient is perfectly justifiable. During the chronic movements of the affection, alcohol taken with cod-liver oil, or in small amounts with the food at meal-times, conduces not so much to the comfort as to the well-being and recovery of the patient.

The question as to the propriety of the daily use of alcohol by healthy

men is at present a very serious one, involving so many moral and politico-moral issues that it cannot be fully discussed here. Suffice it to state, as obvious inferences from our present knowledge of the physiological action of alcohol, that the habitual use of moderate amounts of alcohol does no harm; that to a certain extent it is capable of replacing ordinary food, so that if the latter be scanty, or even if it be coarse and not easily digested, alcohol, in some form or other, is of great advantage; that in all cases it should be taken well diluted, so as not to irritate the stomach, and that wine or malt liquors are certainly preferable to spirits. The experience of arctic explorers has certainly shown that alcohol has no heat-producing power, so that at a time when it was believed to have such influence by physiologists the Northern navigators had learned that the free use of spirits, far from enabling a man to withstand habitual exposure to intense cold, very materially lessened his power of resistance. On the other hand, the experience of almost every trout-fisherman or sportsman has satisfied him that spirits do have power to prevent "catching cold" under sudden and unaccustomed exposure to wet and cold, and that benumbed extremities will become warm and have their proper feelings return under the influence of a glass of whisky. There is, however, nothing strange or contradictory in these experiences, and they are both in strict accord with the present knowledge of the physiological action of the drug. As is often the case, the facts were practically made out, however, before science could solve the apparent paradox. As has been abundantly shown, alcohol has no heating power; but the chill of sudden exposure, the suffering benumbed extremities, the bronchitis that perhaps follows, all mean simply this: that, as the result of the cold, the blood leaves the surface and the extremities, the circulation fails in the outposts, and, as a consequence, suppressed perspiration—*i.e.*, suspended function of the skin—and internal congestions result. The relief afforded by the spirits, as well as the prevention of sickness, is due simply to the power of the remedy in maintaining the circulation and keeping the external surfaces well supplied with fresh blood and well warmed by the constantly-renewed currents from the interior furnaces of the body.

ADMINISTRATION.—Almost enough has been already said upon this point, but a few further remarks seem appropriate. When stimulants are used to sustain the sinking powers in poisoning or in disease, the amount given should be almost solely regulated by the effects. Thus, in snake-bite it may be necessary to give a pint of whisky in the course of an hour; and in low fevers I have seen the greatest benefit result, and life apparently saved, by the exhibition of a quart of spirits a day. The rule is always to be governed by the effects. In poisoning, one, two, three, or four ounces, as the case may seem to need, should be exhibited every ten minutes, until some effect is produced or matters become hopeless. In low fevers half an ounce to an ounce should be given every one, two, or three hours, *pro re nata*, the practitioner watching the results, as already spoken of.

The question of choice, of course, comes up in every case so far as the

spirits are concerned. I have never been able to perceive any difference in their action (gin, of course, being excepted), save only that sometimes one agrees better with the stomach than the other. This has seemed to me to depend simply upon the personal likings of the patient, to which therefore the choice may well be left.

In sudden collapse, some of the wines with a very high *bouquet* are believed to be more stimulating, on account of the ethers which they contain; but I have had no experience with them. In convalescence and for habitual use in health, wines are preferable to spirits,—more agreeable, more tonic, and less apt to lead to excessive indulgence.

When a mild stimulant is wanted in the beginning of fevers, especially if milk-punch seems "too heavy," wine whey may be sometimes used with advantage. It is made by pouring a half-pint of sherry or madeira into a pint of boiling milk, stirring thoroughly, and after coagulation has occurred straining off the whey, which may or may not be sweetened, according to the taste of the patient. *Champagne* for use in medicine should be very dry, and is especially useful in patients with delicate stomachs, especially if nausea or vomiting actually exists.

TOXICOLOGY.—Alcohol not rarely produces death as an acute poison.

The acute form of alcoholic poisoning in its minor degrees is, unfortunately, an hourly occurrence almost in every village, but that fatal results are not absolutely so rare as is generally believed is shown by the fact mentioned by Taylor, that in four years (1863–67) thirty-five deaths from this source have occurred in England and Wales. The absolute diagnosis of acute alcoholic poisoning when the patient is simply seen in the advanced stage of deep coma cannot be made out. The odor of liquor upon the breath or about the person is simply a proof that the subject has been drinking, not that the symptoms are caused by alcohol. The manifestations are merely those of profound compression or congestion of the brain, of apoplexy, of opium-poisoning; and a man who has been drinking only moderately may have been struck down with apoplexy or poisoned with opium. Dr. Hughlings Jackson has recorded a case in which the alcohol impregnated not only the breath but the urine also, and in which the patient was left to sleep it off; but at the post-mortem a clot was found covering nearly the whole of one hemisphere. After death in acute alcoholic poisoning the stomach is usually found very much congested, and sometimes ecchymosed. The treatment consists in the evacuation of the stomach, the use of the alternate hot and cold douche, and the usual mechanical methods of arousing a narcotized patient.

TEREBINTHINA—TURPENTINE.

White turpentine is the concrete oleoresin obtained by incising the *Pinus palustris* and other species of pine. The supply in the American market comes almost exclusively from North Carolina and other of our Southern States. It is rarely, if ever, itself used in medicine, but by distillation is

separated into a volatile oil and a resin (*Rosin*), which is officinal under the name of *Resina*. Rosin is used in medicine solely as an addition to certain plasters, chief among which is the *Emplastrum Resinæ*, *Adhesive Plaster*, or, in ordinary language, "*Sticking Plaster*," so constantly used by surgeons.

OLEUM TEREBINTHINÆ—OIL OF TURPENTINE.

This is a yellowish, highly-inflammable oil, of a strong peculiar odor and a hot biting taste, moderately soluble in alcohol, freely so in ether, very slightly so in water. By heating with muriatic acid it is converted into a red liquid and a white crystalline substance, which, from its resemblance to camphor, has received the name of *artificial camphor*. Turpentine is remarkable for having the property of absorbing oxygen and converting it into ozone.

PHYSIOLOGICAL ACTION.—Turpentine is a powerful irritant, causing in a very short time inflammation in any tissue with which it may be brought in contact in its undiluted form.

When taken by a healthy person in moderate doses, it produces a sense of warmth in the stomach, soon followed by exhilaration, and, if the amount be sufficient, giddiness and even a species of intoxication. The pulse is increased in force and frequency. The turpentine escapes from the body through the lungs and kidneys, imparting its own odor to the breath, and that of violets to the urine. Although several recorded instances prove that turpentine is capable of producing death, yet cases of serious poisoning by it are rare, and a lethal result exceedingly so. The symptoms noted in poisoning by it are most of them constant, but vomiting and purging are present in some cases and not in others. Unconsciousness is generally complete, and in some cases is accompanied by dilated pupils; the urine is very much lessened in quantity, often bloody, not rarely suppressed; the skin is sometimes dry, sometimes moist; the pulse is feeble, rapid, and generally regular. The intense irritation of the urino-genital organs in some cases has been indicated by constant priapism and efforts at micturition.

The lethal dose must be very large, but is not definitely known, since recovery from four ounces in an infant fourteen months old has been reported. In Dr. Maund's case (*Annuaire de Thérapeutique*, 1846), death was supposed to have been produced in an intemperate woman by six ounces; and Philip Miall has recorded an instance of death produced in an infant fourteen weeks old by turpentine, of which half an ounce was thought to have been taken (*London Lancet*, March, 1869).

The symptoms produced by turpentine show that it is a stimulant to the nervous and arterial systems; but we have no definite knowledge of the exact methods in which it acts. The only attempt at a close physiological investigation of the subject which has come under my notice is that of Dr. Hoppe (*Journal für Pharmacodynamik*, Bd. i. p. 105), who concludes, on what seems to me insufficient evidence, that the vaso-motor nerves are the first to feel its influence.

Upon the kidneys the action of turpentine is very marked, causing, according to the size of the dose, increased secretion, or symptoms of renal irritation, such as pain in the back, and strangury, with a diminished excretion of bloody urine.

THERAPEUTICS.—Externally the oil of turpentine is very much employed as a powerful counter-irritant. It is useful more especially when it is desired to act upon a large extent of surface. When a very intense permanent local impression is required, a blister is to be preferred. Thus, in *pleurisy* a blister may be used, in *bronchitis*, turpentine stupes. In preparing the latter the turpentine should first be warmed by setting the vessel containing it in hot water, then a piece of flannel which had just previously been saturated with hot water and wrung as dry as possible should be dipped in the turpentine, and again wrung out. It is then ready for application, and may be left on from fifteen minutes to half an hour, according to the sensitiveness of the patient's skin.

Another local use of the oil of turpentine is as an addition to enemata. From a teaspoonful to an ounce of it mixed with double its amount of olive oil renders opening enemata much more active, especially in causing the expulsion of flatus. Turpentine enemata containing much of the oil in a small bulk are also constantly used with good effect in arousing the system from stupor arising from narcotic poison or similar causes.

In *ulceration of the bowels*, turpentine taken by the stomach is often very efficient, probably acting locally in the intestine, and in *simple gastric ulcer* the very best results are sometimes derived from its use. In very large doses (fʒss to fʒi with an equal amount of castor oil) it is a very efficient vermifuge, especially against the round worm.

Internally, turpentine may be used as a stimulant in low fevers, especially when the tongue is dry and red.

In *typhoid* or *enteric fever* it without doubt acts as a local stimulant to the ulcerated bowel, besides influencing the general condition of the system. There are two conditions or stages in the disease named in which it is especially useful,—indeed, is of incalculable service. About the end of the second week the tongue sometimes becomes very dry, red, chapped, perhaps coated in the centre with a brownish fur, and at the same time marked meteorism develops. Ten drops of turpentine every two hours during the day and every three hours during the night will in the majority of cases remove the bad symptoms noted. That the action of the oil is largely a local one is shown not only by the arguments of the introducer of the practice, Dr. George B. Wood, but also by the value of the same treatment when diarrhœa persists after the acute stage of the fever has passed. When convalescence is protracted, when there is a constant tendency to the recurrence of diarrhœa, when, in other words, the ulcers of Peyer's patches are slow to heal, turpentine acts almost as a specific.

In *typhoid bronchitis* and *pneumonia*, especially as intercurrent in typhus

fever and similar diseases, turpentine applied externally and taken internally is often very useful.

The same may be said of the low forms of *puerperal fever*. In this disease the abdomen should be kept covered with fomentations of the oil and of warm water alternately, the counter-irritant being used as constantly as a proper regard for the skin of the patient will allow. Internally it should be given in very large doses (℥ss) every two hours.

In *hemorrhages* from the stomach, bowels, or lungs, turpentine has acquired celebrity, but is hardly so much used as formerly. It is in the ataxic cases that it is useful. I have very rarely employed it, as the *oil of erigeron* has seemed even more efficacious, and is much more pleasant to the patient. In *purpura hemorrhagica* turpentine has been highly praised.

DIGITALIS.

The leaves of the *Digitalis purpurea*, or fox-glove, of the second year's growth. These are large leaves of a dull pale green, with whitish down underneath, and have a bitter nauseous taste and a faint narcotic odor. *Digitalis* yields both to water and alcohol. M. Homolle discovered in it a peculiar bitter principle, which has been abundantly proven to contain the medical virtues of the crude drug, and, under the name *Digitaline* (*DIGITALINUM*), is now officinal in the U. S. Pharmacopœia. As prepared according to the directions of the latter, it is a whitish or yellowish powder, odorless, but of a very bitter taste, nearly insoluble in ether and in water, readily soluble in alcohol and in acids. With muriatic acid it makes a yellow solution, which soon changes to green.

The officinal preparation is made by a process copied after that of M. Henry and M. Homolle. It is probably impure, and crystallizes with great difficulty, and imperfectly. Nativelle, in 1871, announced (*Bulletin de l'Acad. Roy. de Médecine*, 1871, vol. xxxvi.) to the Academy of France that he had discovered a process by which crystallized digitaline could be prepared, and, on the report of a commission, received the grand prize of Orfila (six thousand francs).

Crystallizable digitaline (*Journal de Chimie Médicale*, 1873) occurs "in short and delicate needle-shaped crystals, and possesses an intense and persistent bitter taste. It is but slightly soluble in water, soluble in twelve parts of cold and six of boiling alcohol of 90°, less soluble in absolute alcohol, and nearly insoluble in ether; very soluble in chloroform. It is rapidly dissolved by a solution of chloral hydrate, the solution becoming greenish blue in color. The concentrated mineral acids dissolve it, hydrochloric acid producing an emerald-green color, sulphuric acid a green which if subjected to the action of bromine fumes changes to a dark red, nitric acid a yellow, nitro-muriatic acid a yellow changing to an obscure green, and a mixture of equal parts of sulphuric and nitric acids a rose color changing to a deep violet. When heated on platinum, it melts, swells up, becomes brown, and disappears without

leaving any traces. It contains no nitrogen, but is composed of 51.33 per cent. of carbon, 6.85 per cent. of hydrogen, and 41.82 per cent. of oxygen."

According to Ch. Blaquart (*L'Union Pharmaceutique*, Nov. 1872), ten per cent. of crystallizable digitaline can be extracted from the crude drug, which probably contains twelve per cent. of it. The question whether this substance is the pure active principle of digitalis is of course an important one. The French commission reported as the result of physiological experiments that it produces in man and animals effects similar to those of the amorphous digitaline, but that it is much stronger than the latter. This conclusion has, however, met with some opposition. M. Gubler (*Bulletin de l'Acad. Roy. de Médecine*, vol. xxxvii. p. 404) denies that this crystallized digitaline is stronger than the amorphous preparation, and M. Vulpian in the subsequent discussion asserted that in experimenting he had found them of equal strength. Dr. Roucher (*Les Mondes*, July 18, 1872) even asserts that the crystallized digitaline is a complex body; but this is probably a mistake. Recently Ch. Blaquart (*loc. cit.*) has arrived at the experimental conclusion that it is almost exactly the physiological equivalent of the amorphous principle. It seems probable that some specimens of amorphous digitaline are equal, or very nearly so, in strength to the crystalline variety; but nothing has, I think, been brought forward to disprove that the latter is the active principle of the drug, and that it is more constantly pure than the other preparation. About one-ninth of a grain of it is said to have been given to an adult without causing a toxic effect.*

PHYSIOLOGICAL ACTION.—To the therapist the interest in the physiological action of digitalis centres chiefly upon the circulation. The drug does, however, exert a direct influence upon the apparatus of voluntary motion which is worthy of notice. In toxic doses it lowers reflex activity, and induces lassitude, prostration, muscular tremblings, and sometimes convulsions. That the muscles themselves are affected has been proven by the researches of Vulpian, of Dybrowsky and Pelikan, and of Gourvat, all of whom have found that the muscles of frogs poisoned with digitalis respond more feebly than is normal to galvanic currents. The nervous tissue has, however, been found by Gourvat to be more susceptible than the muscular, the nerves losing their functional power sooner and more completely than the muscles.

As the result of an elaborate experimental study (*Reichert's Archiv für Anatomie*, 1871, p. 252), Dr. A. Weil concludes that digitalis first lessens reflex activity by directly—*i.e.*, independently of its action on the circulation—exciting the inhibitory reflex centres of Setschenow, and after a time by directly paralyzing the spinal cord. The experiments upon which this conclusion was based are divided into two series, in the first of which it was found that after small toxic doses of the poison great diminution in the reflex activity of the frog was apparent in from ten to twenty minutes, and continued until

* For the process of manufacture, see *Boston Medical and Surgical Journal*, p. 35, 1873.

the death of the batrachian; but that this diminution for from twenty-five minutes to an hour was immediately suspended by section of the cord high up, the reflex activity returning at once to its normal state; that after large doses the reflex movements were almost abolished in five minutes, and continued until death, but at any time during the first ten or twenty minutes could at once be restored by section of the upper cord; that, both after large and small toxic doses, a time finally came when division of the cord had no power to restore the lost reflex functions. The second series of experiments were directed to discovering whether the action upon the inhibitory reflex centres and the cord was a direct one, or was simply the result of the altered circulation. In this part of the investigation, the hearts of frogs were cut out, or their motion arrested by the local application of a concentrated solution of the nitrate of potash, or rendered slower by a dilute solution of the same salt, and the effects of these various procedures upon the reflex activity were studied. It was found that slowing of the heart's action did excite the Setschenow's centre, but not to nearly so great an extent as did digitalis, and that minute doses of digitalis sometimes stimulated the Setschenow's ganglion and lowered reflex activity before the heart was sensibly affected. In regard to the spinal cord it was proven that when the heart was killed by the local action of potash the reflex functions of the spinal ganglia remained intact for a much longer period than when digitalis was administered.

The action of digitalis upon the heart of the frog was, I believe, first investigated by M. Vulpian (*Comptes-Rendus de la Société de Biologie*, 1855, p. 70), who has been followed by numerous observers, among whom may be mentioned W. Dybkowski and E. Pelikan (*Zeitschrift für Wissenschaftliche Zoologie*, Bd. xi., 1862), A. B. Meyer (*Arbeiten aus dem Physiologischen Institut zu Zürich*, quoted by Boehm), Legros and Legroux (quoted by Gourvat), Claude Bernard (quoted by Gourvat), Rudolf Boehm (*Pflüger's Archiv für Physiologie*, Bd. v., 1872), Homolle (*Archives Générales de Médecine*, July, 1861), Gourvat (*Gazette Médicale de Paris*, 1871), Fothergill (*Digitalis*, London, 1871), Fagge and Stevenson (*Proceedings of the Royal Society*, London, vol. xiv.).

The statements of these investigators agree in all essential points. One or two of them have occasionally noted a primary brief acceleration of the heart's action; but the rarity of its occurrence shows that it has been probably produced by some extraneous unnoted influence.

The first distinctive action of the drug is a marked lessening of the number of cardiac beats per minute, due to a prolongation of the diastole, which may be complete, but is more generally divided by an abortive attempt at ventricular contraction. The systole is abnormally energetic, so that the ventricles become white as the last drop of blood is squeezed out of them. As the action of the drug becomes more intense, the rhythm of the heart is very much affected, the ventricle and auricle no longer beating in accord. At the same time the diastole generally becomes imperfect, one portion of the ven-

tricle maintaining its systolic spasm, while the rest dilates. Thus, the extreme apex may remain hard and white during the diastole, and even hernial protrusions of the ventricle may occur. Finally, the heart is arrested in systole; and as the muscle so hardens, of course all its power of responding to electrical or other excitants is lost.

In some rare instances, instead of the above series of phenomena, the diastolic periods throughout are prolonged and quiet, and after several periods of relaxation, lasting for ten or twenty seconds, final diastolic arrest may occur.

As both Boehm (*op. cit.*, p. 163) and Dybkowsky and Pelikan (*loc. cit.*) have found that the slowing of the heart's beat, the increased energy of contraction, the irregularity and final systolic arrest, are produced by digitalis after division of the vagi and destruction of the spinal cord, and as both Ackermann (quoted by Boehm, *op. cit.*, p. 158) and Boehm have found that the paralyzing of the peripheral ends of the vagi by atropia does not prevent the phenomena just alluded to, it is evident that the drug acts directly upon the heart-muscle itself, a conclusion which is confirmed by Eulenburg and Ehrenhaus (quoted by Dr. T. Lauder Brunton, *On Digitalis*, London, 1868, p. 51), who found that digitalis, when locally applied, acts at once upon the heart. On the other hand, the inhibitory activity of the peripheral ends of the pneumogastrics is without doubt increased by the drug. There is no stage in which stimulation of the vagi does not cause diastolic arrest. Indeed, Dybkowsky and Pelikan have seen galvanization of the nerves produce such relaxation in the auricles after the ventricles had already become permanently contracted. Further, Boehm has found that a stimulation of the pneumogastrics which is insufficient to make itself felt before poisoning will, after the exhibition of digitalis, cause diastolic arrest lasting for many minutes.

It appears, therefore, that the peripheral cardiac inhibitory apparatus shares in the stimulant action of digitalis; and, as Boehm has found that diastolic arrest never takes place in frogs poisoned with the drug, after section of the vagi, it is probable that this rare mode of death is really due to super-excitation of the inhibitory cardiac nerves.

Rudolf Boehm (*op. cit.*, p. 170) has investigated the influence of digitalis upon the working power of the heart when freed from all connection with the central nervous system. By using the method of Ludwig and Coats, he obtained as a constant result that the amount of work done was increased by small doses of digitalis; that after large doses a similar increase was followed in a short time by very great diminution in the expenditure of power by the heart, a diminution apparently due to imperfect diastole and consequent non-admission of serum into the viscus.

The elaborate experiments of L. Traube (*Gesammelte Beiträge zur Pathologie und Physiologie*, Bd. i., Berlin, 1871) upon warm-blooded animals showed that in dogs moderate doses of digitalis produce increased arterial pressure, with lowering of the rate of the cardiac pulsation. When toxic doses were used, these phenomena were followed by increase of the pulse-

frequency and fall of the arterial pressure, which, however, did not commence at the same time, since the maximum pressure was not reached until the pulse had risen above the original, normal point.

Boehm (*loc. cit.*) has confirmed these results, and has also noticed a very marked dicrotic pulse, evidently due to an abortive ventricular contraction during diastole, precisely as occurs in frogs.

The experiments of Brunton (*loc. cit.*) and of Gourvat (*loc. cit.*) also are in accord with those of Traube; so that it may be considered proven that in mammals moderate doses of digitalis produce rise of arterial pressure with diminished pulse-rate.

Prof. L. Traube has found that, after section of the vagi, digitalis is in warm-blooded animals, with rare exceptions, incapable of reducing the pulse-rate, and, contrariwise, that when the pulse-frequency has been reduced by the drug, section of the nerves causes an immediate and very marked rise in the rate of pulsation. I believe Boehm* has experimentally confirmed this; and in the single experiment of Gourvat a similar result was attained.

The conclusion would seem to be inevitable, that in mammals the reduction in the pulse produced by digitalis is directly or indirectly owing to an excitation of the peripheral inhibitory apparatus. The occasional reduction of the pulse-rate after section of the pneumogastric shows, however, either that the inhibitory nerves in some animals find another path than the pneumogastriacs, or else that there is an additional—sometimes inoperative, sometimes efficient—cause of the reduction of the pulse-rate. As it has been shown that digitalis is capable of slowing the beat of the isolated heart of the frog, it would appear probable that it may exert a similar influence at times, in mammals, upon the cardiac muscle or its contained ganglia.

Although digitalis does increase the muscular energy of the heart, it seems scarcely possible that the enormous rise of pressure produced by it can be owing to this alone. This *a priori* reasoning has received experimental confirmation from Malan,† Fothergill (*loc. cit.*), Gourvat (*loc. cit.*), Ackermann (*Ueber die Wirkungen der Digitalis*, in *Volkmann's Sammlung Klinischer Vorträge*, No. 48, Leipsic, 1872), and Boldt (*Inaugural Dissertation*, *Schmidt's Jahrbücher*, March, 1872). The first three of these investigators have found that the arterioles of the frog's web as seen under the microscope undergo very decided contraction after the systemic use of digitalis; and Ackermann states that if the abdomen of a rabbit be opened so as to expose the arteries of the mesentery, a very marked contraction, even to the partial obliteration of the lumen of the vessels, can be readily seen to follow the exhibition of digitalis.

Boldt experimented upon curarized frogs after the manner of Cohnheim, and found that the first effect of the digitalis was marked contraction of the arterioles.

* His language is such as to leave the point somewhat doubtful. *Op. cit.*, pp. 188, 189.

† Quoted by Fothergill (*op. cit.*).

According to Boehm, Traube found that, if the spine be divided, digitalis is powerless to increase the arterial pressure, although lessening, as usual, the pulse-rate. The same authority also states that Bezold has seen an excessive fall of the arterial pressure ensue immediately upon the division of the spinal cord in an animal under the influence of digitalis. Further, in his own experiments Boehm has attained similar results, or, in other words, has found that after separation of the small vessels from the vaso-motor nerve-centre, digitalis does not increase arterial pressure.

These experiments would appear to prove that digitalis acts upon the vessels by stimulating the vaso-motor centres in the base of the brain; but they have been contradicted by Ackermann (*op. cit.*, p. 397), who states that he has many times cut the spinal cord and without exception found a very marked rise of arterial pressure follow the injection of digitalis. Unfortunately, none of these experiments have, that I am aware of, been published in detail, and it is therefore impossible to analyze or to reconcile them. It must be considered undetermined whether Professor Ackermann is or is not correct in asserting that digitalis acts directly upon the peripheral vessels themselves.

From the evidence which has been brought forward, it may be considered as definitely proven that in mammals digitalis in therapeutic doses is a powerful stimulant to the circulatory system.

The following proposition expresses our present knowledge, and probably is very close to the truth:

Digitalis in moderate doses stimulates the musculo-motor portion of the heart (probably its contained ganglia), increases the activity of the inhibitory apparatus, and causes contraction of the arterioles. As a consequence of the first action, the cardiac beats become much stronger; as the result of the last, there is narrowing of the blood-paths, and to the passage of the vital fluid an increased resistance, which, acting on the already excited inhibitory system, aids in the slowing of the pulse. Toxic doses of digitalis paralyze or weaken, more or less completely, each of the three systems, and cause rapidity of the pulse and falling of the arterial pressure.

According to my own experience, decided therapeutic doses of digitalis, in man as in other mammals, produce great reduction and sometimes dirotism of the pulse, and increase the size and force of the wave; at the same time the arterial tension is augmented. Poisonous doses induce, after a time, increase of the pulse-rate, with smallness and weakness of the wave and lowered arterial pressure.

Sphygmographic studies of the effect of digitalis upon persons suffering from various acute and chronic diseases have been made by M. Legroux, M. Bordier (*Bulletin Thérapeutique*, 1868, p. 110), Constantine Paul (*Bulletin Thérapeutique*, 1868, p. 193), and Paul Lorrain (*Journal de l'Anatomie et de la Physiologie*, 1870). The problems offered by these gentlemen are so complex as to render a detailed study almost impossible; but, as a whole, their tracings seem to confirm my personal experience. Paul Lorrain calls

attention to the fact that when the drug has reduced the pulse-rate very greatly a second abortive systole can, on auscultation, sometimes be heard occurring during the long diastole, and some of his sphygmographic tracings are markedly dicrotic. It is evident that in man the second systolic movement occurs precisely as in animals; and it seems very certain that the proposition framed for the lower mammals applies also to man.

When the pulse has been reduced by digitalis to forty or fifty a minute, the change from the recumbent to the erect position will not infrequently suffice to alter at once its character, so that it will become feeble, small, and one hundred and fifty per minute. The experiments of Traube, which have already been mentioned, afford an explanation of this phenomenon so simple that it can scarcely be else than true. The action of the drug in such a case is verging upon the point at which the pulse-rate increases and the arterial pressure falls, owing to the partial paralysis due to over-stimulation. Whilst the patient is recumbent, the line is not passed over, but the additional stimulation of the erect position carries the circulatory system beyond the limit of simple stimulation, and the over-effects of the drug are at once manifested.

The influence of digitalis upon the urinary secretion in health has been studied by numerous observers, with such diverse results as to prove that the action of the drug is inconsistent and varying. Thus, Jörg, Hammond (*Proc. Biol. Dept. Acad. Nat. Sciences*, Phila., Dec. 1858), and Brunton (*loc. cit.*) have found the secretion more or less decidedly increased, and Homolle (*Archives Générales*, July, 1861), Winogradoff (*Virchow's Archiv für Anatomie*, Bd. xxii., 1861), Stadion, and, according to Brunton, also Krahmer, Kluyskens, Vassal, and Shohl, have either found it uninfluenced or diminished.

The urea in the apparently very careful experiments of Winogradoff (*Virchow's Archiv*, Bd. xxii.), of Stadion (*Prager Vierteljahrsschrift für die Praktische Heilkunde*, 1862, Bd. lxxii.), and of Hammond (*loc. cit.*), was diminished, while in the almost equally elaborate experiments of Brunton (*loc. cit.*) it was increased. All three observers noted lessening of the chlorides.

It is very certain that *toxic* doses of digitalis lower the temperature in healthy men and animals a number of degrees. It would seem, however, that the fall of temperature is generally, if not always, preceded by a rise, as has been noted by Bouley and Reynal, by Duméril, Demarquay, and Lecoq (quoted by Brunton), by Hirtz, by Legros (*Thèse*, 1867, quoted by Gourvat), and by Gourvat (*Gazette Médicale de Paris*, 1871, p. 572).

The effect of *therapeutic* doses in the normal condition has not been closely studied, that I am aware of. The tendency of our present knowledge is to connect the changes in temperature induced by digitalis with the changes of the circulation; and it seems very possible that therapeutic doses in health will be found to increase bodily heat, although in fever they may diminish it.

THERAPEUTICS.—The chief clinical use of digitalis is in diseases of the heart; and it logically follows, from what has been said in regard to its physiological action, that it should be useful in loss of cardiac power.

When the muscle of the heart is for any reason unequal to the task set it, the systoles become rapid and imperfect, and by this irregular action, the ventricles neither completely filling nor emptying themselves, increase the embarrassment. Under these circumstances, digitalis, by lengthening the diastolic pauses and increasing the force of the systolic contractions, causes the ventricles to fill themselves in the one and to completely empty themselves in the other act. By subduing irregular action through the inhibitory nerves, by energizing the muscular power of the heart-walls, the remedy is of incalculable service, and, increasing arterial tension all over the body, causes the disappearance or the lessening of symptoms due to low pressure in the arteries.

It is a logical necessity, if our reasoning as to the physiological action of digitalis has led to a correct result, that the drug should be of the greatest service when the lesion is simply loss of cardiac power; and clinical experience tallies with this *a priori* argument. In *simple dilatation*, or in *simple failure of the cardiac muscle* without valvular lesion, the results of the use of digitalis are most favorable.

On the other hand, in *simple hypertrophy* digitalis does harm, and should never be used. It must be borne in mind that although this agrees with what the experimentalist has proven to be the action of digitalis, yet it was discovered independently as a clinical fact by practitioners. Thus, Niemeyer, who ridiculed experimental therapeutics because he would not take the trouble to study them deeply and practically, and was therefore incapable of understanding them,—Niemeyer says, “Digitalis in pure uncomplicated hypertrophy is unsuitable.”

Valvular lesion of the heart, as is well known, gives rise under unfavorable circumstances to dilatation, but in favorable cases to hypertrophy, or rather in the great majority of cases to hypertrophy with dilatation. Following out the principles already inculcated, it might seem at first that the use of digitalis in hypertrophied hearts with valvular lesion ought to be reprobated. Clinically, however, it is known that often digitalis does good in valvular lesion with enlargement of the heart. The results of logical deductions from our physiological conclusions as premises are not, however, at variance with this. It must be borne in mind that structural and functional hypertrophy are different things; by this is meant that although a heart be enlarged and absolutely stronger than normal, yet it may be, relatively to the work required of it, *weak*. Thus, if 1 represent the normal work of the heart and 1 its normal power, if the former be increased to 4 and the latter to 3 the heart is really in the position of a weak organ, although possessed of three times its original strength. Hence it is that digitalis is often useful in valvular disease with hypertrophy. In the vast majority of cases the heart with diseased valves is in the position just spoken of; but sometimes the work advances only to 2 and the strength to 3; then the hypertrophy becomes excessive, and digitalis will increase the difficulty. In almost all cases the increased power of hypertrophy, unless the muscle be degenerated, renders

effectual smaller doses than can be used in dilatation, and also increases the danger of the over-action of large doses.

In *mitral insufficiency* and in *mitral stenosis*, digitalis is often of great service. It is evident that in both instances the valvular lesion leads as its first result to pulmonic hyperæmia. How does the digitalis lessen this? In the case of *stenosis*, the diastole being lengthened by the remedy, the auricle is afforded more time to empty itself into the ventricle through the narrowed orifice, and at the same time is strengthened in its contracting power; evidently, then, the left ventricle when its systole occurs will have much more to contract on than before the digitalis was administered, and the amount of blood in the systemic circulation will be increased,—*i.e.*, the amount in the pulmonic circulation will be diminished; further, the right ventricle will have greater power afforded it to force the blood through the lungs,—*i.e.*, to resist the recoil from the left auricle to which the impeded valve gives origin.

In *mitral insufficiency* the mechanism is different, but the result is the same. The increased power of the systole will throw proportionately more blood through the aortic orifice than through the partially open valve. The opening at the insufficient mitral valve is much smaller and more obstructed than the aortic orifice. As the force or rapidity of the current increases under the action of digitalis, the friction becomes greater at both orifices, but the ratio of increase is evidently far higher in the small choked mitral leak than in the wide aortic opening. Hence the large orifice constantly gains upon the smaller as the cardiac force is increased, and, more blood passing into the systemic circulation, the pulmonic is relieved. Again, the right ventricle shares the stimulant action of the drug and acts more strongly upon the pulmonic circulation, resisting the direct backward flow from the auricle.

In *aortic constriction* digitalis is useful when the heart-power begins to fail. In these cases compensatory hypertrophy, with slowness of action, is very apt to occur, or even to become excessive: much more frequently does this happen than in mitral disease. In such cases digitalis does harm, and its use in them has given rise to the opinion that in aortic disease digitalis is not to be employed. It is scarcely necessary to show how both in *aortic insufficiency* and in *stenosis*, when the heart-muscle fails, when the hypertrophy is not compensatory, digitalis gives relief.

From the considerations which have been brought forward, it is very evident that a knowledge of the relation of the heart-muscle to the work required of it in any individual case is much more necessary to the therapist than to know what valve is diseased.

In "*irritable heart*" of soldiers, a disease or condition of cardiac irritability evidently connected with muscular weakness, and very probably dependent upon exhaustion of the inhibitory nerves, Dr. Da Costa (*American Journal of the Medical Sciences*, Jan. 1871) found that in the early stages of the affection digitalis not only acted better than any other remedy, but even, when

administered continuously for some time, often effected a permanent cure. When hypertrophy had taken place, the drug was of little use.

The relief afforded by digitalis in not too inveterate cardiac disease is often in a measure permanent, because the drug may aid very materially in the production of compensatory hypertrophy. Dilatation is certainly more apt to occur when the muscular fibre is lax and acting feebly than when it is toned up and in vigorous play; secondly, the stimulus to action in a muscle is almost of necessity directly or indirectly a stimulus to its nutrition; lastly, and most important, owing to the origin of the coronary arteries being so near the heart, and to the circulation in the cardiac walls being so intimately connected with the alternate pulse of the viscus, it is evident that improved systemic circulation means in a far more intense degree improved blood-supply to the cardiac muscle.

If in *aneurism*, or in general *capillary atheroma*, there be increased resistance to the circulation, and the heart have not sufficient power to meet this, digitalis may be useful, but must be employed with caution. It undoubtedly increases arterial pressure; and this increased pressure may prematurely rupture an atheromatous cerebral capillary or tear open the thinned wall of an aortic aneurism. The use of digitalis for the purpose of "quieting the circulation" in aneurism is of very doubtful expediency.

In *cardiac dropsy*, digitalis is of service probably not only by regulating through the heart the circulation, and by evacuating the surplus fluid through the kidneys, but also by an action upon the vessels. Clinicians do not allow enough for the rôle of the vaso-motor nerves in dropsy. Without saying more as to the clinical side of this question, which I have discussed elsewhere (*American Journal of the Medical Sciences*, July, 1871), it may be allowable to allude to the experiments of Ranvier (*Comptes-Rendus*, 1869, p. 1327), who found that when the vena cava was tied in a dog, and the sciatic nerve of one side cut, œdema occurred only in the leg whose nerve was divided.

The use of digitalis in large doses as a cardiac stimulant in *syncope* or in *sudden collapse* is probably in its infancy. One or two cases have been reported in which the happiest result has followed its exhibition. The great trouble is the slowness of its action; but this difficulty could probably be met by the hypodermic use of digitaline. I am not aware that this has been tried.

Closely allied to the last use of digitalis is its employment in *poisoning* by substances such as *muscaria*, *delphinia*, and *aconitia*, which arrest the heart in diastole. Boehm has shown (*Pflüger's Archiv*, Feb. 1872) that in digitalis-poisoning of the frog, even when systolic cardiac arrest has occurred, these substances will often restore the cardiac movements,—a proof that real antagonism exists in their action; and Dobie reports a case (*British Medical Journal*, Dec. 1872) of recovery after the ingestion of an ounce of Fleming's tincture of aconite, apparently due to the hypodermic injection of twenty minims of tincture of digitalis and the exhibition by the mouth of three doses

in an hour of a mixture of tincture of digitalis (one drachm each dose), brandy, and ammonia.

To maintain the heart's action in various acute diseases, such as *adynamic pneumonia* and *adynamic fevers*, digitalis is without doubt useful. It can have no effect upon the diseases themselves, but may help to sustain the heart during any crisis or a period of strain upon it. The clinical results obtained by H. M. Hankel (*British and Foreign Medico-Chirurg. Rev.*, vol. xxvi. p. 513) are certainly not encouraging: he noticed, however, in a number of cases that the pulse became more regular, fuller, and slower; but this of course is not of necessity any great gain.

With the idea that digitalis is an active *antipyretic*, it has been prescribed in various acute diseases, sometimes with asserted good results. As already stated, toxic doses of digitalis at first elevate the temperature; and proof is wanting that in healthy men therapeutic doses have any decided influence in depressing the temperature. There is, therefore, no good physiological basis for the antipyretic use of digitalis; at the same time, it is very possible that it may directly or indirectly lower the temperature in disease. Clinical proof of this is, however, still wanting. The strongest evidence in favor of such action is furnished by the records of Prof. Wunderlich (*Manual of Medical Thermometry*, Sydenham Soc. Translation, p. 325), according to which from half a drachm to a drachm of digitalis, given in divided dose during three or four days in the second or third week of severe *typhoid fever*, immediately produces a slight fall of temperature in a large proportion of the cases, and sometimes a considerable fall. This fall is said not to last more than a day, when the temperature rises again, but in cases favorably affected does not regain the original height; the pulse is very much lowered in frequency, and remains about uniform for four days. It is evident that at least in some of these cases of Wunderlich's the drug was given about the time natural defervescence would be expected to occur, and that the slight reduction of temperature brought about at such time does not argue very strongly in favor of the proposition that digitalis is a powerful antipyretic in disease.

Far more extensive and complete observations must be made upon a rising, not a falling, temperature, before any satisfactory conclusion can be reached. At present the antipyretic use of digitalis should be purely tentative.

The property of causing contraction of all unstripped muscular fibres has been attributed to digitalis, and whilst the probabilities are certainly such as to invite investigation, yet we have no definite knowledge upon the subject. Mr. Dickenson (*Medico-Chirurg. Trans.*, vol. xxix.) claims that it has a powerful action in causing the uterus to contract and to arrest hemorrhage,—a few minutes after an ounce and a half of the infusion is swallowed in menorrhagia, severe pains resembling those of the first stage of labor coming on, with a momentary profuse discharge of blood and clots, if there be any present, followed by arrest of the flow for hours. Stadion (*Sydenham Soc. Year Book*, 1862, p. 451) claims that digitalis is capable of temporarily but completely

annulling the activity of the sexual organs, and that it may be regarded as a true anaphrodisiac.

The use of digitalis as a diuretic will be considered under that heading.

TOXICOLOGY.—In poisoning by digitalis, the first symptom of any severity is generally vomiting of mucus and bile, very violent and very often repeated. At the same time a feeling of heat of the head, disordered vision, and vertigo manifest themselves. The pulse at this time in the horizontal position may be full and strong and slow, but on the patient's rising becomes weak and rapid. The face is pale. The vomiting continuing, profound prostration comes on, the pulse becomes feeble, small, irregular, although the beat of the heart may be strong and hard. The eyes are very prominent, the pupils fixed and dilated: according to Tardieu, an almost diagnostic symptom is the blue color of the sclerotic. Abundant salivation sometimes occurs. Intense headache and pains in the back or limbs are often complained of. Diarrhœa is very generally present; the urine may be suppressed. The intelligence is often perfect in the midst of profound collapse, but delirium more or less violent finally comes on. Death, usually preceded by stupor or by convulsions, takes place most frequently in one or two days, but has occurred as late as the tenth day, and as early as three-quarters of an hour.*

It is said that in some cases in which a cumulative action during the medical use of the drug has occurred the first marked symptom has been syncope, followed by paralysis of the lower portion of the body, vomiting, diarrhœa, delirium, general insensibility, and death.

In the majority of cases of digitalis-poisoning the patient recovers. When this happens, the symptoms gradually ameliorate. Cardiac weakness, and even a *bruit de souffle*, with more or less exophthalmos, is said to have persisted for weeks in some cases. In poisoning by digitaline, the symptoms are those of rapid digitalis-poisoning,—violent vomiting, intense cephalalgia, and sometimes rachalgia, irregular, feeble, intermittent pulse, paroxysms of suffocation.

The minimum fatal dose of digitalis is not known; a large teaspoonful of the tincture is said to have caused alarming symptoms in a young puerperal woman (Tardieu, *Clinique*, p. 685, Obs. VIII., Paris, 1867); twenty grains of the extract proved fatal on the tenth day (*Ibid.*, Obs. VI.); and two and a half grammes of the leaves in infusion on the fifth day (*Ibid.*, Obs. X.); fifty granules (gr. one-fiftieth?) of digitaline have been recovered from (*Ibid.*, Obs. XII., XIV.); in the only fatal case of digitaline-poisoning I know of (*Affaire Couty de la Pomerrais*), the amount ingested was unknown.

The treatment after the evacuation of the stomach and bowels, and the very free administration of tannic acid, as the best, although unreliable, chemical antidote, should consist in the exhibition of opium, of alcoholic stimulants, and rest in the horizontal position. I know of no recorded experiences with the antagonistic poisons to digitalis, such as aconite or muscarin. As it

* See case reported by M. Barth, quoted by Tardieu.

is possible that whilst so far as the heart is concerned they may be really antagonistic and yet may intensify the action of digitalis on the cord, their use requires caution.

ADMINISTRATION.—Digitalis may be given in substance in the form of pills; the dose being one grain three times a day, and increased until some effect is produced. When a rapid action is desired, one of the liquid preparations, such as the infusion, or else digitaline, should be used.

Infusum Digitalis—*Infusion of Digitalis* (℥i to Oss). Dose, one fluidrachm to half a fluidounce.

Tinctura Digitalis—*Tincture of Digitalis* (℥ii to Oj). Dose, five to twenty drops.

Extractum Digitalis—*Extract of Digitalis*. Dose, one-fourth grain.

In emergencies where single large doses are administered they may be very much larger than those here given. Thus, of the tincture two fluidrachms or even half an ounce may be exhibited; of the infusion, a wineglassful. These large doses, however, should, I think, be employed only by those who have had experience with digitalis given in the more ordinary way.

Occasionally when digitalis is steadily given three times a day continuously, an intense action will be suddenly developed. This probably arises from an accumulation of the medicine in the system, due to elimination not being so rapid as ingestion. Whatever the cause may be, of the fact there can be no doubt.*

For this reason, whenever digitalis is given continuously, at least every two weeks its exhibition should be suspended for three or four days.

* For a report of a case by the author, see *Philadelphia Medical Times*, vol. ii. p. 24.

CLASS IV.—CARDIAC SEDATIVES.


THERE are certain drugs which are used by practitioners to decrease the activity of the circulation; and it is these which are here considered under the heading of *Cardiac Sedatives*. Many, in fact all of them, possess other powers besides those which cause them to be considered under this caption, and none of them are in very close accord in these qualities. There is, however, a *general* resemblance in the action of such as are derived from the vegetable kingdom, in that they are all sedatives to the motor-nervous system and yet all produce convulsions. I have made an especial experimental study of these convulsions (*Philadelphia Medical Times*, vol. iii.), and have found that they are cerebral and not spinal, because they do not occur in any part of the body separated by section of the cord from cerebral influence. Further, they are probably due to disturbance of the circulation at the base of the brain, for the following reasons, the truth of each of which has been experimentally determined: first, lessening of the circulation at the base of the brain will cause convulsions; secondly, the convulsions produced by the cardiac sedatives do not occur until the arterial pressure is reduced about one-half; thirdly, if the disturbance of the cerebral circulation be artificially increased by tying the carotids previous to poisoning, or in any other way, the convulsions come on sooner and are more violent; fourthly, in some animals the convulsions caused by arresting circulation at the base of the brain are feeble and ill defined, whilst in others they are violent, and I have found that in species of the first order cardiac sedatives produce but slight convulsions, whilst in species of the second order they cause violent convulsions.

Of course the chief indication for a medicine of this class is the existence of sthenic arterial excitement; but these drugs vary so much that, like arterial stimulants, they must be studied as separate entities.

ANTIMONY. (Sb.)

The metal antimony is not officinal, but the following substances containing it are recognized by the U. S. Pharmacopœia.

ANTIMONII SULPHURETUM—SULPHURET OF ANTIMONY.



The native tersulphuret, which is included in the *Materia Medica* list as the basis out of which the preparations are made.

ANTIMONII OXIDUM—OXIDE OF ANTIMONY. ($\text{SbO}_3 - \text{Sb}_2\text{O}_3$.)

A grayish-white powder, insoluble in water, wholly soluble in muriatic or tartaric acid. It is prepared by dissolving the sulphuret of antimony in muriatic acid, adding nitric acid, and precipitating with water of ammonia. Its solution in the stomach is dependent upon the acid there present, and consequently, being uncertain in its action, it should not be used internally—although it is capable of producing all the effects of tartar emetic, for the preparation of which it was introduced into the Pharmacopœia.

ANTIMONII ET POTASSII TARTRAS—TARTRATE OF ANTIMONY AND POTASSIUM.

TARTAR EMETIC is prepared by boiling the oxide of antimony in a solution of bitartrate of potassium. It occurs in the form of a white powder, the result of the pulverization of transparent, colorless, slightly efflorescent crystals, which are most commonly rhombic octahedrons. Its taste is variously described: to me it is at first very slight, but after a time styptic and acrid. In some persons it blisters the tongue and lips after a few moments of contact.

Tartar emetic is insoluble in absolute, but soluble in dilute alcohol, soluble in from two to three parts of boiling water, and in from twelve to fifteen parts of water at ordinary temperatures.

It is incompatible with alkalies and with acids, including tannic acid and substances containing it.

PHYSIOLOGICAL ACTION.—Locally applied, tartar emetic is an irritant, acting upon some very delicate and susceptible skins in a very short time. In most instances, however, its continuous application for several days is necessary to produce any effect. At first there is simply a redness, accompanied by some burning pain and the eruption of small papules, which shortly become converted into vesicles and then into pustules. These are irregular in shape and size, varying from one-eighth of an inch to an inch and a half in diameter, and are very painful. Sometimes these pustules give rise to small sloughs, but generally, if the application be withdrawn, they simply give origin to superficial ulcers, which readily heal.

The only symptoms which are induced by small therapeutic doses ($\frac{1}{12}$ gr.) of tartar emetic when exhibited for a short time are a scarcely perceptible diminution of the force of the pulse and an increase of the perspiration.

By somewhat larger amounts of the drug, nausea is induced, and is accompanied in a more decided degree by the phenomena just mentioned. After large doses, prolonged nausea, violent vomiting and retching, with marked reduction of the force of the pulse, great muscular relaxation, and a feeling of faintness, occur. At the same time the saliva is generally increased in amount, and the skin is bedewed with sweat.

By poisonous doses all these symptoms are intensified. The vomiting is

violent, repeated, continuously re-excited by the slightest provocation, and is accompanied by burning in the œsophagus and stomach and by colicky pains in the abdomen. The matters vomited are first mucus, then mucus and bile, and finally, in some cases, blood. With the gastric disturbance occurs violent and frequent serous purging, the discharges resembling those of cholera, but becoming in some cases towards the last bloody. Cramps may occur in the extremity, and, in conjunction with the serous purging, have caused the antimonial poisoning to be mistaken for cholera. The exhaustion is extreme, and finally ends in collapse, with thready or imperceptible pulse, pinched, livid countenance, suppressed voice, profuse cold sweats, lowered temperature, and at last death from asthenia, generally preceded by stupor or convulsions: indeed, Taylor reports cases in which wild delirium was present some hours before death. The urine* in mild cases is increased in quantity, as it is also in the beginning even in fatal cases, but in such towards the close it is generally scanty and bloody, and even suppressed.

It is evident that the symptoms just enumerated can be best studied in detail under several heads. Before entering upon this, however, it is well to premise that the experiments of Viborg (*Stillé's Therapeutics*, vol. ii.), Buchheim, Courten (*Ibid.*), Magendie (*Ibid.*), Ackermann (*Virchow's Archiv*, Bd. xxv. p. 531), Richardson (*London Lancet*, May, 1856), Nöbiling (*Schmidt's Jahrbücher*, Bd. cxl. p. 24), and Radziejewski (*Reichert's Archiv für Anatomie*, 1871), and of others, have demonstrated that tartar emetic acts upon the lower animals precisely as on man.

When a sufficient dose of tartar emetic is injected into the frog (Radziejewski, Ackermann, Nöbiling), the cardiac contractions in a very short time are lessened in frequency and force, and become irregular, the auricles pulsating more frequently than the ventricles, until finally arrest occurs in diastole. After death the irritability of the cardiac muscle is almost, or more frequently entirely, destroyed.

Upon the heart of the mammal the drug acts as upon that of the frog. According to the researches of Ackermann and of Ernst Sentz (*Experimenta de Ratione Interpulsus, etc.*, Diss. Inaug., Dorp. Livon., 1853), the arterial pressure always falls steadily and to an extreme degree. The pulse sometimes seems accelerated at first, but in the great majority of cases is decreased very decidedly in its rate. During this period of slow pulse the diastolic pauses are extremely long, but the individual beat will influence the mercurial column of the cardiometer five times as much as normal.

After a while the pulse suddenly becomes very rapid, the force of the heart-beat is almost completely lost, the arterial pressure falls to a minimum, and

* What is said in the text is, I think, correct, although authorities differ on this point. Trousseau (*Traité thérapeutique*, 4th ed., vol. i. p. 619) affirms that it is suppressed; Husemann, that it *never* is suppressed (*Toxicologie*, p. 854); Tardieu, that it is scanty. For a case in which it *was* suppressed, see Taylor's *Medical Jurisprudence*, London, 1873, p. 309.

in a very few moments diastolic arrest occurs. It is evident that the action of antimony upon the heart is a direct one. The irritability of the muscle is lost, and Ackermann has found that the cut-out heart of the frog is affected by the solution of tartar emetic; further, the experiments of Radziejewski have proven that the peripheral ends of the vagi in antimonial poisoning are early more or less completely paralyzed, so that the diastolic arrest cannot be due to excitation of the inhibitory apparatus. Perhaps the present is as fitting as any other position to notice the theory of Nöbiling, that the action of tartar emetic upon the heart is owing to the potash it contains. This theory in itself is so improbable that it would seem scarcely worthy of discussion, were it not for the fact that Nöbiling asserts that the tartrate of antimony and soda is not poisonous. Dr. Radziejewski (*loc. cit.*, 485) has repeated and extended the experiments of Nöbiling and completely disproved both the asserted fact and the theory based upon it, showing that the soda-salt is as poisonous as the potash-salt.

A prominent symptom in antimonial poisoning is paralysis, affecting to an extraordinary degree the sensory and to a less extent the motor system. In man the anaesthesia which occurs in animals has been overlooked, but in the advanced stages of poisoning it is no doubt present. According to Radziejewski, the paralysis and diminution of reflex action are of spinal origin, as they occur when the Setschenow's centre is previously separated from the cord as well as when access of the poison to an extremity has been prevented by tying the artery. The same authority states that the development of the paralysis occurs in the following order: sensibility towards thermic and chemical irritants, then towards tactile stimuli, then towards locomotion and reflex action. Thus, a rabbit which could still drag itself around suffered its paws to be deeply burned without evincing the slightest evidence of feeling. The contractility of the voluntary muscles is not materially affected.

The influence of antimony upon the temperature appears not to be very marked when the drug is exhibited in ordinary therapeutic doses. Thus, Ackermann found that, after doses severe enough to induce violent vomiting, no alteration in the temperature could be discovered by the thermometer under the tongue. Owing no doubt to the disturbance of nervous and arterial action, there is in these cases, however, a very marked reduction of the temperature of the extremities. Thus, in the cases just alluded to (Ackermann) the heat of the hands was lowered from 0.2° C. to 3.5° C. in various persons.

After poisonous doses of antimony the decrease of animal heat is very perceptible, provided the victim live sufficiently long. Thus, in Ackermann's experiments a fall of only 1.6° C. occurred in rabbits killed in the hour, but in those that lived five hours the depression amounted to 6.6° C.

It cannot be gainsaid that tartar emetic acts as an irritant upon the alimentary mucous membrane. Although cases (*Archives G n rales*, Sept. 1865) have been reported in which no lesion has been found in the stomach or bowels

after death from antimony, yet in the great majority of instances very decided indications of violent inflammation have been present.

Dr. Radziejewski, on the strength of this action, and of two experiments in which he found the greater portion of the ingested antimony in the vomit of the patient, has advanced the theory that the emesis is due to a local action of the drug. The persistent nausea, however, certainly indicates that the remedy does not act like the so-called mechanical emetics. Further, the vomiting induced in the experiment of Magendie, of replacing the stomach of an animal by a bladder and giving tartar emetic, would seem to settle the point completely. This experiment of Magendie has been confirmed by Brinton,* of London, who further proved that when tartar emetic was injected into the vein of an animal it was very freely and rapidly eliminated by the stomach. Dr. B. W. Richardson (*London Lancet*, vol. i., 1856) has corroborated this, and has also found that a similar elimination follows the inhalation of antimonietted hydrogen.

I think, therefore, it must be conceded—first, that the finding of even a large quantity of antimony in the vomit does not prove that its action on the stomach is chiefly a local one; secondly, that the emesis is certainly preceded by absorption, and is probably due to an impression upon the nerve-centres. The purging induced by tartar emetic is without doubt of the same nature as the vomiting, and is an effort at elimination. The respiration in poisoning by antimony is very irregular, with all sorts of variations in the rhythm of the act. In the advanced stages the pauses are often very long, and the inspiration and expiration so forced and prolonged that very generally, in animals at least, marginal emphysema and subpleural ecchymoses are found after death. The origin of the respiratory trouble is probably somewhat complex, the chief factor being the direct influence of the drug upon the respiratory nerve-centres, and minor causes the intense venous congestion due to the failure of the circulation, and the alteration of the blood itself.

Antimony acts directly or indirectly upon the mucous membrane of the lungs, even when given in moderate doses, as is shown by clinical experience and by the experiments of Mayerhofer (*Nothnagel's Arzneimittellehre*, Berlin, 1870, p. 219).

THERAPEUTICS.—There are three indications to meet which tartar emetic is constantly employed. The first of these it fulfils by virtue of its powers as an emetic. The discussion of this may be found in the chapter upon Emetics.

The second purpose for which antimony is used is to depress arterial excitement. It is chiefly in inflammation that tartar emetic is used as an arterial sedative. In combination with more decided diaphoretics it is constantly employed by some surgeons in fever after operations, in gonorrhœa, and in various sthenic inflammatory affections. In pneumonia it has been very largely used, forming an essential portion of the older plan of treating

* *Cyclopædia of Anatomy*, Supplement, p. 319; *London Lancet*, 1853, vol. ii. p. 599.

that disease. According to the method of Rasori, four or five grains a day were at first given, but rapidly increased to twenty-four or even thirty grains daily. Although by the aid of opiates and careful dilution a species of tolerance was often obtained for these heroic doses, yet very properly the plan has been abandoned by modern therapeutists.

As tartar emetic if administered in sufficient quantity to depress markedly the circulation causes generally intense nausea and often purging, I think it is inferior to aconite or veratrum viride when it is desired to depress the circulation very decidedly for any purpose.

Owing to its action upon the mucous membrane of the bronchial tubes, in the first stages of *bronchitis* tartar emetic is an invaluable remedy. After free secretion has been established, other expectorants are, I think, of more value. The value of antimony as a *diaphoretic* depends largely upon its action on the circulation. Minute doses of it are constantly employed to increase the efficiency of fever-mixtures. It must always be borne in mind that it is a powerful depressant, and is therefore to be employed only in sthenic cases.

As a *counter-irritant*, tartar emetic is used only when it is desired to produce a slow, persistent, and at the same time very decided impression. For further discussion of its application to disease, see chapter on Rubefacients.

TOXICOLOGY.—The general symptoms produced by poisonous doses of antimony have been sufficiently described. There is, however, according to authors, a form of antimonial poisoning in which neither vomiting nor purging* occurs, the symptoms being simply intense prostration, cold clammy sweat, a sense of oppression in the chest, with the respiration at first increased, then diminished in frequency and embarrassed; a rapid feeble pulse, after a time becoming slow, intermittent, and irregular; delirium, unconsciousness, tremblings, and clonic and tonic convulsions (Husemann, *Toxicologie*, p. 853).

Tardieu (*loc. cit.*, p. 608) states that in some cases of tartar emetic poisoning a rash exactly resembling that produced by the external application of the drug has appeared all over the body on the fourth or fifth day.

As already stated, in the vast majority of cases there are to be found, after death from antimonial poisoning, very decided traces of inflammation of the stomach and bowels; in some cases, however, these appear to be wanting. The venous system is generally very much engorged, and the viscera are intensely congested. Magendie asserted that in animals poisoned by tartar emetic the lungs are always full of portions apparently hepatized; but Ackermann (*loc. cit.*, p. 544), in twenty experiments, found only some marginal emphysema and subpleural ecchymoses, with, in one or two cases, spots of atelectasis in the lungs.

It is evident, therefore, that the assertion of Magendie is too sweeping; but

* Husemann states this: although vomiting is absent in these cases, purging is generally present. I do not remember to have seen the report of a case in which it was absent.

it is true that, in a large proportion of fatal cases of antimonial poisoning, emphysema, pulmonary apoplexy, atelectasis, or other structural lesions of the lungs exist. The blood usually coagulates imperfectly.

Dr. Saikowsky (*Virchow's Archiv*, Bd. xxxiv. p. 78, 1865), of Moscow, has found that when animals are fed upon antimonie acid (one-half to one gramme daily) or other preparations of the metal for from fourteen to nineteen days, the liver, kidneys, and even the heart undergo a fatty degeneration; also that there is a lessening of the amount of glycogen in the liver, and in some cases even a total disappearance of it. This has been confirmed by Profs. Grohe and Mosler, who state that in the duchy of Brunswick the peasantry give to the geese, when producing the famous fatty livers, a certain quantity of the white oxide of antimony every day.

The minimum fatal dose of tartar emetic is not known. Three-quarters of a grain in a child, and two grains in an adult, have proved fatal; but in the latter case extrinsic circumstances favored the result (Taylor, *Guy's Hospital Reports*, Oct. 1857, an analysis of thirty-seven fatal cases); two hundred grains have been recovered from (case, *Taylor's Medical Jurisprudence*, 1873, p. 309).

Chronic Poisoning.—According to Mayerhofer (*Heller's Archiv*, 1846, quoted by Taylor), the symptoms following the criminal administration of small doses of tartar emetic at intervals are nausea, mucous and bilious vomiting, watery purging, often followed by constipation, small frequent pulse, and asthenia, deepening into death from exhaustion.

The treatment of antimonial poisoning consists in washing out the alimentary canal with large draughts of *tannic acid*,—the best known antidote,—and in the use of opium and of internal and external stimulants.

VERATRUM VIRIDE.

The root-stock of *Veratrum viride*, a coarse perennial herbal plant, indigenous to the Northern United States. It is a large tapering rhizome, an inch or two in length, less than an inch in thickness at the base, and having a bitter acrid taste. It contains the alkaloids *viridia* and *veratroidia*. *Viridia* is so closely united with the inert resin that it is separated from it with great difficulty; in the root-stock it is the more abundant alkaloid of the two.

According to their discoverer, Mr. Chas. Bullock, the alkaloids are almost identical in their reactions (*Proceedings of the American Pharmaceutical Association*, 1867), but are to be distinguished by their solubilities, *veratroidia* being soluble, *viridia* insoluble, in ether. *Veratroidia*, which in causing sneezing when inhaled, and in other properties, resembles *veratria* more closely than does its congener, is distinguished from the *veratria* by its higher melting-point (270° to 275° F.), by its producing intense redness in contact with concentrated sulphuric acid, and by not answering to Trapp's test.

PHYSIOLOGICAL ACTION.—In treating of the physiological action of *veratrum viride*, I shall first speak of the effects of its alkaloids singly.

When an animal is poisoned with *viridia*, the first symptom manifested is sluggishness, as shown by a disposition to be quiet, accompanied by distinct signs of muscular weakness. In a little while peculiar rapidly-repeated thrills run through the muscular system, so that the animal trembles violently. After a greater or less length of time the animal becomes unable to stand, from weakness, and at or before this period violent convulsions appear,—general clonic spasms without rigidity. The convulsions alternate with intervals of relaxation, and as the animal grows more profoundly prostrated are less severe, but they continue in most cases up to death. Even when they are most violent, force is evidently wanting. The animal is totally unable to raise himself from the ground; the pigeon drives himself forward upon his breast, the rabbit on his belly, or lies upon his side and kicks into the air. Sensation appears to be benumbed only very late in the poisoning, and consciousness is preserved almost to the last. The pupils are not affected. There is no purging or vomiting, but always profuse salivation. Respiration ceases before cardiac action, so that death probably takes place from asphyxia.

The circulation is profoundly affected. The pulse is generally, if not always, lessened in frequency, *provided the animal be quiet*. When there are convulsions, or even when the tremors are marked, it becomes very rapid. The arterial pressure is greatly lowered, falling progressively from the beginning to the end of the poisoning. The force of the individual beat appears not to be much altered at first.

It has not yet been experimentally determined exactly how *viridia* depresses the circulation; but, as I have found that previous section of the vagi does not influence the cardiac action of this alkaloid, and that when it is placed directly upon the heart of the turtle it acts at once and powerfully, the probabilities are that it influences the heart itself or its contained ganglia.

I am not acquainted with any experimental facts showing how the drug affects the vaso-motor nerves: the probability is that it paralyzes them; but actual proof of this is wanting.

In frogs, as well as in the higher animals, poisoned with *viridia*, there is a very marked diminution and finally abolition of reflex activity; and, as the functions of neither peripheral nerves nor muscles are interfered with, it is evident that *viridia* is an intensely powerful spinal depressant.* Locally, *viridia* is very feebly if at all irritant.

The general symptoms induced by *veratroidia* resemble those caused by its

* On account of the violence of the convulsions, it has been asserted that *viridia* is a spinal excitant. The convulsions are, however, really of brain and not of spinal origin, as they do not follow poisoning with *veratrum viride* when the cord has been previously cut. See *Philadelphia Medical Times*, vol. iii. Dr. Peugnet has also denied that *viridia* has any influence upon the circulation. As, however, in his experiments he did not use any form of cardiometer, and consequently did not really study the circulation, I do not think any weight should be attached either to his observations or to the conclusions which he draws from them. For a discussion of the matter, see the *Philadelphia Medical Times*, vols. ii. and iii.

congeneric alkaloid, but it is decidedly more irritating than the latter, and always induces vomiting, and occasionally purging. In poisoning by it there are in most cases some muscular twitchings, and finally marked convulsions, but neither of these are so severe and so repeated as in the case of viridia. Death takes place from asphyxia, due to paralysis of the respiratory muscles.

Upon the spinal cord, the peripheral nerves, and the muscles, veratroidia acts very much as does viridia, being a decided spinal depressant.

The action of veratroidia upon the circulation is less pronounced than that of viridia, but has not been at all closely studied. A partial investigation which I made yielded very curious results. The rapidity of the pulse and the arterial pressure were at first decidedly lessened. After a time, the pulse still remaining very slow, the individual heart-beats became endowed with four times their pristine force, and the arterial pressure became normal; then suddenly the pulse-rate quintupled itself, the individual cardiac beats losing much of their extraordinary vigor, but the arterial pressure rising nearly fifty per cent. The most plausible explanation of this series of phenomena is that the latter part of it was due to the influence of the sudden asphyxia induced by the spinal action of the drug. (See *American Journal of the Medical Sciences*, Jan. 1870, p. 47.)

The *resin* of veratrum viride, when completely deprived of the alkaloids, is nearly inert. It seems, however, to be irritating to the digestive organs, and very probably aids in the production of the vomiting occasioned by full doses of the drug.

As the action of the alkaloids of veratrum viride is very similar, and as they are the only active principles of the drug, it is very easy *a priori* to determine what the influence of the drug will be. Sufficiently numerous experiments* have been performed with the crude drug, or its preparations, to show that it acts upon the lower animals as upon man; but it is not necessary here to do more than to allude to them. When taken in small doses by man, veratrum viride first reduces the force without much lessening the frequency of the pulse, but after a time the pulse falls very much in rapidity, sometimes, according to Dr. Norwood, even to thirty-five a minute.

If any exertion be made during this stage of depression, the slow pulse will be suddenly converted into an exceedingly rapid one. The slow pulse is sometimes moderately full, but is always very soft and compressible; the rapid pulse is exceedingly feeble and small, often thready, and may become imperceptible. Severe nausea and vomiting accompany or follow the reduction of the pulse-rate. That the latter is not due to gastric disturbance is, however, shown by the fact that it often precedes, and may exist without, stomacheic symptoms. Thus, Prof. Percy states that he has seen the pulse reduced to thirty per minute without nausea being induced.

* See especially a paper by Prof. S. R. Percy, *Transactions of the American Medical Association*. Reprinted as pamphlet, 1864.

During the stage of depression there is always decided muscular weakness and relaxation.

After a poisonous dose the symptoms above noted are increased in intensity and become very alarming. A running, almost imperceptible pulse,—a cold, clammy skin,—intense nausea, and incessant attempts at vomiting, or retching, or hiccough,—absolute muscular prostration,—faintness,—vertigo,—loss of vision, and semi-unconsciousness, make up the group of extreme symptoms. Various observers also speak of an excruciating præcordial pain; but this I have not seen.

From these symptoms, with what has already been said in regard to the alkaloids, it follows that veratrum viride is a powerful spinal and arterial depressant, exerting little or no direct influence upon the cerebral centres. Under its action the functional activity of the skin is greatly increased, but, as this is a necessary result of the profound arterial depression, there is no reason for believing that the drug has any specific influence upon the perspiratory glands. In a similar manner the excretion of bile is often indirectly increased by veratrum viride, through the severe vomiting which it induces.

No studies have been made as to the action of the alkaloids of veratrum viride upon the vaso-motor system; but Prof. S. R. Percy states that a dilatation of the blood-vessels of the frog's web and bat's wing can be readily seen by the microscope to follow the administration of the drug. As he did not measure these changes, but merely judged of them by the eye, I think the most that can be claimed for them is that they render still more probable what the great lowering of the force of the circulation indicates, namely, that veratrum viride directly or indirectly paralyzes the vessels.

Although careful complete studies of the action of veratrum viride upon animal heat are wanting, yet it cannot be doubted that in health and in disease it lowers the bodily temperature; in what way or to what extent is uncertain.

M. Linon (*Gazette Médicale de Strasbourg*, quoted in the *Bulletin Thérapeutique*, 1869, tome lxxvi. p. 95) states that it abates the temperature, but not so certainly as it does the pulse. Oulmont (*Bulletin Thérapeutique*, 1868, tome lxxiv. p. 153) asserts, as the results of his experiments, that in animals from half an hour to two hours after the administration of such doses as would produce violent symptoms without killing, the temperature fell 2°, 3°, or even 5° (C.?), and remained at this point for twenty-four hours.

THERAPEUTICS.—With our present knowledge of the physiological action of veratrum viride, it is evident that there are only two rational indications for its use, namely, to *reduce spinal action* and to *reduce arterial action*. Owing to the very great effect veratrum viride has upon the circulation, and the numerous drugs which are purer spinal depressants, it is never called for to meet the first indication, and in practice should simply be used to lessen the force of the circulation. The use of the drug in *typhoid fever* and other adynamic diseases is simply an irrational and dangerous practice, founded upon an erroneous idea of the action of the remedy.

Veratrum viride has been recommended in *mania a potu*; and in cases of irritation of the brain from drink, with strong bounding pulse, it may be of great service; but in the true *delirium tremens*, with universal adynamia, it is a thoroughly improper remedy, capable of deepening the prostration into fatal exhaustion: indeed, I have known of death occurring in this disease from its use.

When true sthenic arterial excitement is to be combated in any disease, except it be *gastritis*, *veratrum viride* may be employed as a prompt, thoroughly efficient, and at the same time very safe remedy,—very safe, since it is almost incapable of producing death in the robust adult, unless used with great recklessness and in repeated doses.

In the early stages of *sthenic pneumonia* it offers, I believe, the best known method of reducing the pulse-rate and the temperature, and of ameliorating the disease.*

It is hardly necessary to mention other individual diseases in which *veratrum viride* may be employed to carry out the present indication.

In *peritonitis* its tendency to cause vomiting is very much against its use, and, unless this action can be controlled, should interdict its employment. I desire, however, to call attention to its value in preventing inflammation after severe *abdominal injuries*,—indeed, after any severe injury. Thus, I am cognizant of the case of a woman whose belly was torn open by the horn of a bull; the abdominal walls were rent for about six inches, and the sigmoid flexure of the colon came out and was dragged in the dirt. It was washed, replaced, the wound sewed up, the patient restricted to low diet, and *veratrum viride* administered very carefully so as to keep the pulse as low as possible and at the same time to avoid vomiting, to aid in which opium was also given. Recovery without a bad symptom resulted.†

As an emetic, *veratrum viride* should never be employed.

In chronic *cardiac diseases* it may be used in precisely those cases in which *digitalis* is contra-indicated,—*i.e.*, where there is excessive hypertrophy.

The contra-indications to the use of *veratrum viride* are cardiac weakness or the existence of general adynamia.

TOXICOLOGY.—Although *veratrum viride* is a remedy of great power, capable of producing the most alarming symptoms, yet I believe it to be the safest of all the cardiac depressants; certainly it is far less dangerous than *aconite*. Overdoses of it produce vomiting so soon and so certainly that it is somewhat doubtful whether a robust adult could be killed by a single dose of any of its officinal preparations, especially if prompt and judicious treatment were afforded.

I have several times known a teaspoonful of its fluid extract to be taken, and Prof. Percy cites recoveries after the ingestion of a tumblerful of the

* Compare Oulmont, *Bulletin Thérapeutique*, t. lxxiv. p. 146, and MM. Zuber and H. Hirtz, *Ibid.*, t. lxxvi. p. 468.

† Consult also Dr. C. S. Bishop, *American Journal of the Medical Sciences*, Oct. 1861.

tincture; after thirty grains of the resinoid; after two doses—a tumblerful each—of a syrup representing a pound of the root to the pint. The only death as yet reported is that of a feeble child, eighteen months old, killed by thirty-five drops of the tincture (*American Journal of Pharmacy*, Sept. 1865).

I have seen the most alarming symptoms result from large medicinal doses repeated at short intervals, and have been astonished at the rapidity with which they yielded to treatment.

In cases of poisoning, vomiting should be encouraged by large draughts of warm water until the stomach is well washed out. Then the patient should be forced to lie flat upon the back, with the head lower than the feet, and the efforts at vomiting should be restrained. If they cannot be checked, and if the prostration be severe, on no account should the patient be allowed to rise up, but must be made to vomit into a towel. A full dose of laudanum should be given by the rectum, and brandy or whisky be administered by the mouth. I have noticed that spirits will sometimes be retained only when given undiluted, and in such form will quiet the stomach at once.

If the stomach refuse alcohol in any shape, the rectum should be made use of. Ammonia may be employed as an adjuvant to alcohol, and in extreme cases should be injected hypodermically, or even into a vein. The use of external heat is important, and mild flagellations, rubbing with coarse towels, sinapisms, etc., may be used to keep up the external capillary circulation.

ADMINISTRATION.—In administering *veratrum viride*, it should always be borne in mind that it will do no good in acute disease unless given in increasing doses until its physiological action is manifested. In almost all cases vomiting is to be avoided as far as possible. To do this, small quantities of the drug should be given at short intervals, and corresponding doses of laudanum (five to ten drops) should be exhibited fifteen minutes before each dose of the *veratrum viride*. An hour is generally the best interval between the doses. The drug should always be administered in the following officinal preparations:

Extractum Veratri Viridis Fluidum—*Fluid Extract of Veratrum Viride*.

Dose, one to three drops.

Tinctura Veratri Viridis—*Tincture of Veratrum Viride* (3viii to Oj).

Dose, three to six drops.

The United States Pharmacopœia still retains in its officinal list the root-stock of the *VERATRUM ALBUM*, of Europe—for what reason it is hard to say, as the drug is never, to my knowledge, used by regular practitioners in this country. The extremely close botanical relationship which exists between *veratrum album* and *veratrum viride* would indicate similarity of physiological effects; yet, though this likeness does exist, there is enough difference to cause the entire rejection of *veratrum album* by the therapist. According to Oulmont, the chief difference in their effects upon animals is in the intensity of the action of the *veratrum album* upon the alimentary canal, and the

induction by it of violent inflammation of the whole alimentary mucous membrane; also death is much more apt to result from *veratrum album* than from *veratrum viride*.

In cases of poisoning with *veratrum album* the symptoms* have been—excessive vomiting, generally accompanied by severe abdominal, and often œsophageal, pain, and followed by a very severe diarrhœa; intense prostration and muscular relaxation; very pronounced reduction of the temperature and pulse, the latter being sometimes rapid and almost imperceptible in the advanced stages, and finally becoming extinct; sunken eyes, contracted, anxious countenance, a cold skin clammy with profuse perspiration, and other evidences of collapse. The mind remains clear until the last. When recovery occurs, it may be through a protracted convalescence.

The exact nature of the active principles of *veratrum album* is still involved in doubt. Pelletier and Caventou thought that they found supergallate of veratria in it. Simon asserted that it contained two new alkaloids, *barytina* and *jervina*, and recently Peugnet has affirmed that these are respectively the veratroidia and viridia of Bullock, and that the differences in the physiological action of *veratrum album* and *veratrum viride* are produced by the former containing not only much the larger proportion of veratroidia, but also an active resin. It appears to me very possible, if not probable, that the conclusions of Dr. Peugnet are not correct, and that *barytina* and *jervina* are closely related to, but not identical with, veratroidia and viridia. My reasons are as follows:

1st. Mr. Bullock states expressly (see *United States Dispensatory*, note to page 888, 13th edition) that veratroidia does not respond to Trapp's test for veratria, whilst Dr. Peugnet (*loc. cit.*) affirms that the corresponding alkaloid of *veratrum album* does so respond: consequently, there is either inaccuracy of observation upon one side or the other, or else the two alkaloids are not identical.

2d. The fusing-point of viridia is stated by Bullock to be from 335° to 340° F.; that of *jervina*, according to Dr. Peugnet, 360° F.

3d. The investigations of Dr. Peugnet were not carried far enough to prove the identity of the two alkaloids.

4th. So far as I can make out from the authorities at my command, Simon† claims that there are three alkaloids in the *veratrum album*,—veratria, *barytina*, and *jervina*; and Dr. Mossel (*Sur la Véatrine*, Thèse, Paris, 1868) certainly indicates that *sabadillia* and *barytina* are the same.

The matter undoubtedly needs further investigation before it can be considered settled.

* For cases and analysis of symptoms, see Dr. Peugnet's paper in the *New York Medical Record*, p. 121, 1872.

† I have not had access to Simon's original papers.

SABADILLA.

The seeds of *Veratrum sabadilla* (*Asagraea officinalis*), a plant growing in Mexico.

These seeds occur, in commerce, mixed with the fruit, which is formed of three coalescing capsules, somewhat resembling, although smaller, the fruit of the barley. The seeds themselves are very small, hard, blackish or brownish, slightly winged, inodorous, with a very acrid, burning, persistent taste. They contain gallate of veratria, and are officinal for the manufacture of the alkaloid.

VERATRIA.

The veratria of commerce is almost always more or less impure, and occurs as a grayish-white powder of an intensely acrid taste, and producing, even in the minutest quantity, when smelled, frequently repeated sneezing, which may continue for hours. It has when pure been considered uncrystallizable, but Merck has obtained it in rhombic prisms of about half an inch in length, through the spontaneous evaporation of its alcoholic solution. It is very slightly soluble in boiling water, not at all in cold water; soluble in alcohol, freely so in ether, and still more so in dilute acids.

Veratria dissolves in concentrated sulphuric acid, with the production of a yellow color, changing in five minutes into orange, then into blood red, and in half an hour into a splendid carmine. Masing states that this test is very faint with 0.0026 of a grain. If some bromine be dropped into the freshly-prepared sulphuric acid solution, a beautiful purple results. A more delicate test than either of those yet noted is, according to Masing, that of Trapp, which consists in warming the colorless solution of veratria in concentrated muriatic acid, when a dark-red, very persistent color is produced. This test is said to afford very marked proof of the presence of 0.0026 of a grain of the alkaloid, and to be especially useful when the veratria is impure.

PHYSIOLOGICAL ACTION.—Veratria is exceedingly irritating to any surface it may come in contact with, producing when given hypodermically or endermically severe pain, and when rubbed on the skin a feeling of warmth, followed by prickling, severe pain, numbness, and, if its use be persisted in, a marked redness. On the mucous membranes its action is even more decided. In the nostrils the minutest portion of it produces intense irritation, as shown by repeated sneezing and free discharge, which may be bloody. Upon the tongue a speck causes burning, with free salivation.

When taken internally, in small doses, it produces slowing and weakening of the pulse; more freely administered, indications of gastro-intestinal irritation; and in large doses it is followed by violent vomiting, serous purging, often with intense burning in the mouth and throat, and general muscular weakness. No fatal case of poisoning is on record; but in the experiments of Esche on himself a half-grain of the acetate produced collapse, with a pale, cold, wet skin, pinched features, a rapid, thready, irregular pulse, violent vomiting,

and marked muscular tremblings. Other observers have noted more pronounced indications of convulsions; and according to Bardsley, when absorbed through the skin, instead of purging it produces in some cases very free diuresis. On the whole, the resemblance between the symptoms as induced in man and in the lower animals is, so far as we know, complete.

The phenomena of veratria-poisoning in a mammal are violent muscular twitchings and convulsions, which are often plainly excited by external irritants, severe vomiting, generally but not always accompanied by purging, and disturbance of motion, respiration, and circulation. The pulse is at first, if the dose be not too large, quickened and strengthened, but in a very short time it becomes slower and weaker, and finally very frequent, thready, and irregular. Early there is a marked loss of muscular power, even in the midst of the convulsions, and the latter may give way to the quiet of paralysis, or may continue up to death.

M. Prevost (*Robin's Journal de l'Anatomie*, 1868, t. v. p. 206) has, I think, very well divided the action of veratria in poisonous doses into three stages: first, that of excitation or restlessness; second, that of convulsions; third, that of paralysis. It should, however, be understood that these may after large doses be fused into one. I have seen an animal suffer a convulsion, or perhaps merely give a convulsive shudder, and drop dead.

After death from a very large dose, the muscles are found to have lost more or less completely their irritability, so that they either do not respond, or respond very feebly, to the strongest faradaic currents. That this is due to a direct influence of the alkaloid upon them is proven by the fact, first noted by Kölliker (*Virchow's Archiv*, Bd. x. p. 257), but which I in common with other observers have experimentally confirmed, that if an artery be tied before poisoning, all the muscles supplied by such artery maintain their integrity.

It is evident that veratria is a muscle-poison; but it has other powers, and the subject is best studied in detail system by system.

Central Nervous System.—Upon the cerebrum the action of veratria is not very marked. That the convulsions are not cerebral is shown by the fact, which I have frequently noted, that they are in no wise affected by division of the spinal cord. The spasms must be, therefore, either peripheral or spinal in origin. M. Prevost (*Robin's Journal de l'Anatomie*, 1868, p. 209) has found that convulsions will occur in the frog even when the spinal cord is destroyed, but that under these circumstances the convulsions are not spontaneous, but occur only when an irritation is applied to a part, and are limited to the part irritated. A fact analogous to this was noticed by Kölliker (*Virchow's Archiv*, Bd. x. p. 262, Exp. IX.): in frogs whose nerves were paralyzed by woorari, the exhibition of veratria induced phenomena similar to those just noted. Evidently these facts do not prove, however, that the convulsions in the veratrized frog are not spinal, but only that there is a state of excitation of the muscles. M. Prevost furnishes the following direct proof, however, that the cord in veratria-poisoning is not affected. The hind legs of a frog

were separated from the rest of the body by a very tight ligature, so placed as not to include the lumbar nerves. Some veratria was then introduced into one of the fore legs, and of course found its way into the spinal cord and the anterior portions of the body. Under these circumstances it is evident that the convulsions produced, if spinal, would affect the whole body, but if peripheral would be confined to the anterior part of the frog. It was found that the posterior legs were never affected; that whilst irritation of them caused most violent spasms in the anterior part of the body, only the normal reflex actions occurred in those muscles not reached by the poison. If this experiment be confirmed (and I see no intrinsic reason to doubt its accuracy), to Prevost belongs the credit of having proven that veratria has no action on the motor centres of the spinal cord.

There is, however, an apparent opposition between the experiments of Prevost and those of Kölliker (*Virchow's Archiv*, Bd. x. p. 261). The latter observer noted (Exp. VI.) that when the skull of the frog was opened and a ten per cent. alcoholic solution of veratria dropped on the cord, violent general tetanic convulsions were induced; also (Exp. IV.) that when one crural artery and vein of a frog were tied and the veratria solution placed in the mouth, tetanus ensued, involving the protected limb, and continuing there after it had ceased in the other members. I see no way of reconciling these experiments of Kölliker with those of Prevost except either by supposing that the latter are incorrectly observed, or, what seems more probable, that the poison in the former reached the protected parts by diffusion, although in less quantity than it did the other members: this would also explain the continuance of tetanus in the protected limb after it had ceased elsewhere.

M. Guttman (*Reichert's Archiv für Anatomie*, 1866) is in accord with Kölliker in his experiments, for he states that, notwithstanding the artery of a limb is tied, yet spasms occur in the leg during the convulsive stage of aconite-poisoning: of course the "diffusion" theory would apply to this as well as to the experiments of Kölliker.

The only conclusion to be drawn from the evidence seems to me to be that at present it is uncertain whether veratria does or does not act upon the motor centres of the cord.

In regard to the action of the drug upon the sensitive centres of the cord, our knowledge is by no means perfect. Anaesthesia of the posterior feet was noticed in the frogs experimented upon by Prevost in the manner described; but when the circulation is cut off from the feet of a frog, loss of sensibility always ensues.

Peripheral Nervous System.—The study of the action of veratria upon the peripheral motor apparatus evidently divides itself into a study of the influence upon the muscles and the extreme nerve-endings in them and upon the nerve-trunks.

There can be no doubt that veratria finally destroys the contractile power of the muscle itself, so that it fails to respond to any irritation what-

ever, and soon, becoming stiff, exhibits the acid reaction of post-mortem rigidity. Thus far all recent observers are in accord; and I have frequently witnessed the same phenomenon. Kölliker in some of his experiments (*loc. cit.*) notes that the muscle in the early stage of veratria-poisoning responded inordinately to stimuli. The study of this phenomenon has been especially made by Bezold and Hirt (*Untersuchungen aus dem Physiologischen Laboratorium zu Würzburg, Heft i.*) and by M. Prevost.* When a muscle during the convulsive stage of veratria-poisoning is momentarily stimulated, instead of the usual momentary contraction, a prolonged tetanic spasm results and lasts some seconds: this spasm is induced by the slightest irritation. There is therefore, preceding the stage of muscular paralysis, a stage of muscular hyper-excitability. This is in accord with the experiments related when discussing whether the convulsions of veratria-poisoning were or were not spinal, and it cannot be doubted that to it in great part, if not altogether, the convulsions are due. It can scarcely be doubted that this condition is the result of an action not upon the nerve-endings, but upon the sarcolemma of the muscle. That the muscular paralysis is of similar nature would seem to be proven by the rapid changes which take place in the muscle after death, and by the fact, noted by Guttmann (*Reichert's Archiv für Anatomie*, 1866, p. 498), that whilst frogs apparently dead from nerve-poisons such as atropia, strychnia, and curari often recover themselves after a period of stupor, those poisoned with veratria never do.

When a muscle is dead, galvanization of the nerve of course elicits no response; but it is possible that a substance may be at the same time a nerve-poison and a muscle-poison. Veratria is a muscle-poison; is it also a nerve-poison? Kölliker denies this, but the experimental evidence brought forward by him amounts to almost nothing. Guttmann (*loc. cit.*) asserts that in his experiments, whenever irritation of a nerve failed to elicit a response, direct irritation of the muscle was always equally unavailing. Bezold and Hirt (*loc. cit.*) experimented, with a full knowledge of Guttmann's work, with small and with large doses, and apparently so carefully that it seems to me their results are in all probability accurate and true.

They found (*loc. cit.*, p. 90) that when a *small dose* is used there is at first a very marked increase in the irritability both of the nerve and of the muscle, so that, whether the current be applied directly to the muscle or indirectly through the nerve, contractions take place more readily than normal. After a time, both muscle and nerve lose their irritability, so that no contraction follows either the direct or the indirect stimulation. The process does not go on, however, *pari passu* in the two organs. The irritability increases sooner and is sooner lost in the nerve than in the muscle, so that there is a time when galvanic irritation of the nerve fails to induce contraction, although the muscle still retains its functional power. Moreover, the

* Quoted by Husemann.

upper or spinal end of the nerve dies first, so that at a certain stage irritation of the nerve-trunk close to its origin fails to induce contraction of the tributary muscle, although when applied lower down it elicits a response. It ought, however, to be stated that Kölliker claims that whilst a muscle placed in a weak solution of veratria soon loses its irritability, the nerve retains it under similar circumstances.

As already stated, the action of veratria upon the sensory centres is doubtful; its influence upon the peripheral sensitive nerves has not, that I am aware of, been carefully worked out, but the effects of its local application to the human skin seemingly show that it first strongly excites and then paralyzes them.

Circulation.—After death from a large dose of veratria, the heart is soft, dilated, full of blood, and incapable of responding to galvanism; *i.e.*, the heart-muscle is dead. According to Bezold and Hirt (*loc. cit.*), after a small dose there are quickening of the pulse and rise of the blood-pressure, which soon return to the normal condition; whilst immediate and persistent fall in the number of the heart-beats and in the arterial pressure follows a large dose. If the vagi be divided previous to the poisoning, a large dose produces a temporary increase in the pulse; and a stimulation of the distal end of the cut nerves by a current too slight to be felt in the unpoisoned animal retards very markedly the beat. From these facts it follows that in the uninjured animal, after poisoning by veratria, there is an inhibitory retardation of the pulse, and also an excitation of the peripheral ends of the vagi. That it is not merely the peripheral inhibitory apparatus which is affected was proved by injecting the alkaloid into the carotid,—*i.e.*, into the inhibitory centre,—when there happened an instantaneous and remarkable retardation of the heart-beat, which could only have been caused by excitation of the inhibitory centre. In a later stage of the poisoning the strongest faradaic currents applied to the pneumogastrics fail to affect the heart. It is, therefore, evident that veratria first exalts and then destroys the functional activity of the *par vagum*, as of the spinal nerves.

When the heart is separated from the nerve-centres by section of the *par vagum* and of the spinal cord, veratria produces, according to Bezold and Hirt, at first increase in the pulse and blood-pressure, secondly, lowering of both to the minimum; showing that it exerts upon the internal heart-ganglia, or upon the heart-muscle, its peculiar action of first stimulating and afterwards paralyzing functional activity.

That the poison has similar action upon the vaso-motor centres seems probable from the facts noted by Bezold and Hirt: first, that injection into the carotid after section of the pneumogastrics caused immediate rise of the blood-pressure; second, if the mesenteric arteries have been previously bared, they can be seen to contract. This excitation is followed after a time by vaso-motor paralysis and dilatation of the vessels.

In regard to the *respiration*, Bezold and Hirt conclude, from the fact that

after section of the pneumogastrics even the smallest doses of veratria cause retardation of the respiration without previous increase, that the alkaloid depresses immediately the centre of respiration in the medulla, and finally destroys it.

THERAPEUTIC ACTION.—The study of the physiological action of veratria shows that its rational therapeutic use must be limited. As a heart-sedative, it is much inferior to aconite and veratrum viride, for obvious reasons, and, although it has been used as such, it has not achieved much reputation. When exhibited in full doses it is very apt to give rise to exceedingly disagreeable secondary symptoms, and has no advantage over the medicines just named. Some years ago it was employed in *acute rheumatism*, having been recommended by Turnbull, Bardsley, Piedagnel, Trousseau, and others; but it is not so efficacious in this disease as other far less dangerous remedies by which it has been superseded. The same is true of its employment in *dropsy*, and I know of no condition which would seem to me to call for its internal use.

Bardsley originally employed it in *neuralgia*, especially when arising from cold. He used it both internally and externally. At present it is rarely employed except as a local application. My own success with it has not been very encouraging, but others of larger experience recommend that it be rubbed over the trunks and filaments of the affected nerves in *rheumatic neuralgia*.

As an external stimulant and rubefacient it is sometimes used with good effect in narcotic poisoning; also in various spinal troubles, as an irritant applied to the spine, and to the skin of the paralyzed limbs, to aid in maintaining circulation; but all these indications can, I think, be better met by other means. In regard to the dose of veratria for internal use, it should be borne in mind that one-sixteenth of a grain has produced the most alarming symptoms (Taylor, *Medical Jurisprudence*, 2d edition, London, 1873).

An ointment (*Unguentum Veratræ*, grs. xx to ʒi) is official.

ACONITI FOLIA—ACONITE LEAVES.

ACONITI RADIX—ACONITE ROOT.

The smooth, thin, bright-green leaves of the *Aconitum Napellus*,* or monkshood, a tall perennial, indigenous in Europe, and cultivated in this country for the sake of its spike of blue flowers, are three or four inches in diameter, and cut almost to the base into three to seven three-lobed, wedge-shaped divi-

* All of the species of the genus *Aconitum* are more or less poisonous, although *A. Napellus* is the only one official. For a study of the comparative strength of the various aconites, see Schrott, *Journal für Pharmacodynamik*, 1857, p. 335. He arranges them as follows, commencing with the most virulent: *A. ferox*, *A. Napellus*, with its varieties, *neomontanum*, *tauricum*, and *variabile*, *A. Cammarum*, *A. paniculatum*, *A. Anthora*. The toxic properties of the *A. Anthora* were very weak.

sions. Their taste is bitterish, acid, and after a little while benumbing, giving origin to intense tingling of the lips and mouth.

The root is from three to four inches long, very tapering, about three-quarters of an inch in diameter at the base. Its taste resembles that of the leaf. It is to be distinguished from horse-radish root, with which it has been fatally confounded, by its external brown color and its absence of odor when scraped.

In 1833 Geiger and Hesse discovered in aconite an alkaloid, *Aconitia*, which is undoubtedly the active principle of the drug. This alkaloid is now officinal. As prepared according to the directions of the U. S. Pharmacopœia, it is a yellowish-white powder. In commerce there are several varieties of it, made by different large manufacturers: the *German aconitia*, which is very impure, and, according to Husemann, is less active than the extract; *impure English aconitia*; and the so-called *English aconitia*, prepared by Morson and said to be chemically pure. The latter is a grayish powder.

Recently Duquesnel (*Comptes-Rendus*, vol. lxxiii., 1864) has obtained the aconitia in the form of colorless, rhombic, tabular crystals, soluble in alcohol, benzine, ether, and extremely so in chloroform, very slightly soluble in water, insoluble in glycerine.

The salts of aconite are soluble, and the alkaloid is precipitated by alkalies from their solution in an amorphous state. That aconitia is the only active principle of the root would seem to follow from the experiments of Hottot (*Journal de Physiologie*, 1864).

In 1857 Hübschmann announced the presence in minute quantity of a second alkaloid in the root of *Aconitum Napellus*,—*Napellina*. Schroff* (*Journal für Pharmacodynamik*, i. 3) could find no essential difference between its action and that of German aconitia. T. and H. Smith, of Edinburgh, have found a third non-poisonous alkaloid, *Aconella*, which they think to be probably identical with narcotina; and Flückiger asserts that there are four alkaloids contained in the genus *Aconitum*, namely, *Aconitia*, *Pseud-aconitia*, *Napellina*, and *Lyctonia* (*Sydenham Year Book*, 1869 and 1870).

PHYSIOLOGICAL ACTION.—When applied to a raw surface, or to the skin, aconite, or its alkaloid aconitia, acts as a local irritant and narcotic, soon producing numbness, with tingling, which may persist for a long time. When given in sufficient dose internally, it is a violent poison, acting, so far as is known, similarly upon all animals.

If the dose be large, death may be almost immediate, and, if the alkaloid be given hypodermically, may occur in less than a minute. In such cases the result is apparently due to sudden paralysis of the heart-muscle.

After moderate toxic doses, the prominent symptoms are great disturbance of the respiration, muscular weakness, vascular depression, and finally death,

* From what Schroff says about the material he used in his experiments, it is evident that he had no proof that it was genuine napellina.

with or without convulsions. As I have seen the rabbit after the injection of one-sixth or one-quarter grain of Morson's pure aconitia, the animal commences to jump vertically in a very peculiar manner, and often to squeal piteously. The jumping soon grows less and less powerful, and finally is replaced by severe convulsions, during which the animal often lies prostrate on its side. In the dog, however, the muscles have remained without a quiver during all stages of the poisoning, and it seems certain that the convulsions are an inconstant symptom, dependent upon peculiarities of the individual or species, as well as upon the amount injected.

Dilatation of the pupil very frequently occurs, if it be not indeed a constant phenomenon.

The symptoms which are induced by small therapeutic doses of aconite in man are reduction of the force and frequency of the circulation, a sense of muscular inertia and weakness, and a slight tingling in the extremities or in the lips. If the dose administered be large, all these symptoms are intensified; the muscular weakness is extreme; the tingling is felt all over the body; the pulse is feeble, and reduced to thirty or forty per minute; the respirations are diminished; giddiness and disordered vision may be manifested, especially when the erect posture is assumed. After three or four hours these symptoms gradually subside.

When a poisonous dose has been ingested, the first thing noticed in most cases is a burning or tingling in the throat or in the extremities, soon spreading over the whole body. The pulse rapidly falls in frequency, and in a very little time becomes exceedingly weak, intermittent, irregular, and finally imperceptible; the muscular strength is greatly reduced, and sometimes almost entirely gone; the respirations are shallow, feeble, irregular, and infrequent; the general sensibility is very much benumbed, so that marked anæsthesia of the surface is present; the skin is bedewed with a cold sweat; the countenance is anxious, sunken, livid, and the eyes are often protruded, or are even spoken of as glaring; the pupil is generally dilated, but when there are no convulsions may be contracted; gastric burning is sometimes complained of, and severe vomiting may be present, but the stomach is not rarely retentive. The intellect generally remains unaffected until very near the close, sometimes to the very moment of death.* In the collapse of the latter stages of aconite-poisoning the special senses may be lost, especially the sight. The voice is very generally extinguished. Convulsions occur in some cases, not in others; and certainly in some instances, if not always, the patient is unconscious during their continuance. Death may occur suddenly, especially *directly after some exertion* on the part of the patient, from syncope.

The symptoms which aconite produces in man and in the lower animals are so entirely identical that the conclusions arrived at in regard to the latter may be accepted without reserve as applicable to the former. The action of aconite

* Pereira, however, states that in some recorded cases stupor has occurred.

upon the circulation is very decided. According to Dr. Achscharumow (*Reichert's Archiv*, 1866, p. 255), in the frog a moderate toxic dose of aconitia produces at first a reduction in the number of the heart's pulsations, then an increase in the rapidity of its action, with very evident loss of power, and finally irregular systolic movements, with very long intervening pauses ending in diastolic arrest. Dr. Rudolf Böhm and L. Wartmann (*Schmidt's Jahrbücher*, 1872, Bd. cliv. p. 265) have substantially confirmed these observations.

In the higher animals the exhibition of aconite in sufficient doses yields similar results. In the dog and cat (Böhm and Wartmann, and my own experiments) there is a steady sinking of the arterial pressure; in the rabbit, according to Böhm and Wartmann, this fall is preceded by a brief rise. The rate of the heart's pulsations also undergoes reduction, and there is finally diastolic arrest in these and other mammals.

The method by which the aconite influences the heart is not certainly settled. According to the experiments both of Böhm and Wartmann, it produces a gradual paralysis of the peripheral vagi, a constant increase of the intensity of a galvanic stimulation of the pneumogastric nerves being required to influence the heart as the poisoning deepens, until finally the vagi entirely refuse to transmit any inhibitory impulse.

In a single experiment, Achscharumow (p. 272) found that after section of the vagi in the early stage of aconite-poisoning there was an immediate rise both in the number of the cardiac pulsations and in the arterial pressure. From these data he argues that the slowing of the pulse during the early stage of aconite-poisoning is due to stimulation of the inhibitory centres in the medulla oblongata. Böhm and Wartmann (*loc. cit.*, p. 266) repudiate this conclusion, because, according to their experience, the phenomena of aconite-poisoning occur in the usual manner after section of the vagi, or in atropized animals. It is evident that there is no necessary contradiction in the asserted facts of these observers, as it is possible that the slowing of the pulse may be due to two immediate causes, one having its seat in the medulla oblongata, the other in the heart. Be this as it may, however,—and the point requires further investigation,—it is very certain that aconitia influences directly the heart, or its contained ganglia, for Achscharumow (*loc. cit.*, p. 262) has found that it acts upon the frog's heart removed from the body, and Liégeois and Hottot (*Journal de Physiologie*, p. 520, 1861) have observed the ordinary cardiac phenomena of aconite-poisoning produced by the alkaloid placed directly upon the viscus. Böhm and Wartmann have also noted that in aconite-poisoning the force of the individual beat is lessened. After death the cardiac muscle fails entirely to respond to galvanic irritation, its contractility being lost.

Our knowledge of the action of aconitia upon the vaso-motor nerves is by no means complete. Achscharumow, and more recently Dr. F. B. Nunneley (*Proceedings of the Royal Society*, p. 46, 1870), studied with the microscope the influence of injections of aconitia upon the vessels of the frog's web,

but were unable to detect any alteration of their calibre. The former observer also found that after division of the sympathetic in the neck, galvanization of the peripheral end produced the usual phenomena, even in the most advanced stages of aconite-poisoning. These facts indicate very strongly that aconite does not affect the vaso-motor nerves, and this indication is confirmed by the experiments of Böhm and Wartmann, who found that when in aconite-poisoning a galvanic current was applied to the vaso-motor centres in the medulla, an immediate rise of arterial pressure took place. As stimulation of a sensitive nerve produced at such time no rise of arterial pressure, the conclusion would appear to be inevitable that aconitia, whilst not affecting the afferent vaso-motor nerves or the vaso-motor centres, destroys the conducting power either of the afferent nerves or of the cord, so that in an animal under its influence no impulse can be transmitted from the periphery to the vaso-motor centres in the medulla.

Quite diverse conclusions in regard to the action of aconitia upon the nervous system have been arrived at by different investigators. Achscharumow concludes that the spinal cord is not affected, and that the paralysis and loss of reflex activity induced depend upon the destruction of the conducting power of the peripheral nerve, because he has found that when a frog is poisoned, after the abdominal aorta has been tied, reflex and voluntary activity is preserved in the hind legs long after it has been lost in the anterior portion of the body; and at the same time, whilst the brachial nerves, as tested by galvanic stimulation, have lost their power of transmitting impulses, the protected ischiadic nerves have preserved their functional ability. On the other hand, Böhm and Wartmann declare that the peripheral nerves are not materially affected. From numerous experiments on frogs,—with and without the previous separation of the brain,—and also with section of the spinal cord below Setschenow's reflex inhibitory centre (whereupon in the aconitized frog no increase of reflex activity came on), Böhm and Wartmann conclude that the aconitia first depresses the reflex activity of the sensitive spinal centres and afterwards that of the motor spinal centres, until the cord is completely paralyzed.

A very complete and beautiful investigation of the action of aconitia upon the spinal cord has been made by Dr. Liégeois and M. Hottot (*loc. cit.*, p. 533), who apparently have succeeded in elucidating the matter. According to these observers, in aconite-poisoning loss of sensibility occurs in the frog's legs simultaneously with or even before the disturbances of respiration, and long before the power of voluntary motion is lost, and even when the reflex activity is intact. This sensory paralysis, according to the experiments of the French investigators just quoted, first appears in the hind legs of a frog poisoned with aconitia, and has not its primary seat either in the peripheral nerves or in the spinal cord, for it was found that tying the aorta close to its abdominal bifurcation, so as to prevent access of the blood—*i.e.*, of the poison—to the posterior nerves, did not affect the development of the anæ-

thesia; further, that closing the artery nearer its origin in such a way as to shut off the circulation to the cord and spinal nerves, but to allow the passage of the blood to the cerebrum, did not cause sensory paralysis to come on more slowly than normal when the animal was poisoned with aconite. If the experiments of Liégeois and Hottot be confirmed, the demonstration will be perfect that aconitia paralyzes a sensory perceptive centre above the spinal cord.

Of course it is possible for the peripheral ends of the sensory nerves to be paralyzed either at the same time that the perceptive centre is, or afterwards; and of course, the centre being paralyzed, it becomes very difficult to determine whether the periphery is or is not affected. Liégeois and Hottot assert that this paralysis of the centre occurs before any serious implication of the peripheric nerves, because after aconitic anaesthesia had been produced strychnia was able to induce tetanus; afterwards, however, the extreme peripheric nerves became affected, so that irritation of the skin in the doubly-poisoned frog would not provoke convulsions, even at a time when irritation of the trunk of a nerve would produce general reflex motor disturbance. At last galvanization of the nerve-trunk itself failed to induce response. From these facts Liégeois and Hottot deduce—very logically, I think—the conclusion that aconite induces anaesthesia by paralyzing, first, the perceptive centres; secondly, the peripheral extremities of the nerves; thirdly, the nerve-trunks themselves. The observers alluded to also confirmed this conclusion by other experiments than those already noticed. They found that although aconitia applied directly to a nerve-trunk paralyzes its sensibility, yet when the veins of a frog's leg are tied and the alkaloid injected into the artery and allowed to permeate the tissues of the leg, the skin loses its sensibility long before the nerve is affected.

In regard to motion, Liégeois and Hottot found that in a certain stage of aconite-poisoning the frog lies with his limbs extended, relaxed, and perfectly paralyzed, and yet is capable of executing vigorous voluntary movements and evinces nearly normal reflex activity. They attribute this condition of apparent but not real motor paralysis to loss of sensibility from paralysis of the perceptive centre, as the unpoisoned frog evinces the same phenomena after division of all the posterior spinal roots. After a time the reflex activity is also lost, the power of voluntary movement remaining. Liégeois and Hottot believe that this loss of reflex activity is spinal; but in their experiments upon the conjoint action of aconite and strychnia it was found that at a certain stage, when no amount of irritation of a nerve would induce convulsions, a slight direct irritation of the cord would cause violent strychnic spasms. This would seem to prove that at least the earliest abolition of the reflex activity was due to paralysis of the afferent nerve-fibres.

After a time, in aconite-poisoning, even voluntary movements disappear; but, according to our observers, this is not due to paralysis of the motor nerve-trunk, but is central, since irritation of the nerves causes contractions in the tributary muscles. The motor nerve-trunks are, however, affected by

aconitia to some extent, since Liégeois and Hottot have found that locally applied it destroys their conducting powers, and after death in poisoning by it the motor nerves lose their sensitiveness much earlier than normal.

The action of aconite upon the respiration is very marked, and, as arrest occurs in the frog before the motor nerves are affected by the poison, Liégeois and Hottot believe that the disturbance is centric. Böhm and Wartmann arrive at the same conclusion, basing their opinion upon the fact that previous section of the vagi does not influence the action of the poison upon the respiration. I think there can be no doubt that aconite is a direct depressant and paralyzant of the respiratory centres.

Close studies of the action of aconite upon the temperature in health are wanting. Achscharumow found in fatal poisoning a fall of about 3° C.

MM. Gréhaut and Duquesnel (*L'Union Pharmaceutique*, Aug. 1871) have communicated to the French Academy some experiments upon frogs with crystallized aconitia, whose results are so strikingly different from those of other experimenters as to indicate the existence of some fallacy: possibly the alkaloid used by them was not the same as the amorphous aconitia. They found in the frog, after small doses ($\frac{1}{20}$ milligramme) of their alkaloid, that the heart continued to beat steadily and regularly after all power of spontaneous or reflex movement had been lost, that sensation was preserved as long as any power of motion existed, and that the motor nerve-trunks were paralyzed. After large doses (one milligramme) they observed sudden arrest of the heart's action.

THERAPEUTICS.—Our knowledge of the physiological action of aconite, imperfect though it may be, is sufficient to show that there are only two or three indications to meet which it may be used.

The first of these is to *lower arterial action*, and often, with it, excess of temperature. For this purpose aconite is almost invaluable. I have never used it in those cases, such as *pneumonia*, in which a sudden and very powerful effect is desired, simply because veratrum viride seemed to me safer, more readily controlled, and equally effective. Aconite may, however, be used with very good results in these cases, and especially in such diseases as *peritonitis*, in which it is very important to avoid vomiting. My own experience with it has been in *fevers* of a sthenic type not dependent upon so deep-seated a cause (as an example may be mentioned the febrile movements of severe, acute *muscular rheumatism*), and in the *ephemera* or *irritative fevers* of childhood: in such cases its influence for good is often very decided. In the early stages of *scarlet fever* and other *exanthemata*, when not decidedly adynamic in type, it is very useful. In the reflex fever which sometimes follows the passage of the catheter or bougie (the so-called *urethral fever*) it is very efficient.*

* The following formula affords an excellent combination: ℞ Tr. aconit. rad., gtt. i; Sp. ether. nitric., fʒii; Mist. potass. citrat., q. s. ad fʒi. S.—Dessertspoonful every two hours for a child three years old.

In some cases of *hypertrophy of the heart*, when the valves are perfect, or when, the valves being diseased, the *hypertrophy* is greater than is necessary, aconite is of use to control cardiac excitement. When, however, there is dilatation of the heart or any degeneration of the heart-muscle, it is an exceedingly dangerous remedy, and is also at all times to be avoided if the *hypertrophy* be not excessive.

A second indication, which aconite might be used to fulfil, is to *alloy spasm*. As, however, its influence upon the motor centres and nerves is much less than upon the sensitive centres and nerves and upon the heart, the indication is better met by other remedies.

A third indication, which it would seem from its known physiological action that aconite should meet, is to *relieve over-excitation of the sensitive nerves*. Clinical experience has confirmed this. As long ago as 1834, Dr. Turnbull (*On the Preparations and Medical Employment of Aconitina by the Endermic Method*, London, 1834—*On the Medical Properties of the Natural Order Ranunculaceæ*, London, 1835) called attention to the use of the alkaloid in *neuralgia*; and his estimate of its value has been confirmed by Dr. A. Fleming (*An Inquiry into the Physiological and Medicinal Properties of the Aconitum Napellus*, Edinburgh, 1845) and by other observers.

In cases of *rheumatic neuralgia* dependent upon an acute exposure to cold, and attended with more or less febrile disturbance, in combination with other suitable remedies aconite is often of great service. In *chronic neuralgia*, associated as it always is with a lowered systemic tone, the remedy is less efficient; yet in some cases it seems to give relief. Owing to its very marked local benumbing influence, applied to the painful part it is sometimes very useful. In my own experience, this local use of it has, however, very seldom been effective when, as in *migraine*, the pain is of centric rather than of peripheral origin.

Given in full doses in the reflex *vomiting* of *pregnancy*, aconite is often advantageous, acting probably by benumbing the sensory reflex centres, or possibly the afferent peripheral nerves. I have noticed that relief lasts only so long as decided constitutional effects from the drug are apparent.

TOXICOLOGY.—Aconite is an exceedingly powerful poison, one-twelfth of a grain of the crystallized alkaloid being, according to Duquesnel, sufficient to kill a rabbit in a short time.

The peculiar tingling is the only diagnostic symptom, but it is very characteristic. The first indication for treatment is to evacuate the stomach and wash it well out with the stomach-pump. Alcoholic stimulants should be freely administered, hot and concentrated, and the injection of ammonia into the veins may be practiced. Great care should be used to keep the patient absolutely quiet, upon the back, with the feet a little higher than the head.

Recently attention has been drawn to the employment of digitalis in aconite-poisoning. It was discovered by Dr. J. Milner Fothergill (*Digitalis*, London, 1871, p. 6) that when digitalis is administered to frogs under the

influence of aconite, the heart is visibly relieved from the depression produced by the first poison. Even when all cardiac action had apparently ceased, digitalis had power to recall the systolic movements, until finally a return to the normal state was brought about. In a case reported in the *British Medical Journal* of December 11, 1872, recovery occurred after the ingestion of an ounce of Fleming's tincture of the root. The patient, when first seen, was apparently dying. Twenty minims of the tincture of digitalis were hypodermically injected, and after twenty minutes, the man having revived sufficiently to swallow, a fluidrachm of the tincture with ammonia and brandy was given him, and was repeated twice within the hour. The evidence, though as yet scanty, seems to me strongly in favor of the use of digitalis in aconite-poisoning.

ADMINISTRATION.—Aconite is never used in substance, but in one of the following preparations:

Tinctura Aconiti Radicis—*Tincture of Aconite Root* (℥vi-Oj). Dose, one to five drops, repeated every one to three hours *pro re nata*, its effects being always watched. Fleming's tincture is a stronger preparation (℥xss-Oj).

The tincture of aconite is very frequently added to stimulating and anodyne liniments.

Extractum Aconiti—*Extract of Aconite*. Dose, one-quarter to three-quarters of a grain.

Aconitia.—On account of its intense activity, aconitia should not be given internally; and even its external use requires care. The ointment may be made of the strength of from two to ten grains to the drachm.

ACIDUM HYDROCYANICUM—HYDROCYANIC ACID.

Pure hydrocyanic acid is a colorless, transparent, volatile, inflammable liquid, giving rise to giddiness and headache when smelled, and having, it is said, a burning bitter taste. So poisonous is it that when inhaled it causes death, and it must be handled with the greatest caution: smelling and tasting of it are excessively dangerous proceedings. It is indeed an imperative rule that no one should experiment with anhydrous prussic acid alone, or under any circumstances in summer, or in a warm room, or in an apartment whose open windows and doors do not admit of a free draft of air. The chemist Scheele, the discoverer of prussic acid, is believed to have been killed by the inhalation of the fumes of this material, whose poisonous properties were first pointed out by the Berlin apothecary Schrader in 1803. The anhydrous acid is soluble in water and alcohol, but is never kept in the shops, and is not officinal.

Hydrocyanic acid of common medical parlance is the officinal *Dilute Hydrocyanic Acid* (ACIDUM HYDROCYANICUM DILUTUM), a colorless, watery solution, containing two per cent. of the anhydrous acid. Its odor and taste are the familiar ones of peach-kernels and bitter almonds; its reaction is faintly acid. According to the directions of the U. S. Pharmacopœia, it is prepared

by distilling a mixture of ferrocyanide of potassium, sulphuric acid, and water. or by precipitating cyanide of silver from its watery solution with muriatic acid. With solution of nitrate of silver added in slight excess, one hundred grains of it produce a white precipitate, which, when washed with water until the washings are tasteless, and dried at a temperature not exceeding 212° , weighs ten grains, and is wholly soluble in boiling nitric acid.

The precipitate in this case is the cyanide of silver, and the amount afforded shows that the liquid contains the official percentage of anhydrous prussic acid.

As hydrocyanic acid has a great tendency to undergo spontaneous decomposition, especially under the influence of light, it should be kept in well-stopped, dark-colored bottles.

PHYSIOLOGICAL ACTION.—In warm-blooded animals, poisoning by hydrocyanic acid divides itself naturally into the acute and the subacute; death occurring in the first in at furthest ten minutes, in the second not at all, or else only after the lapse of a longer time than that noted. After a full dose of the strong anhydrous acid, the animal gasps once or twice, and then instantly falls in a tetanic or clonic convulsion, or else drops motionless and powerless upon its side. In either case, at once the signs of asphyxia manifest themselves, and grow more and more intense, until they end in total arrest of respiration. The heart beats irregularly, often at first slowly and strongly, with intervals of suspension of movement, but always becoming weaker and more rapid in its action, until, after the breathing has ceased, its efforts gradually die away. If the dose has been enormous, the heart and lungs may stop acting at once; otherwise the cardiac pulsations may continue some minutes after the arrest of respiration. Ordinarily, three distinct stages are apparent: a first, very brief one, of difficult respiration, slow cardiac action, and disturbed cerebration; a second, convulsive stage, with dilated pupils, violent convulsions, unconsciousness, loud cries, vomiting, often spasmodic urination and defecation, erections, etc.; the third period, of asphyxia, collapse, and paralysis, sometimes interrupted by partial or even general spasms.

The slow form of the poisoning follows the exhibition of the poison in an amount just sufficient to kill. After the ingestion of such a dose, no phenomena are offered for some seconds; then the breathing becomes labored, and the pulse slow and full. The animal perhaps cries out, and muscular tremblings invade the whole body, to give place, in a very short time, to clonic and tonic convulsions, which continue at intervals until the third stage, that of collapse, is developed. The convulsions are less violent and less frequent than those of the acute poisoning; all the symptoms noted as occurring during the second stage of rapid cases are present in the corresponding period of the subacute poisoning, although less violent and less intense in their manifestations. When the third stage is developed, the anaesthesia is marked, affecting first the hind legs, but finally spreading to all parts of the body, and even being complete in the widely-dilated pupil. Death finally results from

failure of respiration. Recovery may occur even after the conjunctiva has lost its sensibility; the return to life by a subsidence of the symptoms is usually rapid, so that generally in from one-half to three-quarters of an hour the animal will be eating, as though nothing had happened. Coullon, however, noted persistence of paralysis, in some cases, for days.

In man, prussic acid produces results closely parallel with those which it causes in the lower animals. The symptoms come on suddenly. In a moment or two the individual falls to the ground insensible and convulsed, the respirations arrested or occurring at long intervals, the eyes salient, the pupil dilated, the mouth covered with bloody froth. If the dose be sufficiently large, death may occur in three or four minutes; if less have been taken, deep insensibility, tetanic or clonic convulsions, dilated pupils, a bloated countenance, cyanosed surface, set jaws, and irregular respiration, constitute the chief symptoms. The breathing is mostly convulsive, with deep, forcible expirations, but in some cases it has been stertorous. Death results from asphyxia. After small toxic but not lethal doses of prussic acid, giddiness, lightness of the head, nausea, a quick pulse, and muscular weakness, are the chief symptoms.

As early as 1814, Dr. F. B. Vietz (*Medicin. Jahrb. d. k. k. Oesterreich. Staates*, Bd. ii., 1814) called attention to the change of color that occurs in the venous blood of animals poisoned with prussic acid; and his observations have been confirmed by E. L. Schubarth (*Horn's Archiv f. Med. Erfahrung*, Berlin, 1824), by J. F. Sobernheim (*Handbuch der prakt. Toxicologie*, Berlin, 1838), and by Dr. Coze (*Gazette Médicale de Paris*, 1849).

In his *Leçons sur les Substances toxiques*, p. 193 (Paris, 1857), Claude Bernard reaffirms the occurrence of these changes, and further states that if the animal die suddenly the blood in the veins and right heart is found of a bright arterial hue at the post-mortem. Notwithstanding all this testimony, J. R. Bischoff (*Ueber Vergiftungen nebst einigen Versuchen an Thieren, welche mit Blausäure, Cyankalium und Arsenik angestellt wurden*, Wien, 1844) and numerous other observers have found that after death from prussic acid, either in man or other mammals, nothing but dark venous blood exists in the body. Of the correctness of this observation there can be no doubt.

Prof. W. Preyer (*Die Blausäure*. Bonn, 1870) has afforded by his experiments an explanation of these apparently contradictory facts. He found that directly after the exhibition of prussic acid to a mammal the blood becomes, even in the veins and in the right heart, of a bright arterial hue, but that after a time this color darkens into the blue of venous blood, and finally, even in the arteries and in the left ventricle, only blood of such character is to be found.

Dr. Carl Gaethgens (Hoppe-Seyler's *Medicinisch-chemische Untersuchungen*, Berlin, 1866, p. 324) has, in a number of experiments, confirmed this, so it must be accepted as a fact. When an animal dies suddenly from cardiac paralysis, during the first stage of poisoning, this excessive arterialization may be found after death, as mentioned by Claude Bernard; and, as Preyer

first noticed, in cold-blooded animals the bright color persists for many hours.

By spectroscopic examination Prof. Preyer (*loc. cit.*, p. 95) found that the dark blood of prussic acid poisoning is absolutely or almost free from oxygen, showing only the absorption bands of deoxidized hæmoglobin, whilst Gaethgens (*loc. cit.*, p. 328) has discovered that the red venous blood of the first stage of the poisoning shows very clearly the absorption bands of oxyhæmoglobin. The first question which arises at this juncture is as to the causes of these changes of the blood, its primary excessive arterialization, its secondary excessive carbonization. Prof. Hoppe-Seyler affirms (*Medicinisch-chemische Untersuchungen*, p. 140, Berlin, 1867) that the appearance of red blood in the veins is because the red blood-corpuscles have been so acted upon by the poison as to have lost their ability of yielding up their oxygen in the capillaries. Dr. Carl Gaethgens (*Ibid.*, p. 325) has by an elaborate series of experiments shown that in the first stage of prussic acid poisoning much less than the normal amount both of carbonic acid and of exhaled oxygen is eliminated. The lessened exhalation of oxygen probably depends simply upon a lessened inhalation of oxygen, owing to the disordered respiration. That the lessened excretion of carbonic acid is not due to the same cause, however, is shown by the fact, determined by Gaethgens (*loc. cit.*, p. 347), that the percentage of the acid in the expired air is less than normal, whilst that of oxygen is greater than normal. If the lessened excretion of carbonic acid were produced by the entrance into the lungs of an insufficient amount of air for the wants of the system, it is evident that the expired air would contain more than its normal proportion of carbonic acid and less than its normal amount of oxygen. The observed phenomena seem to me to prove that during the first stage of prussic acid poisoning oxidation is arrested. They do not demonstrate, however, that the arrest is due to a direct action of the poison upon the blood-corpuscles. The probabilities of such occurrence are rendered very slight by the investigations of Gaethgens himself, for he found that when the experiments were prolonged from seventeen to forty-four minutes, much more than the normal amount of carbonic acid was exhaled, a fact in accord with the excessive carbonization of the blood known to take place in protracted hydrocyanic acid poisoning. As it seems incredible that a substance should one minute paralyze the ozonizing power of the blood-corpuscles and the next minute increase it, it is very improbable that the super-arterialization of the blood in the first stage of prussic acid poisoning is due to a direct action of the poison upon the red disk.

Preyer has proven (*loc. cit.*, p. 85) that when to blood at the temperature of the body hydrocyanic acid is added, the spectrum after a time is altered and new absorption bands appear. These bands are due to the formation of a new compound by the union of the hæmoglobin and the hydrocyanic acid. This substance, *cyanohæmoglobin*, was first discovered by Hoppe-Seyler (*Virchow's Archiv*, Bd. xxxviii. p. 475), and has no ozonizing power whatever;

to its formation, no doubt, is owing the loss of ozonizing power by blood to which hydrocyanic acid is added outside of the body, a phenomenon pointed out by Schönbein* (*Schmidt's Jahrbücher*, Bd. cxl., 1868, p. 161), and indicated even earlier by the researches of Prof. Harley (*London Philosophical Transactions*, 1865, p. 706). The latter observer found that the blood taken out of the veins of a subject forty-eight hours after death from prussic acid poisoning, and thoroughly arterialized by shaking with air, and then allowed to stand, yielded gas containing 19.56 parts of oxygen, 80.44 parts of nitrogen, and 0.00 parts of carbonic acid.†

These facts at first sight seem to prove the theory of Hoppe-Seyler, to which indeed they no doubt gave origin. Preyer has shown (*loc. cit.*, p. 95), however, that the dark blood of prussic acid poisoning has not lost its power of oxidation, for on being shaken with the air it assumes the red arterial hue; and Drs. Lecorché and Meuriot (*Archives Générales*, t. xi., 6e série, p. 539) have determined that artificial respiration will produce the same result in the poisoned animal. Moreover, the spectroscope shows plainly that the hæmoglobin exists in the blood either in its pure state (Preyer, *loc. cit.*, p. 95), or else as oxyhæmoglobin (W. Laschkewitsch, *Reichert's Archiv*, p. 652, 1868), and that no cyanohæmoglobin is present. The reaction between hydrocyanic acid and hæmoglobin is one requiring some time, and evidently does not occur in poisoning. On the whole, therefore, I think that the chemical evidence not only affords no proof, but does not even indicate the truth, of the theory that prussic acid acts in the body directly upon the red blood-corpuscles.

Preyer (*loc. cit.*, zweiter Theil, p. 88) has shown that the excessive oxygenation and the subsequent excessive carbonization of the blood are not peculiar to hydrocyanic acid poisoning, but are equally present after the exhibition of sulphuretted hydrogen, and even after mechanical closure of the mouth and nose. It is possible that an increased arterial pressure, an increased rapidity of circulation, may cause the blood to pass too quickly through the capillaries to allow time for the usual changes; but this has not been

* In the same memoir Schönbein calls attention to the fact that prussic acid destroys also the ozonizing power of living vegetables, such as roots, fungi, etc.

† According to Dr. E. Ray Lankester (*Pflüger's Archiv*, 1869, p. 492), when blood is shaken with cyanogen gas, and allowed to stand for two or three hours, the spectrum changes are exactly the same as after similar treatment of blood with CO. The compound of cyanogen and hæmatin (Cy,Hb) offers not only the identical spectrum of CO,Hb, but, like the latter, is unaffected by reducing agents. After the blood stands awhile, according to Dr. E. Ray Lankester, the spectrum of hydrocyanic acid (H,CN) becomes visible in it, and the Cy,Hb undergoes conversion into the cyanohæmoglobin (Cy,Hb) of Hoppe-Seyler.

Any one desirous of investigating this subject more deeply than can be done in a work like the present should consult especially the papers by Hoppe-Seyler, *Virchow's Archiv*, Bd. xxxviii., and scattered through the *Medicinisch-chemische Untersuchungen*; by Harley, *London Philosophical Transactions*, 1865, p. 706; and by Preyer, *Pflüger's Archiv*, 1868, p. 395.

proven, and at present it must be acknowledged that we are ignorant as to the immediate cause of the blood-changes in these cases.

It is possible, although scarcely probable, that the changes in the color of the blood are due to alteration in the form of the corpuscles. According to Ernst Geinitz (*Pflüger's Archiv für d. gesamte Physiologie*, Bd. iii., 1870, p. 46), outside of the body prussic acid produces in the blood-corpuscles of the frog, first, a shortening of the long and a lengthening of the short diameter, and consequently a rounded form, then granulations, and, finally, a solution and setting free of the nucleus. In frogs poisoned with prussic acid a rounded form of the corpuscles was commonly exhibited, and sometimes granulations were present. M. Geinitz also found that the red disks of mammalian blood, exposed to the vapor of hydrocyanic acid in the moist chamber of Stricker, become first somewhat asymmetrical, then mulberry-shaped, and finally undergo molecular destruction. In poisoning of mammals, according to the same investigator, the granular blood-corpuscles are commonly met with. Preyer (*loc. cit.*, zweiter Theil, p. 91) confirms the observation of Geinitz so far as the action of the poison upon drawn blood is concerned, but both he and Hünefeld (*Der Chemismus in der thierischen Organisation*, Leipsic, 1840) assert that immediately after death from prussic acid the corpuscles offer their usual characters.

Whatever may be the cause of the changes in the blood, the experiments of Lewisson (*Reichert's Archiv*, 1870, p. 352) would appear to prove that the action of the poison on the nervous system is a direct one, and not due to these changes in the vital fluid, for the observer mentioned found that prussic acid acted upon the bloodless "salt frog" as upon the normal batrachian.

The action of hydrocyanic acid upon the *heart* varies according to the dose. In sufficient amount and concentration, it produces instantaneous diastolic arrest, which is either permanent or re-occurs after a few slow feeble beats (Preyer, *loc. cit.*, p. 52, and Drs. Lecorché and Meuriot, *Archives Générales*, t. xi., 6e série, p. 543). As early as 1826, Krimer found that prussic acid placed directly upon the heart of the frog produces arrest of its beat and loss of its muscular irritability. Preyer has confirmed this, and it would seem to be proven that the cardiac arrest spoken of above is due to a direct action upon the heart-muscle or its contained ganglia.

The French observers just mentioned affirm, however, that after cardiac death from prussic acid the heart responds to galvanism. I cannot but think that further investigation will show that this response is very feeble; otherwise the action of the poison on the heart seems at present inexplicable.

The cardiac results of the exhibition of small non-toxic doses are, according to Preyer, simply slowing of the heart's action.

Preyer and Laschkewitsch agree as to the action of large, but not enormous, doses. At first there is a sudden prolonged diastolic arrest of the heart, followed by an augmentation in the rapidity of the cardiac action, and after

this a diminution of the rate,—to the normal number in cases of recovery, to cardiac stand-still in cases of death. As both Laschkewitsch (*Reichert's Archiv für Anatomie*, 1868, p. 653) and Preyer* have found that after section of the vagi this primary diastolic arrest of the heart does not occur, it seems an inevitable conclusion that prussic acid first stimulates and then paralyzes the cardiac inhibitory apparatus, most probably in its centres.

According to Preyer (*loc. cit.*, pp. 17, 18, 19), during all three stages of hydrocyanic acid poisoning the respirations are lessened in frequency, and during the latter moments of life the efforts at breathing are very distant, and finally cease before the arrest of cardiac movements. The observer just mentioned found that, after division of the vagi, normally lethal doses did not kill, and that when death was brought about by the exhibition of larger doses it was by cardiac arrest. From this he deduces the conclusion that the prime respiratory action of the poison is upon the peripheral ends of the vagi. Dr. Preyer's experiments have been at least partially confirmed by Drs. Lecorché and Meuriot (*loc. cit.*, p. 538); but, even if further experimentation should completely establish their truth, the conclusion which he arrives at ought to be considered only a probable one, since the physiologist has not sufficiently studied the relations of the pneumogastrics to respiration to enable us to arrive at settled conclusions. Moreover, the experiments of Prof. Joseph Jones (*New York Medical Record*, vol. ii. p. 459) indicate very strongly that prussic acid has some direct action on the centres of respiration in the medulla oblongata, for he found that whilst to kill an alligator by the administration of prussic acid required a considerable length of time, its application to the medulla produced within one minute a most powerful expiration, ending in permanent contraction of the muscles of respiration and collapse of the lung.

Dr. Kölliker (*Virchow's Archiv*, Bd. x. p. 272) has found that in frogs dead of prussic acid poisoning both the nerve-trunks and the muscles are unexcitable, or that the muscles respond very feebly to direct stimulation. This is in accord with the experiments of Stannius (*Archiv für Anatomie*, 1858, p. 95), who found that when strychnia and prussic acid were given together, the convulsions normally produced by the former poison were altogether absent, or present only in a slight degree. In order to determine whether the nerves are or are not primarily affected, Kölliker experimented by tying the vessels of the thigh, then dividing just below this point all the tissues except the nerve, and administering prussic acid by the mouth. In a number of such experiments he found that always the nerve and muscles below the point of section retained their irritability, but that if the galvanic or other stimuli were applied to the nerve higher up, they failed to elicit any response from the unpoisoned tributary muscles,—positive proof that the nerve-

* Preyer (*loc. cit.*, p. 93) has also noted the same absence in curarized animals poisoned by hydrocyanic acid.

trunks are paralyzed by a direct action of the drug. This is seemingly opposed to the experiments of Stannius (*loc. cit.*), who concluded that prussic acid applied locally to the nerves has no effect upon them. Stannius, however, compared the results of soaking a nerve in water and in a weak solution (three to four per cent.) of prussic acid, and water thus employed is toxic to the nerve-trunks. Kölliker used, in repeating the experiments of Stannius, neutral solutions of phosphate of soda, one containing, the other free from, hydrocyanic acid, and found that the nerves in the poisoned liquid died much sooner than did those in the non-poisoned solution.

Whilst it seems scarcely doubtful that prussic acid does act upon the nerve-trunks, the experiments of Kölliker are in agreement with those of Stannius, that the muscle dies very much more quickly in the solution of the acid than does the nerve, losing its excitability in from seven to eight minutes. This rapid destruction of muscular irritability by the local application of prussic acid was, I believe, first noted by Coullon in 1819.

It is most probable that when given internally prussic acid acts almost as rapidly upon the nerve-trunks as upon the muscles, since Kölliker noted that in some cases galvanization of the nerve was incapable of causing contractions in the tributary muscles, although the latter responded feebly to direct stimulation. This fact has been experimentally corroborated by Funke (*Berichte über die Verhandlungen d. k. sachs. Gesellschaft d. Wissenschaften zu Leipzig*, Bd. xi., 1859, p. 28).

Upon the peripheral sensitive nerves prussic acid probably, if in sufficient concentration, acts as a paralyzant; at least Kölliker (*loc. cit.*, p. 282) found that if the leg of a strychnized frog, whose heart had been cut out to prevent absorption, was put in a four per cent. solution of prussic acid, in a very short time irritation of the immersed skin ceased to produce convulsions.

From the slowness with which, in Kölliker's experiments, the nerve-trunks were affected in frogs poisoned by hydrocyanic acid, it seems probable that he is correct in his conclusion that in these batrachians the poison first paralyzes the brain, and then the reflex centres of the spinal cord, and afterwards the motor nerves. I have not met with any experimental evidence, however, in regard to the order in which prussic acid affects the nervous system. According to Kiedrowski,* as quoted by Preyer, in frogs it first paralyzes the gray, then the white substance of the brain, and the early disappearance of reflex movements is not due to spinal palsy, but to destruction of the functional power of the peripheral afferent nerves.

Preyer also states that the conclusions of Kiedrowski rested upon the following experimentally proven fact, which, if accurate, seemingly renders them

* I have, unfortunately, been unable to obtain access to the original paper of E. de Kiedrowski. Even Preyer appears to know it only in abstract. According to him, it was published in 1858, at Breslau, as a dissertation, under the following title: *De quibusdam experimentis quibus quantam vim habeat acidum hydrocyanicum in nervorum systema cerebro-spinale atque in musculos systematis vertebralis probatur.*

logically inevitable. When a frog is poisoned with prussic acid, and afterwards with strychnia in properly-proportioned doses, there is a stage at which slight irritation of the afferent nerve-roots causes violent general tetanic spasms, although the most intense peripheral irritation fails to elicit response.

It is a question of interest to decide as to the cause of the convulsions in poisoning by hydrocyanic acid. I have found that they do not occur after section of the cord in parts below the point of section, and that they are therefore cerebral in origin: for reasons detailed elsewhere (see p. 129), it is very probable that they are due to disturbed cerebral circulation, and this probability is confirmed by an experiment of Laschkewitsch (*Reichert's Archiv für Anatomie*, 1868), who opened the thorax of a rabbit so as to expose the heart, maintained artificial respiration, and administered prussic acid; directly after arrest of the heart had commenced the convulsion came on. The earlier observation of Coze (*Comptes-Rendus*, t. xxviii., 1849, p. 780) is also to the same effect, as he states that the convulsions did not occur until directly after the arrest of the circulation. In frogs poisoned with hydrocyanic acid, convulsions do not take place.

Preyer states that after section of the vagi convulsions do not generally happen in mammals, but if artificial respiration be performed they come on (*loc. cit.*, p. 69).

In regard to the action of prussic acid upon the blood-vessels, we have very little knowledge; the only records bearing upon the subject that I have met with are two experiments of Wahl (*De Vi et Effectu Acido hydrocyanato ad Curationem attribuendis*, Diss., Bonn, 1865), who found that the blood-pressure rises under the influence of the acid.

THERAPEUTICS.—Our knowledge of the physiological action of prussic acid does not lead to a belief in its wide applicability to the relief of disease, and I think clinical experience has demonstrated that it is of little value except in meeting three indications: first, *to allay cough*; second, *to relieve irritation of the gastric nerves*; third, *to allay irritation of the peripheral sensitive nerves*.

There appears to be in the profession a very wide-spread belief in the power of this remedy to allay cough; at least it is very largely used for this purpose in cough-mixtures, either itself or in the form of cyanide of potassium. I have employed it in a great number of cases in hospital practice, and apparently with good effect, although, as it was always given in combination with such remedies as morphia, it is difficult to say how much of the result was due to it. I do not believe it can compare with such narcotics as opium or hyoscyamus in its ability to fulfil the present indication.

There can be, on the other hand, no doubt as to the value of prussic acid in certain stomachic affections, especially nervous vomiting and gastralgia. When the pain is accompanied by decided dyspeptic symptoms, the remedy will sometimes succeed, but more often fails. Even in the most favorable cases it does not always afford relief; and as the relief when it does occur is immediate,

or at least is very soon apparent, it is useless to persist long in the exhibition of the remedy. In these cases its action is probably local, as it certainly is when the acid is employed to relieve itching in *prurigo* and other cutaneous diseases. For this purpose it is used as a wash (fʒss to fʒi in fʒi), but great care must be taken to avoid constitutional effects, especially when there is any abrasion of the skin. Very serious results are said to have been caused by its absorption when carelessly used in skin-diseases.

Prussic acid has been commended as an arterial sedative; but it is evident that we possess numerous more efficient and far safer remedies of such character.

TOXICOLOGY.—The symptoms of prussic acid poisoning have already been mentioned: those of most value from a diagnostic point of view are the sudden occurrence of unconsciousness; the violent convulsions; the general paralysis; the peculiar character of the breathing, expiration being prolonged and forced; and the rapid results. The odor of prussic acid upon the breath is very often, but by no means always, present. When distinct, it is, of course, of very great diagnostic value. Leaving out of sight the cyanides, the only poison with which prussic acid could well be clinically confounded is nitro-benzole. The distinction is often very difficult, large doses of the latter substance killing almost as quickly as prussic acid, and inducing analogous symptoms. Caspar advises that after death the body be left open, exposed to the air, as the odor of prussic acid disappears rapidly, whilst that of the nitro-benzole is persistent. The diseases with which the poisoning may be confounded most readily are some forms of *apoplexie foudroyante*, and sudden failure of the heart's action. The diagnosis may, during life, be almost impossible. It has been asserted that stertorous breathing does not occur in prussic acid poisoning; but it has been present in several reported cases. (See *Taylor's Principles and Practice of Medical Jurisprudence*, Philadelphia, 1873, p. 363.)

An autopsy, however, ought generally to enable the physician to determine whether the case has or has not been one of prussic acid poisoning, if the symptoms during life are known.

The period at which death may occur after the ingestion of the poison is set down by Lonsdale at from one to fifty-five minutes; but a case is reported in *Guy's Hospital Reports*, 1868, p. 259, observed by Dr. Hilton Fagge, in which the fatal result was put off for at least an hour and a quarter after the ingestion of hydrocyanic acid. After death the body often presents a livid surface, bloated countenance, fixed, glassy eyes with dilated pupils, and clenched fingers; sometimes it offers nothing worthy of note except excessive rigidity, and the face may be very pale. When opened, the odor of prussic acid is generally, but not always, emitted; the mucous membrane of the stomach is very commonly found much congested, and the dark or cherry-colored liquid blood usually everywhere fills up the veins. The heart is soft and flaccid.

The treatment of poisoning by prussic acid is, unfortunately, as inefficient as it is simple. There is no known antidote to it. The stomach should, if

possible, be emptied, and the experiments of Preyer indicate that the hypodermic use of atropia as a respiratory stimulant may be efficacious; the inhalation of the vapors of ammonia, and the free exhibition of ammonia by the mouth and by injection into the veins, may be practiced. Artificial respiration has been found very successful by Preyer in animals when poisoned by small doses of prussic acid, and should always be assiduously practiced. Next to it in importance is the use of the alternate cold and hot douche, about a half of a small bucketful of cold water, and the same quantity of very hot (115° F.) water, being dashed upon the chest in rapid succession.

POTASSII CYANIDUM—CYANIDE OF POTASSIUM (KCy).

This salt is prepared, according to the U. S. Pharmacopœia, by heating together the ferrocyanide of potassium and the carbonate of potassium. It occurs in white, amorphous, opaque masses, having the odor of prussic acid and a taste of similar character, but somewhat alkaline. It is deliquescent, and readily soluble in water. When the nitrate of silver is added to its solution, there falls a precipitate of the cyanide of silver, which is wholly soluble in ammonia.

When the cyanide of potassium is taken into the stomach, the acids there present convert it into prussic acid, and the same change probably occurs, although more slowly, even when the salt is injected directly into the blood-vessels. The physiological, therapeutical, and toxicological properties of this salt are similar to those of prussic acid. Death, however, does not occur so soon as when hydrocyanic acid has been taken, and insensibility is sometimes not manifested for several minutes. Five grains of the salt have caused death in several cases. The therapeutic dose is from one-tenth to one-twelfth of a grain.

CYANIDE OF SILVER (AgCy), a white insoluble powder, is one of the preparations of the U. S. Pharmacopœia. It is used solely for making prussic acid.

VEGETABLE ACIDS.

Although the officinal vegetable acids differ so much from the other substances considered in the present class as not to be poisonous except in enormous doses, and although they are never used to produce a profound impression upon the circulation, yet since they have, or at least are believed to have, the power of lowering the force of the cardiac movements to some extent, and since they are so commonly believed to have a tendency to depress animal temperature as to be usually spoken of as *refrigerants*, the present seems to me a fitting position for their consideration.

ACIDUM TARTARICUM—TARTARIC ACID.

Tartaric acid occurs in large, hard, transparent, six-sided prisms, which are pyro-electric and phosphorescent when rubbed in the dark, and are nearly

free from odor, but have a very sour taste. In the shops the acid is almost always kept in the form of powder. Tartaric acid is the acid of the grape, and occurs in grape-juice as a supertartrate of potassium. When the juice undergoes fermentation and alcohol is developed, the acid salt, not being soluble in the newly-formed menstruum, precipitates, collecting as a dark mass in the wine-casks, whence it is sent into commerce under the name of *argol* or *tartar*. Out of this substance the acid is manufactured by treating with lime, so as to form a tartrate of calcium, and precipitating this new compound in its watery solution by sulphuric acid, sulphate of calcium falling, tartaric acid remaining in solution.

Tartaric acid is soluble in little more than half its weight of hot water and less than its weight of cold water. It is distinguished from all other acids by forming a crystalline precipitate (bitartrate) when added to a neutral solution of potassa.

PHYSIOLOGICAL ACTION.—When applied to a denuded surface, or in sufficient concentration to a mucous membrane, tartaric acid acts as a very decided irritant, and even upon the skin its saturated solution after a time causes redness and burning.

When the drug is taken internally in sufficiently large doses, it acts as an irritant poison, causing violent œsophageal and gastric burning, vomiting, and, it may be, fatal gastro-enteritis. Upon animals it acts, in large doses, precisely as it does upon man. Thus, Mitscherlich states that three or four drachms suffice to kill a rabbit, the evident symptoms being great weakness of the heart's action, difficult and slow breathing, steadily-increasing pains, with slight convulsions before death. According to Devergie, it requires nearly half an ounce to kill a dog when given by the stomach; but Pommer (quoted by Husemann) asserts that one grammé (15.34 grains) injected into the crural vein of a dog will produce death.

Tartaric acid is never used internally by practitioners in such doses as to cause any of the symptoms above detailed, and it is evident that these symptoms throw little light upon its action in therapeutic doses, except to render it somewhat probable that the tendencies of the medicine are to lower cardiac action. This probability is increased very much by the experiments of Bobrick (quoted by Husemann, *Die Pflanzenstoffe*, p. 561), who found that very large doses render the heart's action weaker and slower.

A great deal of interest to the therapist centres in the question as to what becomes of the acid in the system. Unfortunately, our knowledge in regard to this matter is far from complete.

The acid probably is partially burnt up in the body, and is partially eliminated by the kidneys. Wöhler,* in his experiments, found it in the urine in the form of tartrate of lime, whilst Buchheim* and Piotrowski* could find only a very small percentage of the ingested acid in the urine, and conclude

* All these are quoted by Husemann, *Die Pflanzenstoffe*. I have not seen the originals.

that it is mostly destroyed in the body. Dr. Münch (*Archiv des Vereins für Gemein. Arbeiten*, 1863, p. 370) finds that when tartaric acid or citric acid is given it soon appears in the urine. Dr. H. Bence Jones (*Medical Times and Gazette*, vol. ix., 1854, p. 408, and *Lectures on Pathology and Therapeutics*, London, 1867) has found that both citric and tartaric acid cause a very great increase in the acidity of the urine of persons taking them, and are apt also to give rise to the presence of free uric acid in the excretion. Unfortunately, Dr. Jones did not attempt to determine whether the increased acidity was or was not due to the presence of the vegetable acid in the urine.

THERAPEUTICS.—Tartaric acid is very rarely used in medicine, the citric acid almost always being preferred. It may, however, be employed whenever it is desired to render the urine acid, in doses of ten to twenty grains; but it is less valuable than the acid of the lemon.

TOXICOLOGY.—There are, I believe, but two fatal cases of tartaric acid poisoning on record: one reported by Devergie (*Ann. d'Hygiène*, 1851, t. ii.), and one by Prof. Taylor (*Principles and Practice of Medical Jurisprudence*, London, 1873, p. 230), in which death took place nine days after the ingestion of an ounce of the poison dissolved in half a pint of water. The treatment of tartaric acid poisoning consists in the free exhibition of magnesia, of lime, of the carbonate of potash or of soda, or of any article, such as soap, containing an alkali in a suitable shape, which may be at hand. The after-treatment resolves itself into that of gastro-enteritis.

ACIDUM CITRICUM—CITRIC ACID.

Citric acid is the acid of lemon- and lime-juice, from which it is extracted by a process precisely similar to that employed in the manufacture of tartaric acid. It occurs in rhomboidal prisms, which are sometimes very large, are nearly free from odor, but are possessed of a very sour, almost corrosive taste, which, when the acid is in sufficiently weak solution, is quite pleasant. Citric acid is soluble in three-fourths of its weight of cold water, half of its weight of boiling water, soluble in alcohol, insoluble in ether.

It is sometimes adulterated with tartaric acid, which may be readily detected by the addition of a strong neutral solution of the carbonate of potassa to a strong solution of the suspected drug, when, if tartaric acid be present in any amount, a precipitate of the bitartrate is formed.

PHYSIOLOGICAL ACTION.—Citric acid in concentrated solution certainly acts upon abraded surfaces, and upon mucous membranes, as an irritant, but, according to Mitscherlich, is less irritant than tartaric acid, since its concentrated solution has no action upon the sound skin.

No case of poisoning by citric acid has occurred in man, that I am aware of, and Piotrowski (quoted by Husemann, *Die Pflanzenstoffe*, p. 561) took, in six hours, thirty grammes, an hour later fifteen grammes, and an hour later thirty grammes, or nearly two ounces and a half in all, with the induction of no more serious symptom than vomiting. It is, therefore, somewhat doubtful

whether citric acid is capable of causing death in man. This difference in action between it and tartaric acid may depend upon the latter being so much the more irritant of the two; upon the urinary secretion their action is probably similar.

THERAPEUTICS.—Citric acid is sometimes itself employed in medicine, but is almost exclusively used in the form of *Lemon-juice* (SUCCUS LIMONIS), which some, it is true, have thought to be dependent upon citrate of potash for much of its virtue, but which contains, as shown by the analysis of Prof. H. Bence Jones (*Medical Times and Gazette*, vol. ix., 1854), in every ounce twenty-six or twenty-seven grains of free citric acid, and not two grains of the citrate of potash.

Lemon-juice has several very distinct uses in medicine, all of them resting upon clinical rather than physiological data. The chief and most important of these is in the cure and prevention of *scurvy*. During the disease it should be drunk freely in the form of lemonade, three or four ounces of it being taken daily. As a prophylactic against the disease, lemon-juice is simply invaluable, but it is absolutely necessary that it be of good quality. It may be prepared for long voyages in one of two ways: first, boil the juice slightly, strain, allow to cool, pour into bottles up to their necks, fill the vacant space above with pure olive oil, cork tightly, and keep the bottle upright; second, add ten per cent. of brandy, and bottle as before (*Medical Times and Gazette*, 1854, p. 635). Citric acid is of some value in *scurvy*, but is incomparably less active than lemon-juice.

In *acute rheumatism*, benefit may be derived from the free use of lemon-juice as originally proposed by Dr. Rees, of London. One or two ounces of it may be given four or five times a day; but it is certainly less efficacious than the alkalies.

In *catarrhal jaundice*, and in habitual *torpor of the liver*, the free internal use of lemon-juice often aids in effecting a cure. In *fevers*, lemonade often affords a very refreshing and useful refrigerant drink.

ADMINISTRATION.—Lemon-juice, when it can be had, should always be preferred to citric acid; when only the latter is available, an artificial lemon-juice may be made by dissolving in a pint of water an ounce of the acid with which four drops of the oil of lemon have been well rubbed up.

ACETUM—VINEGAR.

The physical properties of vinegar are too well known to need description here. That best suited for medicinal use is in this country prepared from cider, and should have a trace of the taste of cider. It is sometimes adulterated with sulphuric acid, which may be at once detected by boiling with chloride of calcium, which precipitates any free sulphuric acid as sulphate of calcium, without affecting the small proportion of soluble sulphates existing in vinegar.

Vinegar may be substituted for lemon-juice as the basis of an acidulous

drink in fever when the lemon-juice is not to be had ; but as an *antiscorbutic* it is certainly very much inferior to it, and has not, that I am aware of, been tried in *rheumatism*.

By the distillation of vinegar is prepared, according to the U. S. Pharmacopœia, the *Distilled Vinegar* (ACETUM DESTILLATUM), which is used exclusively in pharmacy.

Acetic Acid (ACIDUM ACETICUM) of the U. S. Pharmacopœia is a colorless liquid, having a pungent odor, free from empyreuma, and an intensely acid, corrosive taste. It contains thirty-six per cent. of the monohydrated acetic acid, and has a specific gravity of 1.047. Glacial or monohydrated acetic acid is not officinal. It is a colorless liquid, crystallizing at 34° F., and actively escharotic,—in a measure, no doubt, owing to its properties of dissolving gelatine and gelatinous tissue and of effecting a partial solution of albuminous matters. *Dilute Acetic Acid* (ACIDUM ACETICUM DILUTUM) is officinally prepared by the addition of seven parts of water to one part of acetic acid, and should have the sp. gr. 1.006.

Dilute acetic acid or its equivalent, vinegar, is a useful topical application in various superficial inflammations of the skin, such as "*sunburn*," and in *sprains*. Applied to the skin, it acts as a powerful stimulant and astringent, causing contraction of the vessels and great whiteness. Diluted with two or three times its bulk of water, it is occasionally employed as an injection against *seat-worms* ; but the infusion of quassia is preferable.

The use of acetic acid as a caustic will be spoken of under the heading of Escharotics.

TOXICOLOGY.—Acetic acid in any of its more concentrated forms is a corrosive poison, and death has been produced by it in at least one case (Orfila, *Toxicologie*, t. ii.). The symptoms resemble those caused by mineral acids, and the treatment is exactly similar,—neutralization by an alkali or its carbonate, or by some substance, such as soap, containing an alkali, and the meeting of indications as they arise.

Oxalic Acid, although never used in medicine, has so frequently caused death by its poisonous influence as to merit notice here. As a poison it figures in two forms: that of simple oxalic acid, and that of the *acid oxalate of potassium*, or *salt of sorrel*, or *essential salt of lemons*, as it is variously termed in common parlance. The symptoms produced are a hot acrid taste experienced during the swallowing, a burning in the gullet, soon extending to the stomach, intense abdominal pain, vomiting of highly acid, greenish, blackish-brown or bloody mucus (rarely of arterial blood), collapse, livid surface, cold skin, entire prostration of strength, small irregular pulse, and finally stupor, passing into unconsciousness and death. In some cases the gastric symptoms are very prominent; in others they are nearly wanting, and the chief manifestations are collapse and such nervous symptoms as almost complete general paralysis, numbness, and finally stupor. According to Taylor, the smallest quantity which is known to have caused death is one drachm.

An ounce usually proves fatal, but has been recovered from. In poisoning by oxalic acid, the immediate administration of an antidote is of the utmost importance. As the oxalates of potassium and of sodium are poisonous, neither potash nor soda is available; but fortunately lime or chalk is a perfect antidote to oxalic acid, forming the excessively insoluble oxalate of lime. As time is a matter of so much importance, very often it is best simply to scrape the whitewash off a wall, ceiling, fence, or wherever it may be at hand, rub it up hastily with water, and administer it freely. The after-treatment is that of toxic gastro-enteritis.

CLASS V.—ANTISPASMODICS.

UNDER the name of *Antispasmodics* are grouped in this treatise a number of medicines generally of very feeble powers, but of frequent use. In certain conditions of the nervous system—conditions associated with weakness rather than with simple depression—the nerve-centres appear to be more susceptible than is normal to external impressions, as well as to those impulses which originate in the cerebral centres themselves and are connected with the emotions. As a result of this state, various symptoms arise, of trifling import, but often apparently severe, and always annoying. Such symptoms, in their mildest form, constitute the state of unrest known as *nervousness*; in their severer type they may rise in intensity up to the wildest convulsion of *hysteria*. It is in this class of affections that the so-called antispasmodics are useful. As the condition which they relieve is always associated with weakness, they are often spoken of as “nervous stimulants.” There is, however, but little evidence of their increasing power or functional activity when given to healthy individuals. Some of them act very slightly upon the circulation when given in very large doses, and a few when administered as freely as possible induce slight cerebral symptoms, such as vertigo; but, excepting camphor and Hoffman’s Anodyne, none of them are capable of producing serious poisoning. As any theory of the method in which the hysterical convulsion originates—of its immediate causes and the mechanism of its production—would, with our present knowledge, be at best but an ingenious speculation, I think the safest plan in regard to the action of drugs belonging to the class now under consideration is to accept the teachings of clinical experience as to facts, and to avoid theorizing as to the way in which the results are brought about.

MOSCHUS—MUSK.

A highly odorous, unctuous substance, obtained from the glands situated just in front of the preputial orifice of the *Moschus moschiferus*, or musk-deer of Thibet. The genuine musk-sac is to be distinguished from imitations of it by the hairs being arranged concentrically around a minute orifice. As it occurs in commerce, musk is very greatly adulterated.

PHYSIOLOGICAL ACTION.—Musk appears to act upon the nervous system simply as a mild stimulant and antispasmodic. Jörg and Sundelin have experimented with it upon healthy men with somewhat contradictory results.

According to the first-named observer, twenty grains of it induce exhilaration without lassitude, but, according to the latter authority, may cause giddiness, drowsiness, and lassitude. Both observers noted a slight increase in the frequency of the pulse. It seems to me evident that the action of musk upon the organism is a very feeble and uncertain one. Yet there is considerable clinical evidence that when the nervous system is exhausted it is of service in calming restlessness and equalizing the disturbed balance of nervous power.

THERAPEUTICS.—Musk is at present very little used, but is strongly recommended by some of the older writers in various spasmodic affections, especially in *hysterical convulsions*. In *hiccough* it has been considered a specific.

In the *delirium of low fevers* it is sometimes valuable; and it has been especially commended by Trousseau in the *ataxic pneumonia* of drunkards, when the nervous symptoms are out of proportion to the local disease.

The dose of musk is from five to fifteen grains, which may be given suspended in mucilage.

CASTOREUM—*Castor*.—The preputial follicles of the beaver; pyriform sacs, occurring in pairs, and containing an unctuous material.—the officinal *castor*. This is a substance allied to musk in its physical properties, and apparently also in its medicinal qualities. It is certainly less efficient, however, than musk, and Mr. Alexander (*Pereira's Materia Medica*, American edition, 1866, p. 949) is said to have taken a quarter of an ounce of it without having experienced any effect. It is employed in the same affections as musk, in doses of fifteen to sixty grains, suspended in mucilage. There is an officinal *tincture*, which is stronger in alcohol than in castor.

VALERIANA—VALERIAN.

The root of the *Valeriana officinalis*, an herbaceous perennial of Great Britain. It consists of a short, yellowish-white rhizome, with numerous fibrous roots, of a bitter taste and peculiar odor. The active principle of valerian appears to be the *oil of valerian*, which, according to Pierlot, consists of a mixture of valerian camphene, valerian camphor, valerianic acid, resin, and water.

PHYSIOLOGICAL PROPERTIES.—Upon cats valerian has a very extraordinary effect, attracting them strongly, and greatly exciting their sexual passions. It is possible that this action is suggestive, due rather to the resemblance of the odor to that of the animals during sexual excitement, than to a direct action of the drug.

Valerianic acid given to rabbits in large doses produces, at first, a slight acceleration of the pulse, which, with the respiration, afterwards becomes less frequent than normal, and at the same time lassitude and muscular weakness are developed. Enormous doses kill rabbits somewhat suddenly, or produce fatal gastro-enteritis.

Upon man, large doses (℥ii to ℥iv) are said to produce a feeling of warmth in the stomach, and sometimes nausea, vomiting, and colicky pains. The

pulse is generally slightly quickened, and a sense of exhilaration is induced, accompanied, however, by formication in the hands and feet. Very large amounts cause a feeling of heaviness, and even of pain, in the head.

THERAPEUTICS.—Clinical experience has demonstrated the value of valerian as a means of relief for the milder forms of functional disturbance dependent upon a weak and over-excitabile or an exhausted nervous system. In the state of unrest familiarly known as “*nervousness*,” by soothing and quieting the patient, it will often indirectly procure sleep.

In *hysteria* it has been the most frequently used of medicines, and its action is oftentimes most happy.

It has also been employed, but with more doubtful advantage, in *mania a potu*, and in the *delirium of adynamic fevers*. In these cases it is almost invariably conjoined with more powerful remedies, and it is very difficult to decide how far it assists in procuring the beneficial result.

ADMINISTRATION.—The best preparations of valerian are the *fluid extract* and the *ammoniated tincture*.

Infusum Valerianæ—*Infusion of Valerian* (℥ss to Oj). Dose, a wine-glassful.

Tinctura Valerianæ—*Tincture of Valerian* (℥ii to Oj). Dose, one to three fluidrachms.

Tinctura Valerianæ Ammoniata—*Ammoniated Tincture of Valerian* (℥ii to Oj). Dose, one to three fluidrachms.

Extractum Valerianæ Fluidum—*Fluid Extract of Valerian*. Dose, one fluidrachm.

ACIDUM VALERIANICUM.—*Valerianic Acid* is an oily, colorless liquid, of a caustic taste, and a strong odor, resembling, but different from, that of valerian. It is made by the action of chromic acid upon amylic alcohol, by a somewhat complicated process, and is officinal for the manufacture of *Valerianate of Ammonium* (*Ammonii Valerianus*), a white salt occurring in quadrangular plates, which effloresce in a dry and deliquesce in a moist atmosphere, have the odor of valerianic acid and a sharp sweetish taste, and are very soluble in water and alcohol.

THERAPEUTICS.—This salt was introduced by M. Déclat, of Paris, as a remedy for *neuralgia*. It has since been used very largely for *nervous headache* and in *hysteria*. It appears to be about equivalent to valerian, but, unless it be in nervous headaches, is less efficient. The dose of it is ten grains, which is generally administered in the form of an *elixir*.

ASSAFŒTIDA—ASSAFETIDA.

An exudation obtained by incising the living root of the *Narthex Assa-foetida*, an umbelliferous plant of Afghanistan. It mostly occurs in irregular opaque masses of a dull yellowish- or pinkish-brown, white when freshly broken, of a bitter acrid taste and a strong garlicky odor. Even this *lump*

assafetida is largely composed of tears agglutinated together; sometimes these tears are distinct and separate, when they constitute the variety known as *assafetida in tears*.

Assafetida is composed chiefly of gum and resin, but its properties are in great part due to the volatile oil, of which it contains from 3.5 to 4.5 per cent.

PHYSIOLOGICAL ACTION.—When taken into the stomach, *assafetida* acts as a local stimulant and carminative, and on this account is in some parts of the East used as a condiment. The oil is without doubt absorbed. The evidence as to its action upon healthy men is both scanty and contradictory. Thus, whilst M. Pidoux took half an ounce in a single dose without perceptible effects other than to render his secretions horribly offensive for two days, Jörg and his disciples found that in twenty-grain doses it produced gastric uneasiness and pain with alvine dejections, increased the pulse-frequency and animal warmth, quickened the respiration, and caused headache, giddiness, and erotic excitement.

THERAPEUTICS.—Clinical experience has abundantly proven that *assafetida* is one of the most efficient of the so-called antispasmodics, and may be given to fulfil the same indications as valerian in *functional spasm*, in *hysteria*, and in *nervousness*. It differs from valerian in having a much more decided action upon the mucous membranes. It is an excellent *carminative*, and in the form of injection is constantly used for the relief of *tympanitis*. It also in small doses increases the appetite, and affords relief in the *dyspepsia*, with flatulent colic and costiveness, of the aged or hysterical.

As a *stimulating expectorant and antispasmodic*, it is useful in *whooping-cough* and in *chronic catarrh*. It is especially efficient in palliating the latter affection as occurring in old people, when the difficulty of breathing is paroxysmally increased by spasm of the bronchial tubes.

In *infantile convulsions* and in *severe infantile colic*, *assafetida* enemata (fʒii to fʒss of the milk) are exceedingly useful and harmless.

ADMINISTRATION.—*Pilulæ Assafetidæ*—*Pills of Assafetida* (each containing three grains). Dose, two to four pills.

Mistura Assafetidæ—*Milk of Assafetida* (ʒii to Oj). Dose, half to one fluidounce. For injections, one to three fluidounces.

Tinctura Assafetidæ—*Tincture of Assafetida* (ʒii to Oj). Dose, half to one fluidrachm.

CAMPHORA—CAMPHOR.

Camphor is obtained in China, Japan, Cochin China, the Sunda Islands, etc., by boiling the comminuted wood of the root, stem, and branches of the *Laurus Camphora*, and skimming off the camphor as it rises to the surface of the water when cold. This camphor is then partially purified by sublimation, and comes into commerce as *crude camphor*, which is in grains of a whitish or pinkish color, and is finally purified by sublimation with lime.

Refined camphor (or, as it is commonly called, *camphor*) occurs in hemi-

spherical bowl-like translucent masses, of a fibrous or granular fracture. Its taste is hot and peculiar; its odor very strong and characteristic; it is volatile, inflammable, tough, but readily pulverized on the addition of a few drops of alcohol; melts at 347° F.; is soluble in one thousand parts of cold water,* in one part of strong alcohol, and still more soluble in chloroform; thrown upon water, a granule of camphor floats, and exhibits a rotary motion.

By slow sublimation at ordinary temperatures, camphor can be made to crystallize in handsome hexagonal tables.

PHYSIOLOGICAL ACTION.—Locally applied, camphor is a decided irritant, although when it is taken into the mouth a sense of coolness after a time is experienced, due no doubt to the volatility of the drug: a precisely analogous phenomenon occurs with some other volatile irritants, such as oil of peppermint.

Great differences of opinion have prevailed in regard to the action of camphor upon man, and it is scarcely doubtful that it acts differently upon different persons, or at least that doses which in some cause only exhilaration in others produce general depression. When a moderate dose (five to ten grains) of camphor is taken, a feeling of exhilaration is usually induced, a sense of comfort and quietness, especially marked in those previously suffering from "nervousness;" the pulse may be somewhat accelerated, although it is undoubtedly not markedly affected in the majority of cases, and Trousseau saw it fall after the ingestion of ten grains of the drug. After larger doses (twenty to thirty grains) the pulse is usually lowered in frequency, and giddiness, with a feeling of lassitude, is produced, preceded, it may be, by a short period of exhilarative excitement. After poisonous doses (thirty to sixty grains) the symptoms, which are tolerably uniform, are as follows: faintness, vertigo, confusion of ideas, delirium, violent convulsions, insensibility, apparent general paralysis; a pulse generally small, but sometimes accelerated and sometimes lowered in number; a skin cool, pale or livid, generally bedewed with sweat. In any individual case many of these symptoms may of course be wanting.

No death has occurred in the adult directly from camphor. A young, sickly infant was killed by ten grains of it; and a woman four months gone in pregnancy, having taken three drachms of camphor, suffered the usual symptoms, followed by an abortion which proved fatal.

Much contradictory evidence might be adduced as to the influence of camphor upon the genital organs. The truth evidently is that its action varies according to the dose and the idiosyncrasies of the patient. In the great majority of instances, I think camphor in moderate doses has no decided influence upon the sexual system; at least I have seen many hundred such doses taken and have never yet seen any aphrodisiac effect. In some per-

* By rubbing the gum up with magnesia in water, the latter can be made to take up much more than one part in one thousand.

sons, however, full therapeutic doses are said to cause sexual excitement. In regard to very large doses, the testimony is quite uniform that if they exert any action it is to lessen the erotic feelings.

Upon articulates camphor acts as a poison, and, according to Menghini, in birds it causes epileptiform convulsions, with stupor or delirium. In dogs large doses of it excite violent convulsions, with vomiting and finally death, apparently from paralysis of respiration. Its action upon the circulation has been partially studied by Dr. O. Heubner (*Archiv der Heilkunde*, 1870), who found that in rabbits it never produced a rise in the arterial pressure, but that in the frog, in doses of one milligramme to one and a half milligrammes, it acts as a cardiac stimulant, slowing the pulse-rate but increasing the force and energy of the contractions.

THERAPEUTICS.—Camphor is very largely used internally as an antispasmodic, to quiet restlessness and “*nervousness*.” It is also employed in certain painful affections seen in those persons who are especially liable to the condition of the nervous system just mentioned: thus, it is often useful in *nervous headaches* and in *dysmenorrhœa*. Indeed, in the latter disease, either alone or combined with opium in bad cases, it is a most valuable drug, but must be given freely.

In *diarrhœa* not dependent upon inflammation, in *cholericæ*, and even to some extent in *cholera*, camphor is a very efficient remedy, allaying intestinal pain and spasm, and also checking intestinal secretion. It enters into a large proportion of the popular cholera-mixtures.

In *adynamic fevers* it has been very greatly employed, but is of doubtful advantage: still, a good deal of testimony could be adduced in favor of its usefulness in sustaining the system in the low stages of these diseases; and in nervous restlessness occurring at such times it is often very soothing.

In *abnormal sexual excitement*, and in *chordee*, large doses of camphor very often appear to act most happily.

The drug has also been very frequently exhibited in various spasmodic affections, such as *whooping-cough*, *epilepsy*, and even *puerperal* and *strychnic convulsions*, but is, I believe, at present never so employed. In *hysterical convulsions*, as in other phenomena of the same origin, camphor is a useful antispasmodic.

Externally, camphor is much used in liniments as a stimulant application for bruises, sprains, etc.

ADMINISTRATION.—Large doses (ten to fifteen grains) of camphor are best administered in emulsion, because when given in this way, being very finely subdivided, they create as little irritation as possible, and are rapidly absorbed; smaller doses may be given in pill. As an antispasmodic, the *Camphor Water* (*Aqua Camphoræ*, ℥i to Oj) is usually preferred. Dose, half a fluid-ounce to two fluidounces. The *Spirit of Camphor* (*Spiritus Camphoræ*, ℥ii to Oj) is also officinal. Dose, half a fluidrachm to a fluidrachm. For external use, the Pharmacopœia recognizes the *Linimentum Camphoræ* (camphor

one part, olive oil six parts) and the *Linimentum Saponis*, or *Soap Liniment*,—a mild liniment, very popular by itself, or as the basis of more stimulating preparations.

With iodine and bromine camphor unites to form compounds. According to Laurent, *bromcamphor* occurs in red orthorhombic crystals. These when exposed to the air undergo rapid spontaneous decomposition, but by heating in a closed vessel are resolved into hydrobromic acid and a substance often known by the absurd name of *monobromate of camphor*, a compound in which one atom of hydrogen of the camphor has been replaced by bromine.

Brominated camphor is a crystalline solid, or occurs in large acicular crystals several inches long. It has been introduced by Prof. Deneffe (*Birmingham Medical Review*, April, 1872) as a nervous sedative, and as useful as an antispasmodic, especially in *delirium tremens*. Its value in the latter disease has been confirmed (a single case) by Dr. A. M'Lean Hamilton (*New York Medical Journal*, July, 1872), who also commends it highly in *chordee*. Dr. Wm. A. Hammond (*Ibid.*, May, 1872) recommends it in *hysteria* and in the *convulsive irritation* of teething. The dose for an adult is five grains, repeated cautiously every hour.

SUCCINUM—AMBER.

Amber is a fossil resin found on the southern coasts of the Baltic and in other portions of the world. It is not itself officinal or used in medicine, but by destructive distillation yields an empyreumatic oil, which is included in the *Materia Medica* list of the U. S. Pharmacopœia.

OLEUM SUCCINI—OIL OF AMBER.

This, as first obtained, is thin and yellowish, but soon becomes thick and black, in which state it occurs in the shops. Copal and damar resin are sometimes substituted for amber in its manufacture, and yield a product scarcely distinguishable from the genuine oil. It is not itself used in medicine; but when purified by distillation it yields the officinal *Oleum Succini Rectificatum*,—*Rectified Oil of Amber*,—an amber-colored liquid, of a hot taste, and a very strong, disagreeable odor.

THERAPEUTICS.—Rectified oil of amber is a powerful local irritant, and has been used as a rubefacient in chronic *rheumatism* and similar disorders. It is also an efficient antispasmodic, and as such is used in *hysteria*, in *whooping-cough*, and in *infantile convulsions*. In the *bronchitis* of infants, with severe nervous symptoms, as well as in the last two affections previously named, it is very useful as a counter-irritant and nerve-stimulant when diluted with from one to three parts of olive oil and freely applied over the spine. In obstinate *hicough*, given by the stomach, it is probably, next to musk, the most efficient remedy. Dose, ten to twenty drops, given in emulsion.

SPIRITUS ÆTHERIS COMPOSITUS—COMPOUND SPIRITS OF ETHER.

Hoffman's Anodyne consists of alcohol a pint, ether half a pint, and ethereal oil six fluidrachms. It is a colorless, inflammable liquid, of an aromatic, ethereal odor, and a burning, slightly-sweetish taste. Its specific gravity is 0.815. Hoffman's Anodyne is sometimes offered for sale without the ethereal oil. Forty drops of the genuine preparation will render a pint of water distinctly milky; but if no oil of wine be present, milkiness will not occur.

Ethereal oil, which is used in the above preparation, is a transparent, nearly colorless, volatile liquid, of a peculiar aromatic odor, and sharp, bitter taste. Its specific gravity is 0.91. It is *heavy oil of wine*, prepared by the action of an excess of sulphuric acid on alcohol, and diluted with equal parts of strong ether.

THERAPEUTICS.—We have little or no knowledge in regard to the action of heavy oil of wine upon the system. Clinical experience has shown that Hoffman's Anodyne is more persistent in its effects upon the nervous system than an equivalent amount of ether. It is a most useful antispasmodic in all the disorders for which such remedies are employed, especially when there is a tendency to failure of the circulation. Hoffman's Anodyne is also a very efficient carminative. The dose of it is one or two fluidrachms, repeated in half an hour or an hour if required. It should be given in cold water.

HUMULUS—HOPS.

The strobiles of *Humulus lupulus*, or the hop-vine, cultivated in northern and middle Europe and in the United States. Hops are soft, greenish cones, one to two inches in length, composed of thin, leaf-like, imbricated scales, having a bitter taste and a heavy narcotic odor. At the bases of the scales is a yellowish powder, officinal under the name of *Lupulina*. *Lupulin* is in minute grains, and contains, according to Payen, 2 per cent. of volatile oil, 10.30 per cent. of bitter principle, and 50 to 55 per cent. of resin. Volatile oil of hops is yellowish, and has a strong odor of the drug, and an acrid taste. The bitter principle has been obtained by Lerner in brilliant rhombic columns, of an acid reaction.

THERAPEUTICS.—Hops are a bitter tonic, and a very feeble narcotic, producing, when taken very freely, some heaviness, and perhaps sleep. They are especially useful as tonics in cases of nervous irritability requiring medicines of the class.

In *delirium tremens* they are very largely used to quiet nervous irritability, to aid more powerful remedies in procuring sleep, and at the same time to strengthen digestion.

In *priapism*, in *irritation of the bladder*, and in *abnormal sexual excitement*, hops have been exhibited with asserted benefit. They may be tried in large doses, but very often will fail.

Externally, hops are employed in the form of poultice, and when fresh certainly seem to aid the heat and moisture in allaying pain.

ADMINISTRATION.—The preparations of Lupulin are preferable to those of hops when a narcotic action is desired.

Infusum Humuli—*Infusion of Hops* (℥ss to Oj). A feeble preparation, which may be taken *ad libitum*.

Tinctura Humuli—*Tincture of Hops* (℥v to Oij). Dose, half a fluidounce to three fluidounces.

Tinctura Lupulinæ—*Tincture of Lupulin* (℥ii to Oj). Dose, half a fluidounce to two fluidounces.

Oleoresina Lupulinæ—*Oleoresin of Lupulin*. An excellent preparation, which may be given in pills, if desired. Dose, ten minims to a fluidrachm.

Extractum Lupulinæ Fluidum—*Fluid Extract of Lupulin*. Dose, half a fluidrachm to two fluidrachms.

A *hop poultice* is sometimes made by simply moistening with hot water the hops contained in a gauze bag of the required size and shape. In many cases a mixture of the broken strobiles with an equal part of Indian meal affords a more eligible preparation.

CIMICIFUGA—BLACK SNAKEROOT.

The root of *Cimicifuga racemosa*, an indigenous herbaceous plant, growing abundantly in rich, shady woods, attaining a height of six or seven feet, and readily distinguished by its very large multi-compound leaves and its long-branched spikes of whitish polyandrous flowers, naked when open. The root consists of a knotted head, with numerous fine, brittle rootlets; the odor is faint, the taste bitterish, somewhat astringent and acrid. It has not yet been determined exactly upon what the activity of *cimicifuga* depends. Mr. Geo. H. Davis has found in it a volatile oil, which Prof. Geo. B. Wood thinks is very probably active, since the virtues of the drug deteriorate on keeping. There are also two resins in the root.

PHYSIOLOGICAL ACTION.—Although *cimicifuga* has been most extensively used by American practitioners since the publication of the paper of Dr. Young (*American Journal of the Medical Sciences*, vol. ix.) in 1831, yet we have very little accurate knowledge as to its physiological action. Dr. Chapman (*Elements of Therapeutics*, 6th ed., vol. i.) affirms that in full doses it causes some nausea, more or less relaxation of the surface, vertigo, tremors, and decided reduction of the pulse; and Dr. N. S. Davis (*Transactions of the American Medical Association*, 1848, vol. i. p. 351) dwells very strongly upon its sedative action. In full doses it certainly influences the cerebrum, producing intense headache and giddiness. Although in large doses it vomits, yet its emetic action is never violent, and it never causes more than relaxation of the bowels. Various practitioners have asserted that it acts upon the respiratory mucous membrane.

THERAPEUTICS.—*Cimicifuga* was originally proposed by Dr. Young as a

remedy in *chorea*, and since his time has been very largely used in that disease by American practitioners. In simple *chorea*, such as exists in children, its value is unquestionable. It must be given freely, and in most cases the simultaneous exhibition of iron and laxatives materially aids in effecting a cure.

In acute *inflammatory rheumatism*, *cimicifuga* has been highly recommended by Dr. Davis and other practitioners, but is at present very rarely, if ever, used.

In *chronic bronchitis* it is sometimes used with benefit, especially when the expectoration is free and hectic exists.

ADMINISTRATION.—As *cimicifuga* deteriorates by keeping, the fresh drug should always be used. The powder is preferred by some practitioners; but I have found the officinal *fluid extract* very active. The dose of the fluid extract is from twenty minims to a fluidrachm; of the powder, twenty to thirty grains.

CLASS VI.—ANALGESICS.

IN the class *Analgesics*, in this work, are placed those drugs whose chief clinical use is in the relief of pain. Of course, the *Anæsthetics* might also be discussed under this heading, but, as they make a very marked group of themselves, they are best considered as a separate class. The only drug, besides opium, which seems worthy of a place in the present division, is *cannabis Indica*.

OPIUM.

The inspissated juice of the unripe capsules of the *Papaver somniferum*, or poppy. It is obtained by incising the capsules with a small, sharp knife, and twenty-four hours afterwards scraping off the exuded juice with a blunt blade. Opium is produced in various parts of the world,—chiefly in Turkey, Asia Minor, Persia, and India, but also to a very slight extent in England, Germany, and the United States. Our market is almost exclusively supplied from Asia Minor, with the variety known as *Smyrna*, or *Turkey, Opium*. This occurs in masses from the size of the fist to that of a child's head, irregularly globular, more or less flattened, covered externally with the capsules of a species of *Rumex* or dock, hard externally, softer and of a reddish-brown color within, and of a strong narcotic odor and taste.

Smyrna opium is at times variously adulterated with gum, licorice, and other substances. Such specimens are said generally to want the *Rumex* capsules.

A rough but pretty fair test of the purity of opium is performed by drawing a piece of it across a sheet of white paper. If it be much adulterated, the mark will be continuous,—not interrupted, as it should be. Often the black color, the adhesive consistency, and the sweetish taste will also betray the nature of the sample.

On exposure to the air, opium becomes hard and brittle, and is readily reduced to a powder of a yellowish-brown color. It yields to water, alcohol, and diluted acids, forming dark-brown solutions. Ether does not extract all of its medicinal principles. It is a very complex body, containing the alkaloids morphia, codeia, narceia, narcotina, thebaia, papaverina, porphyroxia, cryptopia, meconia, opiania, paramorphia, besides meconic, thebolactic, and sulphuric acids, extractive matter, gum, glucose, fixed oils, a volatile odorous principle, and other substances of no importance. In regard to the propor-

tions of the more important principles. Messrs. Smith of Edinburgh obtained from 100 parts of fine opium 10 parts of morphia, 6 of narcotina, 1 of papaverina, 0.15 of thebaia, 0.03 of codeia, 0.01 of meconia, 0.02 of narceia, and 4 of meconic acid (*Pharm. Journ. and Trans.*, October, 1865, p. 183). Good opium should yield from nine to fourteen per cent. of morphia.

As meconic acid strikes a blood-red color with a persalt of iron, the latter affords a ready test for opium and the meconates.

PHYSIOLOGICAL ACTION.—When opium is taken in such dose as to produce its mildest physiological effects, it exerts a quieting influence, inducing a peculiar dreamy condition,—very generally a feeling of *bienfaisance*.—during which images and ideas float before the mind, and by their endless and effortless repetition shorten the time, which seems to lose itself in rest. It is commonly asserted that there is a stage of the action of opium in which the activity of the mental faculties is exalted. This may be so in some persons, and especially in those who have accustomed themselves to the use of the drug as a stimulant; but my experience is that in those who do not habitually take opium true mental power is, during all the stages of the action of the drug, diminished rather than increased. The state induced is rather the fabled calm of the lotus-eater than the energetic activity of production. Even in those who are accustomed to the use of opium as an aid to work, I think it is the imagination rather than the reasoning faculties which is excited by it.

After a length of time, varying according to the idiosyncrasies of the patient and the dose of the drug, the condition which has been noted gradually passes into sleep,—either light and dreamful, or natural, or heavy and deepening into stupor, according to the amount of the drug ingested.

After awakening from the sleep, the patient may return at once to his normal condition, but very often experiences a state of depression, as shown by languor, a little headache, nausea, or even vomiting, which may last for some hours.

After very large doses, the first stage of the action of opium is very short, or it may be entirely wanting, sleep coming on almost at once. Thus, I have seen deep coma produced in three minutes by a hypodermic injection of morphia. The symptoms of the second stage of opium-poisoning closely resemble those of congestion of the brain: the pupils are strongly contracted, the face more or less suffused, often deeply cyanosed; the pulse full, slow and strong; the skin generally dry and warm; the respiration slow and deep, and, it may be, stertorous; unconsciousness is apparently complete, but very generally the subject can be aroused by violent shaking or by shouting in his ear, but relapses at once when left to himself. When the patient is aroused, the respirations become more rapid, and the skin often regains almost at once its normal color. Death very rarely occurs during this second stage of opium-

poisoning. When the symptoms do not gradually ameliorate, the third stage, that of prostration, is developed. The coma is now profound, and to arouse the patient may be impossible; the pupils are absolutely contracted, or, as death approaches, are widely dilated; the respirations are slow, feeble, and imperfect, and often interrupted by intervals of death-like quiet; the countenance is at once pallid and cyanosed; the pulse grows more and more rapid and more and more feeble; the skin is cold and moist, finally covered with a clammy sweat. Even yet the patient may recover: if he do so, the return to life is most gradual; if he do not, death occurs generally by failure of the respiration, but amid an almost complete extinguishment of the vital functions.

Although the symptoms which have been narrated are those usually produced by opium, yet in certain individuals the drug provokes quite different phenomena. One of the most common of these departures from the ordinary course of symptoms is an excessive depression following the sleep produced by moderate doses of the medicine. This state is seen, so far as my experience goes, most usually in females of weak, nervous organization, such as are peculiarly liable to attacks of neuralgia. The symptoms are a feeling of weakness and prostration, often accompanied by chilliness, dull headache, and giddiness, but especially marked by intense nausea and frequent vomiting. Very frequently the latter does not occur so long as absolute rest in the horizontal position is maintained: indeed, an almost diagnostic sign of this affection may be found in the fact that the stomach is quiet so long as the patient keeps the head upon the pillow, but the distress occurs at once upon rising up. In some cases this condition of depression even replaces the normal second stage, so that opium, instead of inducing quiet sleep, will provoke, it may be, alarming depression and vomiting, either with or without drowsiness.

Thus, cases have been reported in which one-fourth of a grain, or a somewhat greater quantity, of morphia, hypodermically injected, has been followed at once by syncope, with struggling for breath, and apparently imminent or even present death.*

A second and rarer idiosyncrasy towards opium exists in those persons who are rendered by it very delirious, it may be even wildly so.

In certain cases of opium-poisoning, convulsions, either partial or complete, have occurred amidst the more usual phenomena.

Opium at first sight appears to act so differently upon the lower animals than upon man, that it seems necessary to discuss this action by itself.

In 1826, Charvet (*Pereira's Materia Medica*, vol. ii. p. 1035, Philadelphia, 1854) found that opium acts upon all classes of animals, inducing in

* See Report of the Committee on the Hypodermic Method of Injection, *Medico-Chirurg. Trans.*, vol. i.; see also *Medical Times and Gazette*, 1868, cases reported by Mr. Braine and by Mr. Roberts.

the invertebrata weakness or paralysis of the contractile tissue, with gradual sinking and death; in fishes, a weakened paralytic condition of the muscular system, associated with convulsions; in birds and mammals, paralysis, convulsions, and stupor.

These researches have been much extended by modern investigators, and in considering them we shall confine our attention to the vertebrata.

When one or two grains of opium are injected under the skin of a frog (Kölliker, *Virchow's Archiv*, Bd. x. p. 248; J. F. H. Albers, *Virchow's Archiv*, Bd. xxvi. p. 229), in from six to ten minutes a condition of excitability is induced, so that the least touch produces violent tetanic convulsions, which, a little later, also occur without obvious cause. After a time, these convulsions gradually give way to a deepening paralysis. The breathing, previously disturbed, becomes more and more shallow and imperfect, and finally is suspended.

Morphia acts, apparently, on frogs in the same manner as opium: at least Drs. Richard Gscheidlen (*Untersuchungen aus dem Physiolog. Laboratorium in Würzburg*, Bd. iii. p. 15) and W. Baxt (*Reichert's Archiv für Anatomie*, 1869, p. 128) have found that in large doses it induces the counterpart of the series of phenomena just described. The latter observer noted, however, that when a minute dose (15.25 milligrammes) was employed, immediately following the injection came a brief period of disquietude; one minute afterwards the frog returned to its normal state, in from six to ten minutes suffered a diminution of excitability, and in from twelve to fifteen minutes fell into a stupor which continued from four to ten hours. After awakening, the reflex excitability seemed greater than normal.

According to Kölliker, the opium-convulsions take place after the cord has been divided below the medulla, or even as low down as the third vertebra. In a single experiment, tetanus did not occur after division of the cord at the fifth vertebra; but the quietness was probably simply due to exhaustion, as the frog had already been poisoned for a length of time and had suffered section of the medulla and of the cord below the medulla. These facts seem to prove that the convulsions are reflex and of spinal origin. The convulsive movements which are present late in the poisoning must be, however, of peripheral origin, since in Albers's experiments (*loc. cit.*) they occurred in limbs whose nerves had been previously cut so as to sever all connection with the nerve-centres. Further, both Kölliker and Albers assert that some of the convulsions are epileptiform,—*i.e.*, of cerebral origin. If this be so, opium must induce in the frog three kinds of convulsions; but it can scarcely be doubted that those of reflex origin are the chief.

Kölliker, from his investigations, concluded that opium does not act upon the peripheral nerves of frogs; but the recent very elaborate and apparently accurate experiments of R. Gscheidlen (*Untersuchungen aus dem Physiolog. Laboratorium in Würzburg, zweiter Theil*, 1869, p. 1) have shown that morphia in small doses increases the excitability of the motor nerves and

afterwards depresses them ; after large doses the period of excitation is short, that of depression soon coming on ; and after enormous doses diminution of functional activity is at once manifested. Both Gscheidlen and Kölliker agree that neither the contractile power of the muscles nor the excitability of the motor nerves is destroyed by opium or morphia, although Albers (*Virchow's Archiv.* Bd. xxvi.) asserts that both are extinguished. Gscheidlen calls attention to this disagreement, and states that he has verified his own results by frequent experimentation with enormous doses of the alkaloid.

Experiments upon the sensory nerves are always unsatisfactory, but Gscheidlen (*loc. cit.*, p. 17), employing the method of Pflüger, found that morphia locally applied intensifies and protracts the excitability of an afferent nerve in cases of strychnic poisoning.

The peculiar convulsions prove that in the frog during opium-poisoning the reflex function of the cord is excited. Gscheidlen has confirmed this by direct experiment, and has also proven that in the latter stages, when the motor functions are depressed, the paralysis is largely of spinal origin, the reflex activity of the cord being greatly lessened.

Our knowledge of the action of morphia upon the nervous system of the frog may be summed up as follows :

Morphia in minute, non-toxic doses causes sleep, followed by augmentation of reflex activity.

Morphia in large, toxic doses produces violent convulsion, followed by paralysis ; the convulsions are chiefly spinal, and due to a heightened spinal activity, but are to some extent probably, also, of cerebral origin, and later in the attack arise from a direct action of the alkaloid upon the muscle or the nerve-endings therein ; the paralysis is caused by a depression of the cord and a diminution of the conducting power in the nerves.

Dr. S. Weir Mitchell has shown (*American Journal of the Medical Sciences*, Jan. 1869, and Jan. 1870) that birds, as represented by pigeons, chickens, and ducks, are very insusceptible to the toxic action of opium and its chief derivative, morphia. It appears to be impossible to kill a pigeon by opium given by the mouth, and of morphia from eight to fifteen grains are required to produce a fatal result ; but when given hypodermically from two to three grains of the alkaloid suffice. These results have been in great measure confirmed by Dr. B. W. Richardson, and are no doubt accurate. The symptoms induced have been very uniform : they are unsteadiness, labored breathing, increasing signs of apnoea, *unaltered* pupils, and, finally, general convulsions and death. No true hypnotic effect has been observed, but a curious and very great rise of temperature just before death was noted in one case. As Flourens affirms that a single grain of the aqueous extract of opium will throw a sparrow into a profound stupor, it can scarcely be allowed as proven that the drug acts upon all birds as upon those experimented with by Dr. Mitchell.

Upon dogs morphia acts very much as upon man.* In very many cases, if not in the majority, eight to ten grains of the alkaloid injected into a dog of moderate size will cause deep sleep, amounting to coma, so that the animal will remain in any position in which he may be placed. The length and depth of this sleep are, of course, proportionate to the dose: when at all profound, it is accompanied by marked insensibility to pinching and other forms of external irritation. A repetition of irritation, and especially a sudden loud noise or shaking, will, however, arouse the animal, precisely as in man. Indeed, sometimes the dog, even when comatose, seems more than normally sensitive to sudden noise, trembling and starting in an almost convulsive manner. After awaking, the dog shows unmistakable signs of nervous and psychical depression. In walking, the hind legs are dragged, as though semi-paralyzed; the eyes are haggard; the naturally brave animal cowers in a corner or seeks to hide himself, no longer recognizing his master, and does not return to his natural condition for many hours. After smaller doses, the dog exhibits effects proportionately less intense.

It has been shown by Harley that in some dogs, precisely as in some people, morphia fails to exert its usual hypnotic action, but produces great depression, as evinced by faintness, prolonged nausea, and retching, interrupted only by intervals of dreamy delirious somnolency.

In the horse (Harley, *loc. cit.*), two or even three grains of morphia hypodermically injected produce sometimes a slight drowsiness, sometimes no perceptible effect. Doses of from four to six grains cause great restlessness and accelerated pulse. The mouth is moist, the temperature of the skin and its secretion increased; the animal paws continually, and treads about in his stall with an almost rhythmical movement. After twelve grains, Harley noticed in some cases very great excitement, as shown by marked increase in the rapidity of the heart's action, by muscular rigidity and tremors, by the animal walking rapidly to and fro, slobbering and sweating profusely. In another horse, after an immediate strong erection of the penis and copious emission of semen, heavy sleep came on, interrupted after the third hour by the usual symptoms of excitement. Thirty-six grains of the acetate caused in a powerful hunter deep comatose sleep, commencing in fifteen minutes and lasting for three hours, when it was replaced by intense restlessness and severe delirium, continuing for seven hours. During this time the animal was perfectly blind.

Barbier's (quoted by Prof. Stillé, p. 656) previous experiments upon the horse had yielded results similar to those of Harley. He used larger doses, and found that four drachms of the aqueous extract of opium produced violent tremblings, apparent insensibility to external irritants, convulsions without coma, and death. One hundred grains of the acetate of morphia killed a horse by convulsions in three hours.

* Harley, *The Old Vegetable Neurotics*, p. 107, London, 1869. Claude Bernard, *Archives Générales*, p. 437, vol. ii., 6th series, 1864. J. J. Reese, *American Journal of the Medical Sciences*, Jan. 1871.

In the mouse, according to the experiments of Harley, the first effect of an injection of from one-twentieth to one-twelfth of a grain of morphia is a tonic cramp-like contraction of the muscles, especially of the trunk, of such character that periods of forced rest alternate with a slow, laborious creep, which seems to originate not in the limbs but in the trunk itself. There is in this state no tendency to somnolency, but, on the contrary, an abnormal sensitiveness to loud sounds, which cause the mouse to resume for a moment active running movements. The breathing is irregular, the pulse accelerated, and finally stupor develops itself, and coma deepens into death by apnœa; or, otherwise, recovery, preceded by convulsive movements of the hinder part of the body, is gradually brought about.

In reviewing the action of morphia upon the lower animals, it becomes very evident that whilst we are not at all in a position to explain all of the symptoms, yet two classes of phenomena are everywhere discernible,—*i.e.* the spinal and the cerebral,—and that the higher in the scale of life any given animal may be, the more marked are the brain-symptoms. These cerebral phenomena are mostly sleep and stupor; but, as is well known, in some human individuals morphia acts as a delirifacient; and it seems very probable that the peculiar restlessness of the horse under the influence of the alkaloid is due to delirium, and not to spinal excitement.

When looked at in this manner, it seems to me that morphia does not act so differently as is generally believed upon the lower animals and upon man. The immensely higher cerebral organization of the latter, with the immensely greater sensitiveness which it involves, makes the man correspondingly more susceptible to the cerebral action of the drug: hence not only is he affected by much smaller doses of the alkaloid than are the lower animals, but, as the spinal symptoms are triumphant in the frog because its spinal system is vastly more developed than its cerebral, so in man the cerebral symptoms mask the spinal, because in him the brain is more developed than the cord. The two creatures—man and the frog—occupy the two extremes of the series; between them is probably to be found every gradation.

The action of opium upon dogs and rabbits is sufficiently close to that upon man to enable us to reason from experiments upon the former as to the influence of the alkaloid upon the circulation and respiration in the latter. Indeed, so far as these functions are concerned, morphia appears to act identically in both instances.

In man, the circulatory phenomena are a slight primary evanescent acceleration of the pulse-rate (see Nothnagel, *Handbuch der Arzneimittellehre*, Berlin, 1870, p. 8), succeeded by slowing and increased fullness and force of the pulse, which is followed by a return to the normal pulse, or a great increase of rapidity and loss of strength during the third stage. R. Gscheidlen has found in rabbits and dogs after the injection of morphia, first an increase in the pulse-rate, then a decrease, and finally return to the normal pulse, or else increased rapidity.

The slow, full pulse of the second stage of opium-poisoning is due to an action of the drug upon the inhibitory cardiac nerves, as is also in great measure, if not entirely, the increased arterial pressure; for Gscheidlen (*loc. cit.*, p. 45) has experimentally demonstrated that after section of the vagi morphia is powerless to lower the pulse, and also that division of the nerves during the second stage of morphia-poisoning is followed by an extraordinary rise in the pulse-rate. That the peripheral ends of the vagi are stimulated was proven by the fact that cardiac arrest took place when the distal ends of the cut nerve were more feebly irritated than would suffice to affect the unpoisoned animal; that also the inhibitory cerebral centres are stimulated was demonstrated by the instantaneous very great fall of the pulse-rate, amounting in some cases to one-half in less than half a minute, which ensued upon the injection of a large dose of the alkaloid into the carotid,—*i.e.*, into the brain and the inhibitory centres.

The rapid, feeble pulse of the third stage of opium-poisoning Gscheidlen found to be due, at least in a measure, to paralysis of the peripheral vagi; for at such time stimulation of the peripheral end of the cut nerve was powerless to affect the heart.

The experiments of Gscheidlen also indicate that morphia exerts first a stimulating, then a depressing, influence upon the intracardiac motor ganglia, since, after isolation of the viscus by section of the cord, sympathetic, and pneumogastrics, life being sustained by artificial respiration, a large dose of morphia induced a momentary increase in the number of the cardiac contractions, followed by a marked decrease, and finally extinguishment, of the same.

The question of the action of morphia upon the vaso-motor system is of great interest, but cannot at present be fully answered. Gscheidlen believes that it first stimulates and then depresses it, and asserts that after the injection of a large dose the arterioles in the mesentery can be seen to contract, and later (third stage) to dilate.

The objections to this sort of evidence are sufficiently stated elsewhere in this book; and the rise of arterial pressure, which he also adduces as an argument, may be accounted for without calling upon the aid of the vaso-motor nerves. While, therefore, it is possible that morphia does exert the influence he claims for it, the question must be still considered as *sub judice*: that the vaso-motor system is not paralyzed, even *in extremis*, is shown by Gscheidlen's experiment (*loc. cit.*, p. 52), in which electrical stimulation of the cord at such time induced immediate rise of the arterial pressure.

Death occurs from opium, in the great majority of cases, by failure of the respiration; and that such failure is due to a direct action of the poison upon the respiratory centres in the medulla, is proven by the fact that morphia affects the breathing of dogs and rabbits whose pneumogastrics have been cut, as much as it does those whose nerves are entire (Gscheidlen, *loc. cit.*, p. 64).

Since morphia locally applied does not affect the pupil, it follows that its

constitutional action upon the latter is through the nerve-centres. It is probable, but has not, that I am aware of, been experimentally proven, that the contraction of the pupil is due to a stimulation of the oculo-motor nerve-centres, and that the dilatation of the pupil, as death approaches, is due to a paralysis of the same. Indeed, it cannot well be otherwise; for if the primary contraction were due to paralysis of the sympathetic, the secondary wide dilatation would be impossible; the dilating force—*i.e.*, the sympathetic—having been withdrawn, the pupil would not widely expand even if the contracting force—*i.e.*, the oculo-motor—were paralyzed.

In birds (Dr. S. Weir Mitchell, *loc. cit.*) the pupil is not affected; in horses it is widely dilated (Dr. Harley, *loc. cit.*); and in dogs it dilates before contracting (Dr. Reese, *loc. cit.*, apparently confirmed by Experiment number eight, Harley, *loc. cit.*, p. 109), or sometimes remains unchanged (Harley, *loc. cit.*, p. 111). At present these anomalies cannot be explained.

Morphia is probably all eliminated by the kidneys; it has been found in the urine of animals poisoned with it by Dr. Hilger (Gscheidlen, *loc. cit.*, p. 32), and Bouchardat (*Schmidt's Jahrbücher*, Bd. cxx.) has detected it in the same excretion from a person who had taken only three-quarters of a grain of the aqueous extract of opium.

The amount of the urinary secretion is said to be sometimes increased by morphia, but more generally it is diminished. Retention of urine after a full dose of opium is not rare, and without doubt depends upon the blunting of the sensibility of the bladder.

Upon the digestive tract opium exerts a very marked influence, checking secretion and causing constipation, acting in these respects more efficiently than does morphia.

THERAPEUTICS.—The chief indications for the use of opium are considered below, *seriatim*. Nearly all of them flow evidently from the known physiological action of the drug; others, however, although established by clinical experience, and undeniable, are not so plain in their philosophy.

1. *To relieve pain.* As an analgesic, opium is without a rival in the *materia medica*, except it be the anaesthetics. It is used to allay pain arising from any cause whatever, except acute inflammation of the brain, and is preferred to the anaesthetics whenever the pain has any permanency. In *painful spasm* it is especially useful, as it seems very frequently to quiet the motor as well as the sensory disturbance.

2. *To produce sleep.* Sleeplessness occurring in acute disease, and not dependent upon cerebral inflammation, may very frequently be relieved by opium. While it is often necessary to use the drug freely, care should be exercised not to overwhelm the nerve-centres, in such affections as *delirium tremens*, by enormous doses. In habitual sleeplessness great caution must be used in the employment of opium, not so much on account of the disturbance of digestion which it is liable to cause, as for fear of producing the "opium habit." Chloral is perhaps a more generally applicable hypnotic than

opium. Be this, however, as it may, I have found the combination of morphia and chloral singularly efficient. In low fevers, adynamic delirium often coexists with sleeplessness, and is then best met by opium.

3. *To allay irritation.* In various forms of nervous crethism, opium is most valuable; but when the affection is at all chronic, the dangers of the opium habit should not be lost sight of. On the other hand, in acute cases, as in the excitement which so frequently attends *hæmoptysis*, the drug should be used freely.

In many cases of disease, opium is serviceable by sustaining the system against an irritation for the time being irremediable, by blunting the sensibilities. In this way it is useful in the advanced stages of *smallpox*, and in various surgical affections, in which it also does good by allaying pain. In various local irritations, opium is continually employed, as in the *colic* caused by undigested food; in *bronchitis*, it is thus used to quiet cough.

By allaying irritation and pain, opium affords relief in most cases of inflammation; but in certain varieties of the affection it seems to do much more than this, exerting, in some way at present difficult to explain, a life-saving influence. In *peritonitis*, after due depletion, or in cases not requiring depletion, it should always be exhibited in large doses at regular intervals, in such a way as to keep the patient in a state of decided narcotism.

In severe *acute vomiting*, opium is one of the most reliable remedies. It is best used in the form of suppositories. Although, by checking secretion and peristaltis, opium usually causes constipation, yet when *obstruction* of the *bowels* is produced by spasm due to an irritation or inflammation, by relieving the latter the drug will sometimes act as a most efficient laxative.

4. *To check excessive secretion.* For this purpose opium is very largely employed in *diarrhæas*, and is very efficient either alone or in combination with various remedies. In *enteritis* and in *dysentery*, although no less frequently used than in diarrhæa, it is of service rather as an antiphlogistic and analgesic than by checking secretion. In *diabetes insipidus*, the combination of it and gallic acid is, I think, the most generally successful remedy.

In true saccharine *diabetes*, opium is of very great value in many cases, often ameliorating the symptoms, and, in conjunction with restricted diet, sometimes even effecting a cure. Of course, however, like all other known remedies in this disease, it most frequently acts simply as a palliative. According to Dr. Pavy (*Medical Times and Gazette*, June, 1869, p. 641), it affects the quantity of the urine before diminishing the sugar in it.

In severe *mercurial ptyalism*, opium often seems to check the discharge, but certainly is not nearly so powerful in this regard as atropia.

5. *To support the system.* Opium appears in low fevers, and in various protracted adynamic illnesses, to afford actual support to the system in some way not as yet made out. This is especially the case when, from any reason, sufficient food to keep up life cannot be taken or retained.

6. *As a sudorific.* In the form of Dover's powder, opium is very largely used

when it is desired to produce sweating, as in the early stages of a "general cold," or other forms of *muscular rheumatism*. With its use should generally be conjoined such measures as "soaking the feet," covering warm in bed, and the free drinking of hot lemonade or hot teas.

TOXICOLOGY.—Sufficient has already been said concerning the course and symptoms of poisoning by opium in ordinary cases. Sometimes in adults trismus and other convulsive manifestations are added to, or in a measure replace, the usual phenomena, and in children the drug appears at times to overpower the nerve-centres at once, so that the second stage is very much shortened or aborted, and symptoms of collapse, with unconsciousness, are developed very rapidly.

The positive diagnosis of opium-poisoning from the symptoms alone is often impossible.* In some cases of congestion of the brain, or of apoplexy, or of uræmia, the phenomena are identical with those sometimes seen in opium-poisoning. I have thought that inequality of the pupils is proof that a case is not one of narcotism; but Prof. Taylor has recorded an instance of opium-poisoning in which it occurred (*Manual of Medical Jurisprudence*, 7th Am. ed., 1873, p. 205).

The indications in the treatment of poisoning by opium are: first, to evacuate the stomach; second, to maintain respiration; third, to keep up the circulation when failing. The first indication may be met in three different ways: by an emetic, by the stomach-pump, and by Dieulafoy's aspirator. There is often in narcotic poisoning great difficulty in getting an emetic to act, owing to the obtunding of the sensibility of the nervous system by the drug. For this and other reasons, so palpable as not to need mentioning, a prompt stimulant emetic should be used; antimony, on account of its depressing influence, should always be avoided. *Mustard flour* is almost always to be had at once, and is very efficient. A heaped tablespoonful stirred up in a tumblerful of warm water should be exhibited as soon as possible, and, if it fail to act in fifteen minutes, should be repeated; then a powder of thirty grains each of sulphate of zinc and ipecacuanha may be given, to be repeated once or twice, at intervals of fifteen or twenty minutes. Large draughts of warm water should be administered in the intervals, and also between the acts of vomiting, so as thoroughly to wash out the stomach.

The stomach-pump is of no value when the solid drug has been ingested, but, if at hand, is preferable to emetics when a fluid preparation has been taken, because of the promptness of its results.

To maintain respiration is the ultimate object of all the measures which are commonly undertaken for the purpose of arousing the system in opium-poisoning. Unconsciousness in itself is of no moment, but as it deepens the sensibility of the respiratory centres grows less, and consequently the involuntary

* See Clinical Lecture by the author on the Diagnosis of Apoplexy, *Philadelphia Medical Times*, vol. iii.; also paper by Dr. Wilks, *Medical Times and Gazette*, 1863.

breathing is less rapidly or perfectly performed. More than this, when at all awake, a patient suffering from opium-poisoning can be made to supplement the almost suspended automatic breathing by voluntary respiration, and every effort to induce him to do this should be used. It is often surprising how an apparently unconscious man can be made to breathe by a command shouted in his ear. To keep a patient awake, walking, flagellations with small, *fine* twigs, shaking, shouting, and various other methods which may suggest themselves, should be practiced. Care should always be exercised not to carry these useful measures unnecessarily far, and perhaps add physical exhaustion to the natural prostration of the third stage.

The cold douche affords an excellent method of arousing the patient and at the same time of especially stimulating respiration. The simplest method of application is to support the head and shoulders of a patient stripped to the waist over a common wash-tub, and to dash the water over the chest and head. The effect is much greater if ice-cold water and water a little hotter than the hand will bear (115° F.) be used in quick succession. In the way of drugs, atropia is the only known stimulant to the respiratory centres, and as such is of the greatest value in opium-poisoning when there is evident failure of respiration. (For discussion of its use, see the article on Atropia.)

When other measures fail, artificial respiration should be resorted to, and steadily maintained until all cardiac action has ceased or recovery taken place. Whenever life is in evident danger from the insufficiency of natural respiration, the use of artificial respiration should not be postponed, but it should be practiced to supplement nature. Life has undoubtedly been saved in this way.

In regard to the amount of opium which will cause death, the smallest fatal dose on record is half a grain of morphia in the adult.* According to Dr. A. Calkins (*Quarterly Journal of Psychological Medicine*,† 1868, vol. ii. p. 739), four grains‡ of crude opium placed in the ear have caused death; also four grains by the mouth in more than one case.

According to the authority just quoted, out of twenty-nine reported cases in which a fluidounce of laudanum was taken, nine died. The maximum doses from which recovery has taken place without emesis are fifty-five grains of the solid opium and six ounces of laudanum. In a babe a day old, one minim of laudanum (E. Smith, *London Lancet*, 1854), and in one aged nine months, a few drops of paregoric (Wood, *Boston Medical and Surgical Journal*, 1858), have proved fatal.

ADMINISTRATION.—Whenever it is desired to produce very decided narcotism by the use of repeated doses of opium, the drug should *always be*

* Two cases (hypodermically used), *Medico-Chirurg. Trans.*, vol. i. One case (enema), Calkins.

† Dr. Calkins's very elaborate paper may be consulted.

‡ Taken from *Journal de Chimie*, 1831. Assuredly there must have been some mistake in this case.

given in liquid form, since opium pills sometimes become very hard and undergo solution so slowly that they may accumulate in the alimentary canal. On the other hand, in diarrhoeas, or in sickness of the stomach, old opium pills are thought by some to act better than more soluble forms of the drug.

Many persons cannot take opium on account of the very great secondary nausea and depression which it produces. It has been supposed that these disagreeable after-effects are due to the narcotina in opium; but this can hardly be, seeing that they often follow the use of the pure alkaloid, morphia. The deodorized tincture of opium agrees with some individuals better than any other preparation of the drug; and, as first pointed out by Dr. Da Costa, by giving a drachm of the bromide of potassium with twenty-five drops of it, the after-effects of the narcotic are often entirely avoided. In many neuralgic women the knowledge of this fact is an inestimable boon; in others, however, the unpleasant symptoms are not averted by the bromide.

Children always bear opium very badly, and to them only the weaker liquid preparations should be given. Dover's powder should especially be avoided. It is probable that in its manufacture on the large scale the ingredients are sometimes not thoroughly mixed: at least I have seen cases in which the symptoms caused by it were seemingly so out of proportion to the dose as to suggest that more than the officinal amount of opium was present.

In *acute vomiting* from any cause, in *dysentery*, in *strangury*, and other irritations of the urino-genital organs, great advantage is often to be gained from the use of opium by the rectum. Suppositories made out of the extract (gr. ss to i), or enemata of laudanum (gtt. xxx to xl), may be used in these cases. The latter should be made by adding the narcotic to a tablespoonful of starch-water.

The dose of opium for an adult is from one to three grains; for a child a year old, one-twenty-fourth of a grain.

Confectio Opii.—*Confection of Opium* contains only one grain of the drug in thirty-six of the mass.

Pilula Opii.—*Pills of Opium* are made with soap, and contain each one grain of the drug.

Pilula Saponis Composita.—*Compound Pill of Soap* is in a pilular mass, containing in every five grains one of opium.

Extractum Opii.—*Watery Extract of Opium* is twice the strength of the crude drug.

Tinctura Opii Camphorata.—*Purgative* has in every fluidounce one grain of opium, besides benzoic acid, oil of anise, and camphor, and, in consequence of the last ingredient, is more constipating than the other preparations of opium, and hence is preferred in diarrhoea-mixtures. It is also much used in cough-mixtures. Dose, fʒi to fʒi.

Tinctura Opii—Laudanum.—*Tincture of Opium* (grs. xxxvijss to fʒi). It should never be forgotten that one drop of the tincture of opium equals only half a minim. Dose, ℥xii to ℥xiv.

Tinctura Opii Deodorata—*Deodorized Tincture of Opium* (grs. xxxvii to f℥i). *Deodorized Laudanum* really consists of a watery extract of opium, with the odorous principle extracted by ether, and enough alcohol added to preserve the preparation. Dose, ℥xii to ℥xv.

Tinctura Opii Acetata—*Acetated Tincture of Opium* (grs. xlvi to f℥i). Dose, ℥viii to ℥x.

Vinum Opii—*Wine of Opium*—*Sydenham's Laudanum* (℥i to f℥i). Dose, ℥vi to ℥viii.

Acetum Opii—*Vinegar of Opium*—*Black Drop* (grs. lxxv to f℥i). Dose, ℥v to ℥vii.

MORPHIA.

This alkaloid occurs in minute, colorless, shining crystals, according to Guy melting at 330° F. and subliming at 340° F.; insoluble in cold and almost so in boiling water; only slightly soluble in cold alcohol and ether; freely soluble in boiling alcohol and in the fixed and volatile oils.

The following are some of the most sensitive and characteristic tests. In a solution of the alkaloid in concentrated sulphuric acid, which has been allowed to stand from ten to twelve hours, or has been heated for half an hour to 100° C., or momentarily to 150° C. and allowed to cool, the faint reddish violet changes at the point of contact to a deep-blue violet upon the addition of dilute nitric acid or of a crystal of saltpetre. Morphia, with concentrated sulphuric acid, makes a colorless solution, which, on strong heating, becomes red, violet, dirty green. With concentrated nitric acid it makes a red color, and finally a yellowish solution. With the neutral chloride of iron, morphia strikes a blue color, perceptible only when the test contains one part of the alkaloid in six hundred. Less characteristic, but much more sensitive, is the iodine test, with which, according to Husemann, one-tenthousandth part of morphia can be recognized. Iodic acid, in the form of a mixture of iodate of potash and sulphuric acid, is to be added to the suspected solution. If morphia be present, iodine will be set free, and can be recognized by the starch test.

The *Acetate* (MORPHIÆ ACETAS), *Sulphate* (MORPHIÆ SULPHAS), and *Muriate of Morphia* (MORPHIÆ MURIAS) are all of them official. The first is a white powder; the last two occur in snow-white feathery crystals. They are all soluble in water, and are of a bitter taste.

THERAPEUTICS.—The salts of morphia differ in their therapeutic value from opium chiefly in that they act with less power as sudorifics and in checking secretion in the bowels, and consequently constipating.

The smallness of their dose and their perfect solubility fit them for hypodermic use. Almost the only purpose for which they are used in this way is to relieve pain. The advantages of the method are the quickness of the results and the increased power of relieving suffering which the remedy seems to acquire. In cases of severe pain, hypodermics are invaluable; but it must be borne in mind that sometimes they cause most unpleasant symp-

toms. I have seen most alarming results from the injection of a sixth of a grain, and half a grain has produced death. In females, unless very robust, the maximum dose should be the eighth of a grain; in men, the sixth to the quarter. The dose of a salt of morphia, corresponding to a grain of opium, is one-quarter of a grain.

Liquor Morphice Sulphatis—*Solution of Morphia* (gr. i to fʒi). Dose. one to three teaspoonfuls.

Magendie's Solution of Morphia contains sixteen grains to the fluidounce: it is not official, and should not be kept in the apothecaries' shops.

NARCEIN.

This alkaloid, which is not official, was discovered by Pelletier in 1832. "It crystallizes out of its watery, alcoholic, and dilute acid solutions in long, white, four-sided, rhombic prisms, or in bunched masses of fine acicular crystals, odorless, and of a taste at first bitterish, but later styptic." (Pelletier, Hesse, Winckler—Husemann, *Die Pflanzenstoffe*.) According to Pelletier, it is soluble in 375 parts of water at 13° C.; according to Hesse, in 1285 parts at 12° C.; whilst Dr. S. Weir Mitchell found that a specimen prepared by Merck dissolved in 1000 parts, one prepared by Powers & Weightman in 4000 parts, and one of unknown European manufacture in 2100 parts, of distilled water at 60° F. Its saturated solution in boiling water on cooling is filled with crystals.

Concentrated nitric acid dissolves narcein with a yellow color, and the solution on being heated gives off reddish fumes; iodine makes with it a bluish-black mass, which forms a colorless solution in boiling water, but on cooling separates; with concentrated sulphuric acid narcein strikes a brown color, and finally makes a clear yellow solution (Husemann).

PHYSIOLOGICAL ACTION.—According to Baxt (*Reichert's Archiv*, 1869, p. 126), three or four centigrammes (0.46 to 0.62 gr.) of narcein, when injected into a frog, produce, in from ten to fifteen minutes, a semi-comatose condition, in which the batrachian makes no resistance or effort when laid upon his back or in other unnatural position. The respiration and circulation are not disturbed. The frog can be aroused by strong irritation, and when awake seems perfectly conscious. In three to six hours he comes out of his lethargic condition apparently unaffected. Albers (*Virchow's Archiv*, vol. xxvi., 1863), using larger doses, found that one grain produces in the frog sleep, reflex and also spontaneous convulsions, and, after seventy-four hours, death.

Dr. S. Weir Mitchell (*American Journal of the Medical Sciences*, Jan. 1870) has found nine grains of the alkaloid to have very little effect upon pigeons, causing only a little abnormal quietness.

In Baxt's experiments upon rabbits and guinea-pigs, fifteen centigrammes (2.3 grs.) had no perceptible influence; and on dogs Dr. S. Kersch (*Schmidt's Jahrbücher*, Bd. cxli. p. 15), and also Dr. Harley (*The Old Vegetable*

Neurotics, p. 143), found moderate doses (26 mgr. Kersch) equally inert. In a mouse (Harley) one-half grain caused tranquil sleep, with, after a time, tremors, from which the animal recovered, to be taken suddenly, some hours later, with fatal convulsions. At the post-mortem the tubules of the kidneys were found completely choked up with the alkaloid, which had crystallized in them and produced a mechanical suppression of urine, to which death was probably due. Schroff (quoted by Rabuteau), Frommüller (*Schmidt's Jahrbücher*, Bd. cxli. p. 15), Harley, Mitchell, and Da Costa (*Pennsylvania Hospital Reports*, 1868), have found narcein to act very feebly upon man. Harley gave one grain hypodermically, Da Costa two and one-half grains by the mouth, and Mitchell took five grains himself, with the result of only causing some headache; and Frommüller has exhibited as much as twenty grains with equally negative results.

These investigations are in close concord, and conclusive; and that of Dr. Da Costa upon man was very full and exhaustive.

Opposed to these results are, however, a number of seemingly equally conclusive investigations made by various French and German authorities. Much of the interest that has been manifested in the alkaloid arose from the assertion of Claude Bernard, in 1864 (*Archives Générales*, 6e sér., t. iv. p. 459), that it is the most pleasant and certain hypnotic of any of the opium alkaloids. He experimented upon dogs, guinea-pigs, rats, pigeons, sparrows, and frogs, and in all of them there was produced deep sleep, closely resembling natural sleep, with benumbed, but not destroyed, sensibility. The irritability of the morphia-sleep was wanting, and no secondary depression followed. Seven to eight centigrammes (1.65 grs.) was sufficient to throw a dog into the profoundest slumber. As early as 1852, Lecomte (Husemann, *Die Pflanzenstoffe*, p. 176) had affirmed that 0.1 gm. (1.53 gr.) of narcein thrown into the jugular vein of a dog would produce deep sleep, and very recently Rabuteau has confirmed the results of Bernard. Further, Behier (*Bulletin Thérap.*, t. lxxvii.), Debout (*Ibid.*), A. Eulenburg (*Schmidt's Jahrbücher*, Aug. and Oct. 1866), Liné (*Journal de Pharmacie et de Chimie*, 4e sér., t. iii.), also assert as the result of experience that narcein, in doses of one-half grain to a grain, produces in man pleasant, persistent sleep. Rabuteau (*loc. cit.*) also agrees with this, except in placing the dose somewhat higher.—viz., at from ten to twenty centigrammes (1.53 to 3.06 grs.). Oettinger (*Das Narcein als Arzneimittel*, Diss. Inaug., Tübingen, 1866) also asserts that decidedly larger doses of narcein than of morphia are required to obtain any action.

As seemingly these opposing results are all true, the only possible explanation is that different substances were used by the different sets of observers under the one name.

As the greatest care was practiced by Harley and Mitchell in obtaining the alkaloid pure, and as Claude Bernard states expressly that the substance used by him was soluble in twenty parts of water, it is most probable that the former observers really had, and that Bernard did not have, the narcein of Pelletier.

It is, at any rate, a very plain deduction from the above facts, that if it be so difficult—nay, impossible—to obtain in commerce a uniform reliable narcotin, it is not proper to use it as a medicine.

CODEIA.

Albers found that one grain of the muriate of codeia would produce in the frog, twenty minutes after its injection, tetanic cramps, alternating with convulsions; evidently reflex, since the slightest touch would call them forth. After a time the convulsive excitement grew less and less; the fore feet lost their sensitiveness first, but finally a paroxysm could no longer be provoked by touching the hind feet. The pupils at this time were widely dilated. Death occurred by failure of respiration, the heart continuing to beat a quarter of an hour after the extinguishment of all other movements.

In Woldemar Baxt's (*Reichert's Archiv*, 1869, p. 125) experiments, three centigrammes (0.45 gr.) produced in the frog deep sleep, lasting three or four hours. On awaking, the frog seemed more sensitive than natural to external irritation. Six centigrammes produced sleep more quickly, and following the deep sleep a stage of excessive sensitiveness, during which external irritation produced repeated cramp-like contractions. M. Wachs (quoted by Husemann, *Die Pflanzenstoffe*, p. 163) observed phenomena similar to those detailed by Albers, but noted a peculiarity of gait in the frog preceding the convulsive stage, owing apparently to disturbance in the innervation of the adductors.

Wachs also found that .010 gramme (0.15 gr.) produced in the pigeon only increased rate of respiration and sleepiness, whilst larger fatal doses caused restlessness, inability to stand, movements backward and in a circle, disturbance of respiration with gasping, cramps of single muscles, and finally convulsions, mostly clonic, frequently repeated, and followed by an adynamic condition, terminated by sudden death. In Dr. S. Weir Mitchell's experiments (*American Journal of the Medical Sciences*, Jan. 1870, p. 26), seven grains produced violent non-tetanic convulsions, ending fatally in one minute; and one grain caused similar symptoms, terminating fatally in eight minutes.

According to Husemann *loc. cit.*, p. 133, Kunkel, in 1833, from his experiments arrived at results similar to the recent ones of Bernard. The latter observer found that five centigrammes (.075 gr.) produced in dogs a sleep similar to, but not so profound as, that of morphia, with less benumbing of sensation, and not followed by any symptoms of depression.

These results have been called in question by Harley, and it seems most probable that Claude Bernard used something more than codeine in his experiments. Certainly Harley found that one to two grains of the alkaloid produced in the dog disturbance of respiration, languor, and convulsive twitchings, but no sleep. Moreover, Husemann states that, after fatal doses, Wachs observed similar symptoms in the dog and in the rabbit, as follows: falling of the head, trembling, spasmodic movements of the eyes and lips, rarely trismus,

movements in a circle and backwards, weakness of the legs, with hurried respiration and prominence of the eyeballs as prodromes; later in the poisoning there were severe convulsions, after one or many paroxysms of which great weakness developed, ending in death.

As a hypnotic in man, codeia has been used by Magendie, Berthé, Aran, Krebel, Reissner, Robiquet, and others, some of whom assert that the sleep produced by it is followed by nausea or other symptoms of depression, whilst others deny this. It does not seem necessary to discuss in detail the researches of the authorities alluded to, but it may be well to give the results of one or two of these observers more in full.

Robiquet in a series of experiments found that doses of 0.01 to 0.03 gramme (.15 to .46 gr.) produced a feeling of contentment, calmed nervousness, and induced refreshing sleep, whilst 0.1 to 0.2 gramme (1.53 to 3.07 gr.) caused deep sleep, followed by nausea and vomiting; 0.1 gramme (1.53 gr.) caused in children very alarming symptoms.

Harley has found in a number of experiments that, given by the mouth, codeia is a very uncertain and feeble hypnotic, four grains producing simply accelerated pulse, contracted pupils, and some giddiness, followed by nausea and vomiting. When the drug is given subcutaneously, "sommolency," he states, "is a more prominent effect, but only occurs in certain individuals." His hypodermic dose is one to two grains.

On the other hand, Dnorzak and Heinrich found that codeia in doses of 0.1 gramme caused gastric uneasiness and pain, some salivation, nausea, heat, and feeling of weight in the head, some confusion of thought, very marked reduction of the pulse-rate, and very marked tremors affecting the whole body; and Mitchell took five grains of the alkaloid without inducing other effects than a rise of twenty per minute in the pulse rate, nausea, slight giddiness, and a sense of heaviness about the head.

It is very evident that these various investigators have not had the same principle; and, whilst it appears probable that codeia is a feeble hypnotic, there seem to be as yet no grounds for believing it in any way superior to morphia, whilst the uncertainty as to the constitution of any specimen makes its clinical use extremely unsatisfactory.

The remaining active principles of opium being objects of physiological rather than of clinical interest, I shall allude only to their physiological actions.

Narcotina.—Although in Dr. Albers's experiment (*loc. cit.*, p. 244) one grain of narcotina proved fatal to a frog without the production of convulsions, yet the united testimony of Claude Bernard (*loc. cit.*, p. 462), of Baxt (*loc. cit.*, p. 124), of Rabuteau (*loc. cit.*, p. 266), and of Mitchell (*loc. cit.*, p. 23), would seem to show that in the lower animals this alkaloid produces, when given in sufficiently large dose, active spinal convulsions. In small dose (one-half grain) it caused, according to Albers and to Baxt, in frogs

a semi-comatose state. In larger dose (1.2 grains) the last observer found it to induce in the same batrachian very decided convulsions, similar to those of morphia-poisoning. Dr. Mitchell states that from two to three grains produce violent and fatal convulsions in the pigeon. Claude Bernard ranks narcotina above morphia and codeia, next to papaverina, as a convulsant in the lower animals. Yet Orfila found thirty grains necessary to kill a dog, and Baxt has given about two grains to rabbits and to guinea-pigs without producing any symptoms.

Man is no more sensitive to narcotina than the lower animals, if indeed he be so sensitive. Twenty and thirty grains of it have frequently been taken without effect, and doses of one hundred and twenty grains are said to have been exhibited with no greater result.

Thebaia.—Since the days of Magendie, it has been known that thebaia produces in all the lower animals powerful convulsions, resembling those of strychnia. Orfila, Albers, Baxt, Claude Bernard, F. W. Müller (*Das Thebaia*, Diss. Inaug., Marburg, 1868), Harley, Fraser, Mitchell, Rabuteau, and other observers, all agree as to the very great similarity between the action of this alkaloid and that of strychnia. Albers found 0.03 to 0.045 gramme (.45 to .67 gr.) sufficient to throw a frog into violent tetanic spasms, and in Harley's experiments two grains proved fatal to a bitch. In man it would probably act in sufficiently large doses as upon animals. Rabuteau took ten centigrammes (1.5 grains), with the production only of a feeling of lightness in the head.

Papaverina.—A great deal of discrepancy exists among observers as to the physiological action of papaverina, obviously dependent upon variations in the purity of the specimens which they have used. Schroff and Hoffmann believe it to be inert in man, since the latter observer took about seven grains without any effect being induced.

Albers and Claude Bernard claim that on animals it is a convulsant resembling thebaia, whilst Baxt asserts that in frogs it produces profound sleep, with great slowing of the heart's beat, but without tetanus, and that it even acts as an antitetanic in the poisoning of codeia and of morphia. He also finds that rabbits and guinea-pigs bear enormous doses of it. Rabuteau says two to three centigrammes produce violent convulsions in the frog, but twenty-five centigrammes in the dog cause no symptoms; yet in Baxt's experiments four to ten centigrammes served to cause profound coma in the latter animal. These contradictory statements are simply irreconcilable, and no opinion can be arrived at until new researches, prefaced by a rigid chemical study of the alkaloid used, are entered upon.

Porphyrocin, according to Albers, acts upon the frog as the most powerful of all the convulsant opium alkaloids. Baxt (*loc. cit.*, p. 123) found it in doses of two to three milligrammes (.030 to .045 grain) to throw the frog into a semi-comatose condition, followed in fifteen to twenty minutes by

most violent convulsive excitement. In sparrows one milligramme (.015 gr.), and in pigeons a larger dose, produced violent tremors, lasting ten to fifteen minutes, and followed by a state of semi-coma. In guinea-pigs ten to twenty milligrammes (.15 to .30 gr.), and in rabbits one-fourth to one-third larger dose, caused violent tetanic cramps; larger doses, speedily fatal convulsions. According to Schroff, 1.5 grains of it are without influence upon man.

According to Harley, in dogs *Cryptopia* produces wild delirium, with dilated pupils, followed by tetanic spasms. In the mouse, small doses produce some delirious excitement, followed by somnolency; while large doses cause heavy sleep, and death from failure of the respiration. A grain and a half injected beneath the skin caused in some persons intense drowsiness; in others, very slight symptoms.

Meconia, according to Albers (*loc. cit.*, p. 248), produces in (.045 grm.) the frog mild tremors, lessening of sensation, and finally death. On the higher animals its poisonous action is not very great. Orfila injected 0.06 grm. (0.9 gr.) into the jugular vein of a dog, without effect. Harley gave two grains hypodermically to the same animal, with no further result than a little abnormal quietness. According to the same observers, subcutaneous injections of fourteen grains have no effect on the horse, but one-sixth of a grain produces very decided hypnotism in the mouse. On man, Dublane (*Pharm. Centralblatt*, 1832), Schroff (*Medicin. Jahrb.*, 1870), Frommüller (*Klinische Studien narcot. Arzneien*), and Harley have found meconin, when given by the mouth, inert in doses varying from one to eighteen grains. Harley, however, claims that in doses of one to two grains given hypodermically it acts upon man as a very excellent hypnotic; but Frommüller has injected nearly two grains, with entirely negative results.

In regard to the action of *Pseudomorphine* and *Opianine* we have little or no knowledge.

Sertürner, the discoverer of *Meconic Acid* (quoted by Albers), found it in his experiments to be actively poisonous; but Sömmering and Albers have come to an opposite conclusion. Large doses (one to two grains) do, however, affect frogs, although slowly, inducing a stupor-like condition, with convulsions.

CANNABIS INDICA—INDIAN HEMP.

CANNABIS AMERICANA—AMERICAN HEMP.

Under the respective names which head this article, the U. S. Pharmacopœia recognizes in its Materia Medica list the dried tops of the *Cannabis sativa*, or common hemp plant, as it grows in India and in our own country. These substances are official for the manufacture of their extracts.

EXTRACTUM CANNABIS INDICÆ—EXTRACT OF INDIAN HEMP.

The alcoholic extract of Indian hemp is a blackish, resinous extract, of a decided narcotic odor and a peculiar taste. In the East, hemp and its educts

are used as narcotic stimulants. *Ganjah* is the dried plant as sold in the bazaars of Calcutta for smoking. *Churros* is the resinous exudation with the epidermis, etc., scraped off the leaves. *Hashish* is an Arabian preparation of the drug. The effects of the hemp upon the system would suggest that its active principle is an alkaloid, but all attempts so far have failed to separate anything from the resin, which represents the activity of the plant, and is known as *cannabin*. It is best obtained by precipitating the saturated tincture with water containing an alkali. Besides *cannabin*, hemp contains also a trace of volatile oil.

PHYSIOLOGICAL ACTION.—When given in full doses, *cannabis Indica* produces a feeling of exhilaration, with a condition of reverie, and a train of mental and nervous phenomena which varies very much according to the temperament or idiosyncrasies of the subject, and very probably also, to some extent, according to the nature of his surroundings. The sensations are generally spoken of as very pleasurable; often beautiful visions float before the eyes, and a sense of ecstasy fills the whole being; sometimes the venereal appetites are greatly excited; sometimes loud laughter, constant giggling, and other indications of mirth are present. Some years since, in experimenting with the American extract, I took a very large dose, and in the essay upon the subject (*Proceedings of the American Philosophical Society*, vol. xi. p. 226, 1869) the result was described as follows:

“About half-past four P.M., September 23, I took most of the extract. No immediate symptoms were produced. About seven P.M. a professional call was requested, and, forgetting all about the hemp, I went out and saw my patient. Whilst writing the prescription, I became perfectly oblivious to surrounding objects, but went on writing, without any check to or deviation from the ordinary series of mental acts connected with the process, at least that I am aware of. When the recipe was finished, I suddenly recollected where I was, and, looking up, saw my patient sitting quietly before me. The conviction was irresistible that I had sat thus many minutes, perhaps hours, and directly the idea fastened itself that the hemp had commenced to act, and had thrown me into a trance-like state of considerable duration, during which I had been stupidly sitting before my wondering patient.

“I hastily arose and apologized for remaining so long, but was assured I had only been a very few minutes. About seven and a half P.M. I returned home. I was by this time quite excited, and the feeling of hilarity now rapidly increased. It was not a sensuous feeling, in the ordinary meaning of the term; it was not merely an intellectual excitation, it was a sort of *bien-être*,—the very opposite to *malaise*. It did not come from without; it was not connected with any passion or sense. It was simply a feeling of inner joyousness; the heart seemed buoyant beyond all trouble; the whole system felt as though all sense of fatigue were forever banished; the mind gladly ran riot, free constantly to leap from one idea to another, apparently unbound from its ordinary laws. I was disposed to laugh; to make comic gestures; one very

frequently recurrent fancy was to imitate with the arms the motions of a fiddler, and with the lips the tune he was supposed to be playing. There was nothing like wild delirium, nor any hallucinations that I remember. At no time had I any visions, or at least any that I can now call to mind; but a person who was with me at that time states that once I raised my head and exclaimed, 'Oh, the mountains! the mountains!' Whilst I was performing the various antics already alluded to, I knew very well I was acting exceedingly foolishly, but could not control myself.

"I think it was about eight o'clock when I began to have a feeling of numbness in my limbs, also a sense of general uneasiness and unrest, and a fear lest I had taken an overdose. I now constantly walked about the house, my skin to myself was warm, in fact my whole surface felt flushed; my mouth and throat were very dry; my legs put on a strange, foreign feeling, as though they were not a part of my body. I counted my pulse and found it one hundred and twenty, quite full and strong. A foreboding, an undefined, horrible fear, as of impending death, now commenced to creep over me; in haste I sent for medical aid. The curious sensations in my limbs increased. My legs felt as though they were waxen pillars beneath me. I remember feeling them with my hand and finding them, as I thought at least, very firm, the muscles all in a state of tonic contraction. About eight o'clock I began to have marked 'spells,'—periods when all connection seemed to be severed between the external world and myself. I might be said to have been unconscious during these times, in so far that I was oblivious to all external objects, but on coming out of one, it was not a blank, dreamless void upon which I looked back, a mere empty space, but rather a period of active but aimless life. I do not think there was any connected thought in them; they seemed simply wild reveries, without any binding cord,—each a mere chaos of disjointed ideas. The mind seemed freed from all its ordinary laws of association, so that it passed from idea to idea, as it were, perfectly at random.

"The duration of these spells to me was very great, although they really lasted but from a few seconds to a minute or two. Indeed, I now entirely lost my power of measuring time. Seconds seemed hours; minutes seemed days; hours seemed infinite. Still, I was perfectly conscious during the intermissions between the paroxysms. I would look at my watch, and then after an hour or two, as I thought, would look again and find that scarcely five minutes had elapsed. I would gaze at its face in deep disgust, the minute-hand seemingly motionless, as though graven in the face itself; the laggard second-hand moving slowly, so slowly. It appeared a hopeless task to watch during its whole infinite round of a minute, and always would I give up in despair before the sixty seconds had elapsed. Occasionally, when my mind was most lucid, there was in it a sort of duplex action in regard to the duration of time. I would think to myself, It has been so long since a certain event,—an hour, for example, since the doctor came; and then reason would say, No, it has been only a few minutes; your thoughts or feelings are caused by

the hemp. Nevertheless, I was not able to shake off this sense of the almost indefinite prolongation of time, even for a minute. The paroxysms already alluded to were not accompanied with muscular relaxation. About a quarter before nine o'clock, I was standing at the door, anxiously watching for the doctor, and when the spells would come on I would remain standing, leaning slightly, perhaps, against the doorway. After awhile I saw a man approaching, whom I took to be the doctor. The sounds of his steps told me he was walking very rapidly, and he was under a gas-lamp, not more than one-fourth of a square distant, yet he appeared a vast distance away, and a corresponding time approaching. This was the only occasion in which I noticed an exaggeration of distance; in the room it was not perceptible. My extremities now began to grow cold, and I went into the house. I do not remember further, until I was aroused by the doctor shaking or calling me. Then intellection seemed pretty good. I narrated what I had done and suffered, and told the doctor my opinion was that an emetic was indicated, both to remove any of the extract still remaining in my stomach, and also to arouse the nervous system. I further suggested our going into the office, as more suitable than the parlor, where we then were. There was at this time a very marked sense of numbness in my limbs, and what the doctor said was a hard pinch produced no pain. When I attempted to walk up-stairs my legs seemed as though their lower halves were made of lead. After this there were no new symptoms, only an intensifying of those already mentioned. The periods of unconsciousness became at once longer and more frequent, and during their absence intellection was more imperfect, although when thoroughly roused I thought I reasoned and judged clearly. The oppressive feeling of impending death became more intense. It was horrible. Each paroxysm would seem to have been the longest I had suffered; as I came out of it, a voice seemed constantly saying, 'You are getting worse; your paroxysms are growing longer and deeper; they will overmaster you; you will die.'

"A sense of personal antagonism between my will-power and myself, as affected by the drug, grew very strong. I felt as though my only chance was to struggle against these paroxysms: that I must constantly arouse myself by an effort of will, and that effort was made with infinite toil and pain. I felt as if some evil spirit had control of the whole of me, except the will-power, and was in determined conflict with that, the last citadel of my being. I have never experienced anything like the fearful sense of almost hopeless anguish and utter weariness which was upon me. Once or twice during a paroxysm I had what might be called night-mare sensations; I felt myself mounting upwards, expanding, dilating, dissolving into the wide confines of space, overwhelmed by a horrible, rending, unutterable despair. Then, with tremendous effort, I seemed to shake this off, and to start up with the shuddering thought, Next time you will not be able to throw this off, and what then! Under the influence of an emetic I vomited freely, without nausea, and without much relief. About midnight, at the suggestion of the doctors,

I went up-stairs to bed. My legs and feet seemed so heavy I could scarcely move them, and it was as much as I could do to walk with help. I have no recollection whatever of being undressed, but am told I went immediately to sleep. When I awoke, early in the morning, my mind was at first clear, but in a few minutes the paroxysms, similar to those of the evening, came on again, and recurred at more or less brief intervals until late in the afternoon. All of the day there was marked anæsthesia of the skin.

"At no time were there any aphrodisiac feelings produced. There was a marked increase of the urinary secretion. There were no after-effects, such as nausea, headache, or constipation of the bowels."

The sense of prolongation of time which I experienced was to me very remarkable, but is not uncommon in these cases. It is evidently due to the immense rapidity of the succession of ideas. The mind, without doubt, measures time by the duration of its own processes, and when an infinitude of ideas arise before it in the time usually occupied by a few, time becomes infinitely prolonged to the mind. It is a lifetime in the minute.

In addition to symptoms above noted, or sometimes even replacing them in a measure, are in some cases marked disturbances of motility. Convulsions have been noticed by Dr. Lawrie (*Stillé's Therapeutics*, vol. i. p. 772), and local spasms, with salaam convulsions, by Dr. F. H. Brown. According to Dr. O'Shaughnessy, the induction of catalepsy is not rare among the Hindoos.

A very common mental phenomenon, not yet mentioned, is a condition of double consciousness, a sense of having two existences, of being at the same time oneself and somebody else. Whatever may be the symptoms of the first stage, sooner or later, if the dose be sufficient, drowsiness comes on. Generally, before it is marked, partial anæsthesia, often with partial loss of strength, is manifested, especially in the lower limbs. The pupils are dilated, the pulse is quickened, and finally the subject falls into a heavy sleep, out of which he generally awakes hungry, without any of the wretched gastric sensations or the malaise felt after an opiate. Confusion of thought, however, may persist for some hours. Cannabis exerts no constipating influence upon the bowels, and appears to increase, rather than decrease, the excretion of the kidneys.

THERAPEUTICS.—Leaving out of sight the employment of the medicine by alienists, hemp has been used in this country chiefly for the *relief of pain*, but also to some extent as a *hypnotic*. As an analgesic, it is very much inferior to opium, but may be tried when the latter is for any reason contra-indicated. In full doses, in *neuralgic* pains, it certainly often gives relief. It has been very largely employed to induce euthanasia in the advanced stages of *phthisis*, and constitutes, it is said, a popular nostrum employed for that purpose.

In *tetanus*, Indian hemp has been used quite largely, and until within a short time was, after opium, one of the few known drugs of service. Dr. Roemer (*St. Louis Medical and Surgical Journal*, p. 363, 1873) has collected thirty-five cases, with twenty-one recoveries and fourteen deaths.

ADMINISTRATION.—The action of the preparations of Indian hemp is exceedingly variable, in some cases small doses producing alarming effects, in other instances the remedy seeming almost entirely powerless. There appears to be a very great difference in the susceptibility of persons to its influence; but this cannot explain the wide variance of the clinical results obtained by its use. A large proportion of the extracts upon the market must be inert. It is possible that the crude drug undergoes deterioration during its long sea-voyage from India; at least I have seen the extract carefully made from genuine India hemp, and offering all the physical characters of good extract, yet entirely inert in doses of many grains. I have recently tried three American extracts, all made by prominent manufacturers, and all of them, so far as could be told by inspection, excellent preparations. To some half-dozen persons these were given in increasing doses until each patient took fifteen grains in three hours, and not a perceptible symptom was induced. I have seen an eighth of a grain of an English extract produce in a susceptible woman decided intoxication. So far as my present knowledge goes, although the foreign extracts of hemp are often inefficient, yet they are much more reliable than the American. The only way of administering hemp with satisfaction at present is for the practitioner to try various samples until he gets an active one, and then, being supplied with this, and having learned its proper dose by clinical experiment, to depend solely upon it. Hemp is not a dangerous drug: even the largest doses of its active preparations, although causing most alarming symptoms, do not really compromise life. No case of acute poisoning by it terminating fatally has, that I am aware of, been reported. There is an officinal *tincture* (*Tinctura Cannabis*), every fluidrachm of which contains nearly three grains of the extract.

EXTRACTUM CANNABIS AMERICANÆ—*American Extract of Hemp*.—On account of the great unreliability of the foreign hemp, the question as to the activity of the plant when grown in more temperate zones is one of practical interest. Many years since, Prof. Christison experimentally determined that the hemp-plant of Edinburgh is medicinally powerless, and Prof. Procter found that this is also true in regard to specimens grown in Philadelphia. The summers of the hemp regions in Kentucky approach in heat those of the more temperate parts of India, in which the hashish is produced; and some years since, incited by this, I made some experiments, and found that the Kentucky plant yielded quite largely an extract active in doses of one-half a grain to a grain. Whether future experimentation will show that Kentucky hemp will steadily yield as my sample did, time alone can determine. Certainly the subject is worthy of investigation on a larger scale. In making these experiments, well-matured, large, coarse plants should be alone used. If female plants are selected, the seed should be ripened, or nearly so; if male plants are taken, the pollen should have been shed some time previously.

CLASS VII.—MYDRIATICS.

IN the present group are considered three medicines,—belladonna, hyoscyamus, and stramonium,—whose preparations, when given internally or applied locally to the eye, dilate the pupil, or, in other words, produce mydriasis, and whose action upon the system appears to be almost identical.

BELLADONNÆ FOLIUM—BELLADONNA LEAF.

BELLADONNÆ RADIX—BELLADONNA ROOT.

The leaves and root of *Atropa belladonna*, an herbaceous perennial, a native of Europe, but cultivated in this country, and attaining a height of some three feet. The oval, pointed, entire, smooth, unequal leaves are in pairs, on a short footstalk. The bell-shaped, axillary, pendent flowers are of a dull reddish color. The globular berry is about three-quarters of an inch in diameter, deep purple, with a violet-colored juice, and adherent, green calyx. The dried leaves have a faint narcotic odor, and a sweetish, sub-acrid, slightly nauseous taste. The dried cylindrical branched root is from one to several inches in diameter, much longer, fibrous, externally reddish brown, internally whitish, almost odorless, with a very feeble sweetish taste.

ATROPIA.

The sole active principle of belladonna, discovered by Mein in 1831 and independently by Geiger and Hesse in 1833, occurs in silky prismatic and acicular, often aggregated crystals, of a bitter, burning taste, without odor, soluble in three hundred parts of cold and fifty-eight of boiling water, forty of benzole, thirty of ether, three of chloroform, and eight of alcohol. It is most abundant in the root, and, according to M. Lefort (*L'Union Médicale*, Nov. 1871), in that of young plants. Cyanogen gas passed through its alcoholic solution makes a deep red (Hinterbeyer). The most reliable test is the physiological one,—*i.e.*, the production of mydriasis in a rabbit or a cat by the local application to the eye. It has been found in all the tissues of poisoned individuals, but always exists in greatest abundance, and is most easily demonstrated, in the urine.

PHYSIOLOGICAL ACTION.—When the smallest physiological dose of atropia is administered to man, the only symptom induced is dryness of the throat and mouth, and possibly some disorder of vision. When a little larger

amount is given, this dryness is more intense, and is associated with redness of the fauces, dilated pupils, disordered vision, and possibly diplopia. The pulse is sometimes at first rendered less frequent, but this decrease is very transient, and certainly in many cases cannot be demonstrated at all. Often from the first, certainly after a short time in all cases, the heart's beats, after a large dose of the alkaloid, become excessively rapid, the pulse rising to one hundred and twenty, or even one hundred and sixty; and in a little while a peculiar bright-red flush appears on the face and neck, and may spread over the whole body. As I have seen this, it lacks the punctations of the rash of scarlet fever, and is only in very severe cases followed by desquamation.

Early in the course of the symptoms of atropia-poisoning there is very often forcible expulsion of urine, and erections of the penis may occur; but afterwards there is very generally, Harley says always (*Old Vegetable Neurotics*, p. 207), retention of urine. With the symptoms above enumerated, intellection may remain perfect; but there are generally some lightness of head, giddiness, and confusion of thought, as well as a staggering gait and restlessness. Occasionally, even with doses which may be called medicinal, there are spectral illusions. Drowsiness is not a general or at all characteristic symptom: if present it is apparently always produced indirectly, as by the removal of some cause of previous wakefulness.

When a decidedly poisonous amount of belladonna or its alkaloid has been taken, all the symptoms already noted are intensified, and to them is added a peculiar talkative, wakeful delirium, in which the patient lives in a world of his own, engrossed by the spectres and visions which throng him, and completely oblivious to the surrounding realities. Thus, I have seen a patient remain for a long time stooping and holding fast to the bed-post, to which she talked in the most voluble manner, as though it were an intelligent living entity. Sometimes this delirium is wild, and the patient almost uncontrollably violent. After a time, sleep may come on, and on waking from this complete consciousness may be regained, or the symptoms may gradually subside.

Sometimes, after a very large dose has been taken, severe convulsions appear in a very short time, and persist, with or without furious maniacal delirium, until near death. Sooner or later, however, the delirium subsides into stupor, and the convulsions into paralysis; and when the dose has been enormous, and especially when the alkaloid itself has been taken, stupor, with great muscular relaxation, may occur very early. Lividity of the face, and evident imperfect aeration of the blood, are not seen in atropia-poisoning, except in the stage of most imminent peril. Death is preceded by marked failure both of the heart's action and of the respiratory forces. In most cases, I think, it is actually brought about by asphyxia.* Post-mortem examinations have shown in fatal cases congestion of the lungs, often with

* See a case reported by S. W. Gross, *American Journal of the Medical Sciences*, 1869, p. 401, as a striking instance.

ecchymoses, and a similar state of the membranes and even substance of the brain and cord. According to M. Lemaitre,* congestion of the retina is an almost characteristic lesion of atropia-poisoning.

Upon the lower animals, belladonna to a great extent acts as upon man, although its influence is much less powerful in them, and very much larger doses are required. Seeming differences of action are in most cases simply apparent, not real. Thus, in the dog, as in man, the pulse-rate is very greatly increased by atropia, whilst in the rabbit it is not. As will be shown hereafter, the rise of the pulse-rate in the former is largely due to paralysis of the par vagum. Now, atropia paralyzes the par vagum in the rabbit as much as in the dog, but in the rabbit pneumogastric paralysis, by section or otherwise, is never followed by a rise of the pulse-rate at all comparable to that seen under similar circumstances in the dog. Evidently the action of the drug is identical in the two cases, although the symptoms are different.

In their sensitiveness to atropia animals differ very much, and, as a general rule, herbivora are less susceptible than carnivora. Thus, the rabbit may be fed for days entirely upon belladonna-leaves without injury, and many grains of atropia are necessary to kill him. Birds—at least pigeons—I have found will often recover after the hypodermic injection of two grains of atropia, and three grains by the mouth did not prove fatal. A very curious and at present inexplicable fact, which I have repeatedly verified, is that the pupils in pigeons cannot be dilated by the constitutional use of belladonna.

The close study of the physiological action of atropia can only be made system by system, and I shall now consider the subject under such headings.

Action on the Circulatory System.—In the dog and in the rabbit small doses of atropia cause an increased frequency of the pulse, with rise of the arterial pressure. If, however, larger amounts of the alkaloid are used, and especially if the poison is thrown immediately into the venous circulation, there is an immediate fall of the blood-pressure, although the pulse increases as before. Bezold and Bloebaum (*Ueber die physiologischen Wirkungen des Schwefel-Atropins*, in *Untersuch. aus d. Physiolog. Laborator. in Würzburg*, Heft i.) state that the dose of atropia can be so graduated as to produce at first a temporary rise of the arterial pressure, followed in a short time by a fall.

In atropinized animals, as is stated both by Bezold and Bloebaum (*loc. cit.*, p. 33) and by Meuriot (*De la Méthode physiologique en Thérapeutique et de ses Applications à l'Etude de la Belladonne*, Paris, 1868, p. 73), and as I have frequently seen, section of the vagi is not followed by any increase of the heart's action, and galvanization of the nerve is incapable of influencing the viscus. It follows that atropia paralyzes either the trunk or peripheral filaments of the nerve, and, as Bezold and Bloebaum believe, most probably the latter. The observers just alluded to have found that if atropia be injected into the carotid so as to reach the pneumogastric centres before the

* Quoted by Tardieu, *Sur l'Empoisonnement*, Paris, 1867, p. 752.

periphery, there is an instantaneous fall in the rate of the heart's beat,—a proof that upon the cardiac inhibitory centres atropia acts as a direct stimulant, precisely as it does on the spinal cord. The reason this action is not more manifest under ordinary circumstances is the incapacity of the paralyzed vagus to transmit the central impulse. The fact accounts very beautifully for the primary brief slowing of the pulse stated to occur in some cases of atropia-poisoning. (See *Stillé's Therapeutics*, vol. i. p. 725. Mitchell, Keen, and Morehouse found it in about one-third of the cases after large hypodermic injections; Da Costa in a larger proportion, *Amer. Jour. Med. Sciences*, July, 1865; Miss Mary Putnam in some cases, *New York Medical Record*, 1873.)

When, by the division of the spinal cord and vagi, the heart is isolated from the nerve-centres and the vaso-motor nerves are paralyzed, Bezold and Bloebaum have found that atropia in minute, as well as in large, amount, lessens at once the arterial pressure. Botkin (*Virchow's Archiv*, xxiv., 1862) states that when atropia is applied to the heart of the frog it at once diminishes, and finally arrests, its action, which cannot be re-excited by galvanic or other stimuli; and Bezold and Bloebaum (*loc. cit.*, p. 48) have found that the same thing occurs when a mammal is suddenly killed by the injection of an overwhelming dose of the alkaloid into the jugular vein. From these facts it follows that upon *the heart itself atropia acts as a direct depressant poison*, but for this influence to be apparent a very large amount of the alkaloid must be used.

From what has already been said, it is evident that the increase of the pulse seen in atropia-poisoning is in a measure due to a paralysis of the cardiac inhibitory nerves; but that this is not the only cause is shown by the fact noticed by Lemattre (*Archives Générales*, August, 1865) and confirmed by my own experiments, that after section of the par vagum atropia still causes an increase in the rapidity of the heart's action. Further, Bezold and Bloebaum have found (*loc. cit.*, p. 54) that when the thoracic sympathetic is galvanized, even after large doses of atropia, the heart is very sensibly affected. From these facts it is obvious that atropia acts on the cardiac *accelerator nerve-centres*, or *possibly nerves*, as a stimulant, and, unless it be in lethal doses, *does not destroy the excitability of these nerves*.

The relation of belladonna to the vaso-motor nervous system is of such practical importance that it deserves the closest study. Brown-Séquard says positively that the drug has the power of exciting the muscular fibres of the arterioles; but, as he nowhere details the reasons for this opinion, I think little weight is to be attached to his statement.

In 1857, Wharton Jones, of England (*Medical Times and Gazette*, p. 28, 1857), announced the fact (confirmed by Lemattre, *loc. cit.* p. 52) that if atropia be dropped upon the web of a frog's foot, under the microscope the vessels can be seen to contract. Dr. Hayden (*Dublin Quarterly*, Aug. 1863) in repeating these experiments found that, if the skin were cut just above the foot, no contraction occurred, and concluded that the phenomenon was purely

reflex. Meuriot (*loc. cit.*, p. 39) has obtained in some cases, but not invariably, the results of the last observer. He has, however, discovered that if the nerves of the leg be divided, no contraction ever takes place. Atropia is an irritant, and it is evident that the contraction caused by its local application is simply, as Hayden believed, a reflex phenomenon, precisely similar to that which occurs on the application of any chemical or mechanical irritant. The experiments of Wharton Jones, upon which so much has been based, must, therefore, be eliminated from the evidence on the question. The problem can be solved only by studying the effects of the remedy administered so as to act on the part solely through the circulation. Meuriot, as the result of such studies on the frog's web, has concluded that there is in the first stages of the poisoning a very slight contraction of the vessels, amounting to nothing more than increase of their tonicity; never to any decided lessening of their lumen. Dr. Harley (*The Old Vegetable Neurotics*, London, 1869, p. 220) has witnessed a more decided contraction of the vessels. I have tried the experiment several times, but obtained no decided results. On the other hand, Bezold and Bloebaum (*loc. cit.*, p. 50) have made similar experiments upon the ear and mesentery of the rabbit, and have never been able to detect any contraction of the vessels, and consequently deny its existence. In their experiments it was not possible to use the microscope, which detracts greatly from their value.

The evidence derived from direct observation of the capillaries seems to me to be, on the whole, decidedly in favor of their contraction by minute doses of belladonna. I do not think, however, that much weight is to be attached to evidence of this nature. The alterations in the calibre of the vessels are so slight as to leave great play for the imagination of the observer,—a source of fallacy which probably accounts for the different results obtained by different investigators. Much more decisive proof is, however, obtainable from a study of the arterial pressure. I have found that after section of the vagi atropia still has the power of raising very materially the arterial pressure. As atropia does not augment the force of the individual cardiac beat, and as the increase in the number of the cardiac pulsations caused by it after section of the vagi is comparatively slight, it is exceedingly probable that the rise of arterial pressure just spoken of is due to a contraction of the small vessels. This logical conclusion becomes almost a certainty when it is further known that after division of the cord, and consequent separation of the vessels from their vaso-motor centres, atropia is powerless to produce rise of arterial pressure, a fact vouched for by Bezold and Bloebaum, and which I can confirm from my own experiments (*Amer. Jour. Med. Sci.*, April, 1873). To this cumulative evidence must be added the experimental fact noted by Bezold and Bloebaum, that when a small dose of atropia is injected into the carotid artery—*i.e.*, into the vaso-motor centres—there is an instantaneous rise of blood-pressure.*

* It is proper to state that Bezold and Bloebaum attribute this rise to psychical disturbance, without, however, as it seems to me, good reason.

Viewing all these facts together, I am forced to give assent to the proposition that *atropia*, in not too large amount, is a *stimulant to the vaso-motor centres*; a conclusion in harmony with the action of the drug on all the other motor centres.

All observers agree that in the advanced stage of atropia-poisoning, after the blood-pressure has commenced to fall, there is dilatation of the capillaries. The probabilities seem that this is due to a direct action of the poison on the muscular fibres in the coats of the vessels; for when directly applied to the web of the frog's foot, atropia, after a time, produces an evidently paralytic dilatation; and Bezold and Bloebaum have found that the arterial muscular coats in atropia-poisoning finally lose their irritability, but that so long as they retain it, galvanic stimulation of a sympathetic nerve does not fail to induce contraction in the tributary vessels.

Action on the Nervous System.—In 1862, Dr. S. Botkin (*Virchow's Archiv.* Bd. xxiv. p. 85) found that when the vessels of a frog's leg were tied and the animal poisoned with atropia, whilst paralysis developed itself in the ordinary way in the uninjured leg, the injured leg preserved its motility. He also discovered, on testing with galvanic currents, that the nerve of the leg whose artery had been tied transmitted a very forcible impulse to its tributary muscles under the stimulus of a galvanic current much weaker than could elicit the faintest response from the nerve of the opposite side. He concluded, very logically, that atropia acts as a paralyzant to the motor nerve-trunks themselves, and also, since motion persisted in the protected leg after it was completely lost in its fellows, that this influence of the drug was exerted upon the motor trunks before the spinal centres. These experiments have been confirmed by Lemattre (*loc. cit.*, p. 49), by Bezold and Bloebaum (*loc. cit.*, p. 20), by Meuriot (*loc. cit.*, p. 90), and by Fraser and others, who have proven that atropia, if in sufficient dose, has the power of destroying the excitability of the efferent or motor nerve-fibres, but that it must be in very large quantity, so that in mammals death may be caused by the alkaloid and yet a notable amount of functional power be retained by the motor nerves.

Bezold and Bloebaum, whose elaborate experiments are especially commendable, affirm that very rarely have they been able totally to destroy by atropia the functional power of the motor nerves, and also have shown that both the nerve-stem and the peripheral intra-muscular nerve-endings are affected.

All the experimenters agree that no stage of super-excitability preceding that of depression can be discovered.

The action of the alkaloid upon the spinal centres has only been closely and successfully studied by one observer, Dr. Fraser, of Edinburgh. He discovered* that if a frog receive a hypodermic injection of about one-thou-

* An Investigation into some Previously Undescribed Tetanic Symptoms produced by Atropia in Cold-Blooded Animals, by Thomas R. Fraser, M.D., *Transactions of the Royal Society of Edinburgh*, vol. xxv., 1869, p. 450.

sandth part of its weight of atropia, a condition of perfect paralysis and abolition of reflex actions comes on after a time and lasts from two to four days, to be succeeded by a tetanic stage, with violent convulsions and excessive excitability of the reflex centres. During the first or paralytic stage, electrical excitation of the motor nerves fails to induce muscular contraction; the function of the nerves is, in other words, completely suspended, and as it is regained, *pari passu*, the tetanic symptoms develop. This indicates very strongly that atropia is a spinal stimulant, and that the state of the centres during the first stage is simply masked by the inability of the nerve to transmit impulses. This conclusion appears to me absolutely proven by the following facts, which Dr. Fraser has experimentally determined. The convulsions are certainly spinal and not cerebral, since they occur after section of the cord high up in the neck. Further, if a sciatic nerve be protected from the action of the alkaloid by occlusion of the crural artery, tetanus occurs in the limb during the time when the rest of the body is completely paralyzed.

The experiments of Lemaitre (*loc. cit.*) indicate that belladonna exerts a similar double action in mammals, and Dr. Fraser's investigation confirms this. It would seem, however, that in mammals the spinal action comes on more rapidly, and is less protracted, and that the nerve-action is also less intense, than in the frog; so that whilst the batrachian poisoned by atropia is in the beginning paralyzed and afterwards convulsed, the mammal suffers from convulsions and reflex spasms early in the poisoning, and often exhibits evidences all through of the stimulant spinal action striving to make itself felt through the disabled nerves. The difference in the action of non-fatal doses of atropia upon the frog and upon the mammal is graphically expressed by the following diagrams, which are copied from the paper of Dr. Fraser.

FIG. 1.




FIG. 2.




Fig. 1 represents an experiment made upon a dog: in it the curve of paralysis, $op_1 p_2$, leaves the line of normality, AB, before the curve of spinal stimulation, the first symptoms being those of paralysis. As the curve of $op_1 p_2$ never reaches the line of complete paralysis, CD, the paralytic action is not sufficiently great, in the height of the poison, to prevent the manifestation of the considerable spinal stimulant action.

Fig. 2 expresses the results of an experiment upon a frog: in it, although there was some spinal stimulant action, as evinced by the curve s_4, s_8, s_{12}, s_{16} ,

and although the paralytic curve $p_4, p_8, p_{12}, p_{16}, p_{20}$ never reaches the line of complete paralysis, CD, yet it is manifest that the paralytic action of the drug throughout the experiment masks and covers up the convulsive influence.

It is very certain that in man atropia exerts this double influence; for the records of poisoning-cases are at once the records of convulsions and of paralysis. It would seem that early profound paralysis occurs when a very large dose of the poison has been taken in its concentrated alkaloidal condition, and consequently has been rapidly absorbed and suddenly precipitated upon the nervous system.

In the experiments of Botkin, made by tying the vessels of the frog's legs, as described on page 211, the investigator found that although for a time irritation of the foot of the perfectly paralyzed leg would cause spasms in the opposite limb whose nerve was protected, yet later in the poisoning, although irritation of the foot of the protected leg caused movements in that leg, no irritation of the opposite poisoned foot was able to induce any response whatever on either side. From this he drew the inevitable conclusion that, whilst atropia paralyzes first the motor nerves, yet later it acts also on the afferent nerves.

The very beautiful experiments of Dr. Fraser, already quoted, certainly show that the afferent nerves enjoy a comparative immunity from the influence of atropia in large doses. For when one leg of a frog was protected by tying the crural artery during the period of general paralysis, any irritation of the poisoned leg would produce immediate reflex convulsions in the unpoisoned extremity,—a demonstration that whilst the functions of the efferent or motor nerves were suspended wherever the poison reached, the afferent nerves retained more or less of their activity.

This of course confirms a portion of the experiments and conclusions of Botkin, and does not disprove that atropia acts to some extent upon the afferent nerves, for a very feeble impulse reaching the cord in its excited condition would cause reflex movements.

Bezold and Bloebaum (*loc. cit.*, p. 20) have investigated this subject by exposing the sciatic nerves of a strychnized frog, immersing them for a portion of their course, the one in a solution of atropia, the other in a solution of the phosphate of soda, and then testing the effect of the application of stimuli in causing reflex movements. The result was not very satisfactory, so many sources of fallacy arising—sources not necessary here to point out—as to vitiate greatly the experiments. They certainly found, however, that the nerve-trunk preserved for some time its power of transmitting impulse even when immersed in a two and a half per cent. solution of the alkaloid. Bezold and Bloebaum (p. 25) also repeated the experiments of Botkin, substantially confirming them, but showing that very large doses are necessary to affect seriously the conducting powers of the afferent nerves.

The following experiments of Meuriot (*loc. cit.*, p. 95) prove conclusively that atropia does influence the afferent nerves. He found that if a frog be

bound tightly around the body so as to interrupt the circulation, and then be poisoned by atropia in the front part of its body, at first irritations in any part give rise to general spasms, but after a time in order to get any movements of the hind legs it is necessary to apply an irritant to them. Again, the hinder parts of a frog were so bound by ligatures as to cut off on the one side all communication except by the nerves, and on the other to leave free the nerve and the vessels. A strong injection of atropia was then given, and when the moment came that irritation of the periphery of the leg whose circulation was free would no longer cause reflex spasms, the artery of this leg also was tied, so that both legs, the one atropinized, the other not, were now connected with the body of the frog only by their nerves. Strychnia was given hypodermically, and it was found that, whilst irritation of the atropinized leg had no effect, stimulation of the non-atropinized leg gave rise to general convulsions. These experiments show that atropia acts upon the afferent nerves as well as upon the efferent. Recently, Miss Mary Putnam (*New York Med. Record*, 1873) has asserted that if the general tissue of the frog's limb be tied, the vessels and nerves being left intact, so as to prevent any possible diffusion of the poison through the cellular tissue, the sensitive nerves are alone affected by atropia. As no details of experiments are given, the matter rests simply upon the assertion of Miss Putnam. It is very difficult to imagine why a nerve should be affected differently by a poison when reaching it by different routes, and equally difficult to conceive how the poison should reach more effectively the motor than the sensitive nerves by diffusion. Further, it is also almost as unreasonable to believe that any perceptible diffusion through the cellular tissue should occur when the seat of injection is in a distant part of the body. Taking together the facts that Miss Putnam's assertion is unsupported by detailed experiments, is exceedingly improbable, and is seemingly opposed to an enormous mass of experimentation by some of the most experienced and able investigators who have ever lived, it seems to me that it cannot be accepted. It is probable that atropia acts upon the peripheral filaments of the nerves more quickly than upon the main trunks; and this is corroborated by Miss Putnam, who states that she has seen galvanization of a sciatic nerve cause indications of pain in an atropinized animal when the peripheral sensation was already lost.

There can be no doubt that in the higher animals atropia acts upon the cerebrum as it does in man, producing the same peculiar delirious intoxication often ending in stupor. It is not necessary to discuss this matter further, except to remark that belladonna is *not* a hypnotic. The fact that the exhibition of a remedy is followed by sleep in disease is no proof that the remedy is in a proper sense a hypnotic. No one would give oyster soup such a cognomen, yet in certain exhausted wakeful states of the system such food may bring back sound sleep. If belladonna ever acts as a hypnotic, it is in this indirect way, by removing some cause of abnormal wakefulness. It can never be relied on, like opium, to induce sleep.

Action on the Muscular System.—The voluntary muscles escape unscathed in atropia-poisoning. It is true that Lemattre has shown that the contractility of a muscle may be destroyed by soaking it in a very concentrated solution of the alkaloid; but long before any such action can take place in life the animal is killed; consequently after death from belladonna the contractility of the muscles is found unimpaired.

On the non-striated muscles the action of the drug is much more pronounced, but the exact nature of such action is still uncertain. Meuriot states (*loc. cit.*, p. 112) that if the belly of an animal poisoned by atropia be opened, the intestines will be found undergoing violent contractions, and that belladonna is a powerful excitant of the non-striated muscles. On the other hand, Bezold and Bloebaum affirm that they have experimented upon the intestines, bladder, uterus, and ureters, and that in all cases there is a state of marked sedation from atropia, and that, whether a small or a large dose be used (*loc. cit.*, p. 65), there is produced muscular quietness and relaxation in all these organs,—evidences of sedation not preceded by any stage of excitement, and always accompanied by lessening of the electro-muscular sensibility. They state, further, that by the use of sufficient doses absolute muscular paralysis of the organs alluded to is induced, so that the strongest faradaic currents are unable to cause any motion.

P. Keuchel* has made a most elaborate series of experiments to determine the effect of atropia upon the inhibitory fibres of the splanchnic nerve. It has been shown by Pflüger that galvanization of the peripheral ends of the divided splanchnics causes immediate arrest of the peristaltic intestinal movements, and that the nerves mentioned are to these movements what the par vagum is to those of the heart. Keuchel has found that even when doses of atropia so small as not to affect distinctly the motor nerves are given, galvanization of the splanchnics fails to affect the intestines, and therefore that atropia paralyzes the peripheral inhibitory intestinal apparatus precisely as it does that of the heart.

There is, of course, seeming disagreement in the results obtained by Meuriot, by Bezold and Bloebaum, and by Keuchel; but it is possible they are all correct. It appears certain that, in full doses, atropia paralyzes the smooth muscular fibre of the intestine, bladder, etc., and it may be that in any dose it depresses their activity directly, but that, by paralyzing more quickly the inhibitory apparatus, it sometimes places the intestinal muscular coat in such a position that it will respond more vigorously than normal to external stimuli. If this be so, it is evident that there is real accord in the results of all these investigations.

Action on the Respiratory System.—It has already been stated that in man

* Unfortunately, I have had access only to an abstract (*Schmidt's Jahrbücher*, Bd. cxliii. p. 14) of Keuchel's paper, which is an inaugural dissertation, published in 1868 at Dorpat.

small doses of atropia do not affect the respiration, whereas large doses accelerate it. The same is true in animals; and the acceleration takes place even when the vagi have been previously divided,—a proof that it is due to a direct stimulation of the respiratory centres in the medulla. (Bezold and Bloebaum, *loc. cit.*, p. 59.) When *very large* fatal doses are administered, there is evidently also a paralysis of that portion of the peripheral pneumogastric nerve which is connected with the function of respiration; for if in an animal suffering only from a moderate dose the par vagum be cut, respiration is profoundly affected, but when the alkaloid has been more freely given, no marked influence is exerted upon the expiratory rhythm by section of the pneumogastrics. As death approaches in belladonna-poisoning, the blood, which has preserved its normal coloration, may become very dark, and the patient may at last die of asphyxia,—probably not from failure of the centres themselves, but from the loss of functional power in the respiratory nerves.

Action on the Glandular System.—One of the earliest and most notable effects of medicinal doses of atropia is dryness of the mouth, due to suppression of the secretions of the mucous and salivary glands. The action of the alkaloid upon the skin is similar. It can scarcely be doubted that this arrest of secretion is nervous, and the experiments of Keuchel (*loc. cit.*, p. 15) indicate that it is due to an action upon the peripheral nerve-filaments. As was first discovered by Schiff,* section of the chorda tympani is followed by arrest of secretion of the submaxillary gland, and galvanization of the peripheral end produces a greatly increased flow of saliva. In Keuchel's experiments these phenomena occurred in the unpoisoned animal, but when atropia had been exhibited galvanization of the peripheral chorda tympani was powerless to excite secretion,—proof that the peripheral end of the nerve was paralyzed.

Elimination.—When atropia is exhibited medicinally it probably all escapes from the body through the urine, and even after poisonous doses the amount eliminated through other channels must be almost infinitely small.

Probably on account of this elimination, certainly for some reason or other, after small doses of atropia the urine is increased,—sometimes, according to Harley, doubled. I am confident, however, that this increase varies very much, and is not always marked. After very large toxic doses the urine may be at first increased, but is usually lessened very early, and in the later stages may be entirely suppressed.† Meuriot states that the secretion of urine rises and falls in atropia-poisoning with the arterial pressure.

The experiments of Harley upon men (*loc. cit.*, p. 214) would appear to show that medicinal doses of atropia decidedly increase the solids of the urine, slightly the urea and uric acid, very markedly the phosphates and the sulphates.

In regard to the secretions of the alimentary canal, the action of atropia is

* See *Physiology of Man—Nervous System*, by Prof. A. Flint, Jr., New York, 1873, p. 158.

† See poison-case reported by Dr. Gross (*loc. cit.*), also cases of Dr. Morel (*Annales de la Société de Médecine de Gand*, 1873).

very uncertain. It has been a matter of traditional and clinical belief that they were increased, and Harley gives some experiments which he claims corroborate this; Meuriot, on the other hand, states that they are lessened. I cannot find, however, any experiments that seem to me at all decisive, and clinical evidence certainly indicates that the alimentary secretions, if affected at all, are increased.

Action on Temperature.—In moderate doses atropia causes a pronounced rise in temperature, but in very large decidedly toxic amounts it lessens animal heat. Thus, in the dog, Meuriot has obtained an augmentation of from 1° to 3° C., and Duméril, Demarquay, and Lecomte* of 4° C. In fatal poisoning of the same animal, these observers have noticed a fall respectively of $5\frac{1}{10}^{\circ}$ and of 3° .* In man, Meuriot, in the use of medicinal doses, has observed the temperature to rise $\frac{5}{10}^{\circ}$ to $1\frac{1}{10}^{\circ}$, and Eulenburg* $\frac{1}{2}^{\circ}$ to $\frac{8}{10}^{\circ}$.† Harley has seen in man an elevation of 1° F. As pointed out by Meuriot, the rise and fall of temperature probably correspond to the rise and fall of the blood-pressure.

From what has been already stated, it is evident that the action of atropia in therapeutic and in toxic doses is in a sense quite different.

In full medicinal doses it produces a sort of febrile state, with dryness of the mouth, increased rapidity and force of the circulation, quickened respiration, elevation of temperature, and secretion of febrile urine. The rapidity of the heart's action is due to paralysis of the peripheral inhibitory nerve and to stimulation of the accelerator nerves; the increased arterial pressure, to the increased cardiac action, together with the general contraction of the capillaries, the result of excitation of the vaso-motor centres. The spinal cord is not sensibly affected by these doses; the motor, and probably to a much less degree the sensory peripheral nerves, suffer lessening of functional activity, although the influence of therapeutic doses of atropia upon them must be very slight. If the dose be sufficiently large, the cerebrum is thrown into a condition of mild delirium, resembling also that of fever.

After decidedly toxic doses of atropia the blood-pressure falls, from dilatation of the capillaries, owing to the paralysis of their muscular coats, and from direct laming of the heart-muscle. The temperature also falls; the muscular system is relaxed, and sensation is impaired, from the paresis of the motor and sensory nerves respectively; yet convulsions may now occur from the over-activity of the reflex centres, the predominance of paralysis, or of convulsions, varying with the dose, as the depressing or stimulating influence is the most powerful. Delirium precedes stupor, which in turn precedes death, from asphyxia, caused by failing trunkal nerve functions, or very rarely from syncope, caused by failure of the cardiac muscle.

It is evident that when belladonna is applied to a part it must act locally as

* Quoted by Meuriot, p. 111.

† These figures are probably all of them Centigrade, though it is not stated by Meuriot.

a paralyzant, no doubt overpowering the capillary walls, the sensitive and motor nerves, and even muscular and glandular cell-action; for, except in the case of the latter, experimental evidence has already been brought forward to prove that, locally and freely applied, belladonna is a sedative poison, and clinical evidence points very strongly to its exerting a similar influence upon gland-cells.

Action on the Eye.—Atropia placed in the eye, or given internally, dilates the pupil of all animals except birds. Accompanying this mydriasis are paralysis of accommodation and a lessened intraocular pressure. In regard to the latter, the subject is so intricate that I must refer the reader to the treatises of specialists.*

Before discussing briefly the action of atropia upon the pupil, the fact that a recent American female writer has reasserted the old theory that the movements of the iris are due to erectile tissue, or, in other words, to its blood-vessels, seems to render necessary a few words as to the real motile power of the part. In the first place, it is an indisputable anatomical fact that the iris is largely composed of muscular fibres, and it is a simple common-sense deduction that the muscular fibres are there for the purpose of causing motion, especially since, in many animals, it can be readily demonstrated that whilst some of these fibres are circular, others are radiating, so that by position they become antagonistic. The paper of Dr. Arlt (*Archiv für Ophthalmologie*, 1869) seems to me very decisive. In a very elaborate series of experiments it was found that when the upper cervical ganglion was stimulated the pupil dilated long before any influence upon the vessels was detected, and that on cessation of the stimulation the pupil became natural long before the spasm of the vessels yielded; to my mind a proof that the ganglion has fibres other than vaso-motor,—fibres which control the muscular actions of the iris, and are more sensitive than the vaso-motor filaments; and, secondarily, a proof that the movements of the iris are not due to movements of the blood-vessels.

Space cannot be afforded in the present work for a further discussion of this subject. The reader is respectfully referred for further information to books on the physiology of the eye, and especially to Engelhardt's paper, *Beiträge zur Lehre von den Bewegungen der Iris* (*Untersuchungen aus dem Physiolog. Laboratorium in Würzburg, zweiter Theil*).

In considering the action of belladonna upon the eye, it is necessary to view separately its influence when applied locally and when given internally; and I shall consider these in the order in which they here stand.

It may be first asserted that the dilatation induced by the local application of belladonna or of its principles is a nervous phenomenon, and not due to a direct action of the drug upon the muscular fibres of the iris; for as all of these, both the radiating and the circular, are of the same nature (*non-striated* in mammals), their antagonism is simply due to position; and it seems inconceivable that mere position should affect the relations between a muscle

* See especially *Der intraoculare Druck und die Innervations-Verhältnisse der Iris*. Von Prof. Dr. Stellwag von Carion. Wien, 1868.

and a drug. Moreover, decisive proof is afforded by the experiments of Bernstein and Dogiel, who found that whilst galvanic irritation of the oculo-motor nerve was unable to cause contraction of the pupil in the atropinized eye, yet when the electrodes were applied to the eyes in such a way as to affect directly the iris contraction occurred,—phenomena only explainable by the theory that the nerve-endings were paralyzed, whilst the muscle was unaffected. Of the truth of this observation there can be no doubt, as it has been confirmed by Dr. Gustav Engelhardt (*Untersuchungen aus dem Physiolog. Laboratorium in Würzburg, zweiter Theil*, p. 321).

The statement first made by Wharton Jones (*Medical Times and Gazette*, 1857), that the reason atropia does not dilate the pupils of birds is that their irides have no radiating fibres, has been disproven by the beautiful anatomical researches of Alex. Ivanoff and Alex. Rollett (*Archiv für Ophthalmologie*, vol. xv. p. 1), confirmed by Johannes Diegel (*Max Schultze's Archiv für Mikroskop. Anat.*, Bd. vi. Heft i., 1870). As, therefore, there are radiating fibres in the eyes of birds, no reason can, as yet, be assigned for the non-action of atropia on their irides, unless it be that they are composed solely of striated muscular fibres. It is just possible, indeed, that the observation as to the non-action of the mydriatic upon the irides of birds is incorrect. Donders (*The Accommodation and Refraction of the Eye*, Syd. Soc. ed., p. 584) says that the pupillary action of atropia “is slight in birds in whom it was formerly overlooked.” In my own experiments the most thorough application of very strong solutions to the eyes of pigeons had no distinct effect. Unfortunately, however, accurate measurements of the pupils were not made.

The dilatation of the pupil by the local application of atropia is certainly independent of any nerve-centres farther back than the ciliary ganglion. This is proven by the following facts: Claude Bernard (*Physiologie et Pathologie du Système Nerveux*, Paris, vol. ii. p. 212) and Lemattre (*loc. cit.*) both have found that atropia-mydriasis occurs in animals after section of the oculo-motor, and I have seen it in cases of complete oculo-motor paralysis in man. It also takes place after section of the trigeminus or of the cervical sympathetic, or of both of these nerves, as is shown by the testimony of numerous observers and by my own experiments. In man, I have seen it after paralysis of the sympathetic (*Philadelphia Medical Times*, vol. i. p. 290).

Not only is the dilatation of the pupil by the local application of atropia independent of the central nervous system, but also of the ciliary ganglion, and it is therefore due to an action exerted directly upon the *nerve-endings in the iris*. The experiments of Bernstein and Dogiel, confirmed by Engelhardt, already quoted, are in themselves almost enough to establish the truth of this proposition. More direct evidence is not, however, wanting. Thus, Vierordt* has found that atropia locally applied still causes mydriasis after

* Unfortunately, the only notice I have seen of this capital experiment is in Hermann's “*Grundriss der Physiologie*.” No reference is given, and I have been unable to find the original paper.

the removal of the ciliary ganglion. Prof. I. Hoppe (*Die Nervenwirkungen der Heilmittel*, Leipzig, 1856, zweiter Heft, p. 179) has discovered, and Y. Valentin (*Versuch einer physiologischen Pathologie der Nerven*, Leipzig, 1864, zweite Abtheilung, p. 368) has confirmed, that in the eye of the frog removed from the body atropia will produce dilatation of the pupil. According to Borelli (*Edinburgh Medical Journal*, Nov. 1871), mydriasis is produced by the alkaloid when applied to the eye of a man just dead. Lastly, the presence of the alkaloid in the humors of the atropinized eye has been proven by numerous observers, among whom may be mentioned Lematre (*loc. cit.*, p. 55) and Prof. Donders (*The Accommodation and Refraction of the Eye*, Syd. Soc. ed., p. 588), who have found that the liquids removed from such an eye are capable of causing dilatation of the pupil when placed in another eye.

It having been demonstrated that the mydriasis of the atropinized eye is the result of an action upon the peripheral nerve-fibres, the question arises, Are the ends of the oculo-motor, the contractor of the pupil, paralyzed, or are the ends of the sympathetic, the dilator, stimulated, or is there a double influence, both of these actions occurring? Both Donders (*loc. cit.*, p. 589)* and Stellwag von Carion (*loc. cit.*, p. 92) insist that the paralysis of accommodation is proof of paralysis of the oculo-motor nerve, and it seems to me they do so with truth. However this may be, there is abundant direct proof that the oculo-motor fibres are paralyzed, since the experiment of Grünhagen, showing that galvanization of the exposed oculo-motor nerve does not affect the atropinized pupil, has been repeated, and its results confirmed by Dr. Gustav Engelhardt (*loc. cit.*, p. 321).

In artificial mydriasis there is, then, undoubtedly peripheral palsy of the oculo-motor. The question arises, Is there also stimulation of the dilating nerve? The evidence as to this is not so positive, but to my mind indicates very strongly that there is such an action. Clinical experience certainly shows that the dilatation produced by a mydriatic is not merely a passive movement of relaxation, but is active, capable of tearing up inflammatory adhesions even when of some firmness. Again, the dilatation that occurs after the paralysis of the oculo-motor nerve in man and after its destruction in animals is not at all equal to that produced by atropia, and, indeed, can be largely increased by the action of the drug; further, in the eye separated entirely from the nerve-centres (see above) atropia still causes a wide dilatation; facts which necessitate the belief either that the alkaloid acts upon the sympathetic fibrillæ, or that the peripheral fibres of a nerve are in themselves nerve-centres, acting upon the muscle of themselves even when separated from their centres.

It has been urged against the view here taken that even the widest artificial mydriasis is increased by galvanization of the sympathetic. De Ruiter

* See also Von Graefe, *Deutsches Klinik*, 1861.

states the contrary ; but since Grünhagen, Hirschmann, and Engelhardt separately affirm as the result of personal experiment the correctness of the asserted fact, it must be accepted. Granting its truth, I do not think it warrants the deduction, since it is very conceivable that an agent may excite the peripheral filaments of a nerve very greatly, and yet not to that point that they are incapable of further excitation.

In conclusion, the action of atropia applied to the eye may be summed up as follows: the mydriasis is the result of a direct influence upon the peripheral nerve-fibres, those of the oculo-motor being certainly paralyzed, those of the sympathetic and its ally the trigeminus being probably excited.

In regard to the constitutional action of atropia, it is evident that when the alkaloid is administered internally there are only four possible ways in which it can cause mydriasis, and that these are as follows:

1. By acting alone on the sympathetic centres, as a stimulant. 2. By acting alone on the oculo-motor centres, as a paralyzant. 3. By combining these actions. 4. By being carried to the eye, and acting as though locally applied.

Authors are greatly at variance in their conclusions: thus, Harley (*The Old Vegetable Neurotics*) and Hayden (*Dublin Quarterly Journal*, August, 1863) may be cited as in favor of the first view, and Budge (*Ueber die Bewegung der Iris*, 1855), Braun (*Archiv für Ophthalmologie*, Bd. v. Abth. ii.), and Hirschmann (*Zur Lehre von d. durch Arzneimittel, etc., Reichert's Archiv*, 1863) as favoring the second.

Neither the first nor the second view is, however, tenable: the first, because of a fact which has been asserted by authorities and which I have experimentally corroborated, namely, that atropia given hypodermically causes dilatation of the pupil after section both of the trigeminus and of the sympathetic in the neck: the second, for the reason that after section of the oculo-motor in animals, or after complete paralysis of the oculo-motor in man, the mydriasis is much less than that of atropia-poisoning.

In regard to the third view, I have noticed that the dilatation of the pupil under the constitutional action of the alkaloid after section of the cervical sympathetic is still greater than that which is normally produced by oculo-motor paralysis. At my solicitation, Dr. T. G. Morton, of this city, cut down to the optic nerve in a rabbit and divided all the structures about it. The pupil contracted very much at the time; the cornea was not sensitive, but recovered its sensitiveness in part after some days. Atropia given hypodermically dilated the eye very markedly, but not nearly to the extent of the other eye. If in this experiment all the ciliary nerves were really cut, the proof is conclusive that the mydriasis is not of centric origin; the reason that the pupil did not dilate so freely as the other being the strong contracting influence it was under, and the great reduction in the amount of blood, *i.e.*, the amount of atropia, entering the eye, owing to the division of the blood-vessels.

An experiment of Lemattre, if it be accurate, is also conclusive in proving that the action of the mydriatics upon the pupil, even when administered internally, is a peripheral one. He succeeded in producing mydriasis in normal eyes by placing in them aqueous humors taken from dogs poisoned with atropia, and even from a fetus whose dam had been killed by the alkaloid. Donders, however (*loc. cit.*, p. 589), failed to get the dilatation; and two or three experiments have yielded me the same negative result. It requires no elaborate argument to prove that in this case a negative result does not overpower a previous positive one: still, the experiments of Lemattre need confirmation. In the Pennsylvania Hospital, under the care of Dr. Morton, not long since I saw a man who had been wounded by a railroad accident in such a way that the whole of the temporal bone anterior to the petrous portion was thrust into the side of the head; there was complete paralysis of the facial, of the trigeminus, and of the oculo-motor, as could be readily demonstrated upon the man, who lived some months, finally dying suddenly of abscess of the brain. The carotid canal was so pressed upon that the sympathetic, which passed upwards through it to the eye, must have also been paralyzed. The eye was, as proven by the autopsy, separated from all connection with the nerve-centres, and yet when atropia was given hypodermically the pupil dilated. The proof seemed complete that the mydriasis was owing to a peripheral action.

Our knowledge of the action of atropia upon the pupil may be summed up as follows: Atropia applied locally causes mydriasis by paralyzing the peripheral ends of the oculo-motor nerve, and probably by stimulating the peripheral ends of the sympathetic. Atropia given internally *almost* certainly causes mydriasis, not by influencing the nerve-centres, but by being carried in the blood to the eye itself, and there acting precisely as when applied locally.

THERAPEUTICS.—The results of clinical experience are in strict accord with what is known of the physiological action of belladonna. The chief indication for its use is to *relax spasm*. In the case of voluntary muscles its powers are comparatively feeble, except when it is *thrown directly into the muscle affected*. In this manner Dr. S. Weir Mitchell (*Injuries of Nerves*, Philadelphia, 1872, p. 258) has obtained very marked relief in the fearful spasms following nerve-wounds, and Dr. J. M. Da Costa in rheumatic spasm (*Pennsylvania Hospital Reports*, 1868). The benefit derived in this way is evidently due to the depressing effect of the drug upon the terminal nerve-filaments, with which it comes in direct contact, and to a certain extent also on the muscle itself. When given by the mouth, so small an amount of the remedy reaches the diseased part as scarcely to affect it, and very little or no relief follows. As has been previously shown, the non-striated muscles are more affected than the striated by belladonna, and clinically the drug is found to be even more efficacious in *spasm* of the *involuntary* than of the voluntary muscles: in such cases it is often of value used internally. It must be thus ad-

ministered in *lead colic*.—in *simple spasmodic colic*.—in *spasmodic dysmenorrhœa*.—in *spasmodic constriction* of the bowels with *obstinate constipation*,—in *laryngismus stridulus*.—in *nervous cough*,—in *asthma*.—in *hiccough*,—in *whooping-cough*.—in which, as originally advised by Bretonneau, it has been largely used, and is one of the best known remedies; also, even in the spasm accompanying passage of *renal* and *biliary calculi*, where of course it often fails. Wherever it is possible, however, it should be used locally in spasm of the involuntary as well as of the voluntary muscles. Thus, in *spasm* of the *urethra*, the ointment should be rubbed in along the canal; in *rigid os uteri*, the extract should be applied directly to the os; in *asthma*, belladonna should be inhaled, either by means of the cigarette or of the atomization of a decoction of the leaves; in *spasm* of the *sphincter ani* from *fissure* or other cause, it should be applied directly to the part by poultice or ointment.

It is, no doubt, by relaxing spasm, or rather by lessening irritability, that belladonna acts in that *nocturnal incontinence* of urine which is seen generally in children. It has been taught that this affection is due to a relaxation of the sphincter, but undoubtedly in the great majority of cases its real cause is an irritability of the bladder itself, so that spasmodic contraction occurs under the stimulus of a small portion of urine. A common result of the ingestion of a large dose of belladonna is a paralytic retention of urine, due no doubt to the local action of the atropia in the urine upon the bladder. It is needless to point out more in detail how the indications in nocturnal incontinence are met. Suffice it to say that clinical experience proves the great value of belladonna in such cases.

Under the present indication also probably belongs the use of the remedy in *constipation*. In doses of one-quarter to one-half grain of the extract, added to a laxative pill, belladonna is of great service, and sometimes used alone will cure chronic constipation. It appears to be of most value in subjects of rigid tone; in feeble, relaxed people, strychnia is preferable.

To relieve Pain.—Physiologically viewed, atropia should be of little value for this purpose; and I think clinical evidence bears this out. Dr. Mitchell has had probably the best opportunities ever afforded for testing this, and he says decidedly that it is of little use in severe suffering. My own experience is to the same effect. There is, however, considerable evidence of its value in *neuralgia*, but it is chiefly as to its efficiency when injected immediately in the neighborhood of a painful nerve, or applied as inunction over its course when superficial. In large quantity, belladonna certainly affects the afferent nerves, and, used as above, may readily relieve pain. It is very probable also that at times it cures neuralgia by modifying the circulation in the affected part. Though these things be so, yet belladonna is almost immeasurably below opium as an analgesic. In some forms of *neuralgia* with *spasm* it is of service by a double action.

To impress the Heart and Blood-Vessels.—Under the idea of its contracting the blood-vessels, belladonna has been highly commended by Dr. Harley

(*loc. cit.*) in *pneumonia*, acute *nephritis*, and various other acute diseases. I have had no experience with it in these affections, but the published accounts do not seem to me to indicate that it is of equal value with other remedies or combinations of remedies. In chronic albuminous nephritis I have tried it, as recommended by Dr. Harley (*loc. cit.*), but have failed to derive any advantage from it.

Possibly it is by acting on the blood-vessels that belladonna is so useful in ordinary sore throat, but it is more probable that it does good by relaxing the pharyngeal muscles.

As a stimulant to the circulation, belladonna has probably not been employed as much as it ought. Dr. Graves, however, commends it especially when the pupil is contracted in *typhus fever*, and it has been used with asserted advantage in *erysipelas*, *scarlet fever*, etc. I think this use of belladonna offers a very inviting field for therapeutic investigation.

To arrest Secretion.—Arresting secretion of the salivary glands by paralyzing the extreme branches of the chorda tympani has already been shown to be a physiological action of belladonna, and it follows from this that the drug should be useful in *ptyalism*. I have tried it in several cases of *mercurial salivation*, and found that it arrests almost at once the discharge of saliva, and seemingly facilitates greatly the return to health. In *colliquative sweats* it was originally recommended by Dr. S. Ringer, and I have found it of very great service. A full dose of belladonna extract, or one-sixtieth to one-eightieth of a grain of atropia used hypodermically, at bedtime, will very frequently prevent the usual *night-sweat*. In *colliquative diarrhoea* it has been recommended by M. Delpage, and very probably will be found of service.

Employment in Opium-poisoning.—It is stated that as far back as 1570 it was asserted that opium and belladonna are, in their influence upon the system, antagonistic. In the early part of the present century their employment as counter-poisons was again brought prominently before the profession; but, although a few scattered earlier records of their use as such exist in medical literature, it was not until the paper of Dr. Wm. F. Norris appeared (*American Journal of the Medical Sciences*, Oct. 1862) that general attention was attracted to the subject. Since this publication, very many cases of the use of the one medicine in poisoning by the other have been published.

The opposite actions of belladonna and of opium upon the pupils no doubt first suggested the idea of their antagonism; but in the light of recent experiments these apparently opposite effects upon the eye cannot be considered as proving any antagonism between the drugs, since it is almost certain that the dilatation is due to a peripheric and the contraction to a centric influence. Neither has any positive experimental evidence of the existence of these asserted antagonistic relations been brought forward. Indeed, I have not met with any investigations upon animals sufficiently close and thorough to determine the question of their relations; but the experiments of Prof. Reese (*American Journal of the Medical Sciences*, vol. xi.) certainly indicate

that they are not only not antagonistic in the sense in which an alkali and an acid are, but that they are not antagonistic in the sense in which physostigma and atropia are. Further, I think any one will perceive that our present knowledge of the physiological action of the two drugs renders anything like complete antagonism very improbable. This last point might perhaps be well discussed here in detail: but, on account of limited space, other less obvious matters must have the preference.

When the subject in hand is looked at from its clinical aspect, the appearance is different: to tabulate and discuss the reported cases of opium- or belladonna-poisoning in which the counter-narcotic has been used would require very many pages, and I therefore can only state my opinion that these records establish the therapeutic value of atropia in opium-poisoning. The clashing between this conclusion and that arrived at in the previous paragraph is apparent rather than real. No one would question the value of alcohol in certain stages or conditions of opium-poisoning, and yet no one would claim that opium and alcohol are in any sense antagonistic. In opium-poisoning, death occurs chiefly through failure of the respiration. Atropia is the only known drug which exerts a decidedly stimulating effect upon the respiratory centres. It is evident that in advanced stages of opium-poisoning this property renders atropia an invaluable remedy. In protracted opium-narcosis the cardiac and vaso-motor actions of atropia are of service, but it should never be forgotten that the main influence for good is upon the respiratory centres.

The first improvement from atropia in these cases is usually increased frequency of respiration; and as the breathing becomes less embarrassed the other symptoms ameliorate, largely because of the increased aeration of the blood.

The double nature of profound opium-narcosis must not be lost sight of: the blood is saturated with carbonic acid almost to the dead-line, and much of the unconsciousness, much of the failing circulation, much even of the embarrassed respiration, is due to the presence of the gas.

As soon as the system is in a measure relieved of this load, it begins to rebound; emetics act, consciousness returns to some extent, the circulation frees itself, and the road leading towards health is entered upon.

It is a matter of the gravest practical importance to decide when, how, and in what quantities the mydriatic should be employed. The exhibition of belladonna should, I think, commence so soon as there is decided failure of the respiration. The stomach is so paralyzed in the narcosis that it is uncertain how fast absorption will take place in the viscus; and the drug should always be given hypodermically, in the form of the alkaloid if possible.

The first injection of atropia should be of such size that it could not possibly do harm, and one-fortieth of a grain is in most instances a fair commencing dose. Very generally several repetitions of this are necessary, and the delicate practical point is to decide how often these repetitions shall be indulged in.

I think that very frequently too much atropia is given, and believe that often a great deal of firmness is required in these cases not to use it too freely, especially since reliance is generally placed upon the pupils as a guide. They are, however, a very unsafe guide, as is apparent when it is remembered that whilst opium contracts them by influencing the nerve-centres, atropia probably dilates them by acting on the peripheral nerves.

It must not be forgotten that in doses of sufficient magnitude atropia paralyzes the nerve-trunks, and may thus increase the danger. A cardinal principle should, therefore, be to give no more of the mydriatic than is absolutely necessary. One-fortieth or one-sixtieth of a grain may be injected every fifteen, twenty, or thirty minutes, as the urgency of the symptoms may demand. The judgment should be formed from a bird's-eye view of the whole case, fresh atropia not being given so long as the respiration and other symptoms are undergoing amelioration, but the dose being renewed so soon as any tendency to a relapse is manifested. Thus, if under the influence of atropia in a case the respirations had risen from four to eight per minute, I would not use the counter-poison again until there was manifested a tendency for the respirations to grow less frequent, or unless for a long period there had been no improvement.

As a Local Sedative.—Locally and freely applied, belladonna is a sedative; and, I believe, to glandular as well as to muscular and nervous tissues. In this way it is often very useful in various local inflammations. In the form of a plaster it frequently appears to do good in *palpitations* of the *heart*. Its use locally in spasms and neuralgia has been sufficiently dwelt on. In *mastitis*, or when it is desired to dry up the secretion of milk, its local application to the breast is often very efficacious.

Whenever belladonna is used locally, in order to get its good effects it must be employed freely. At the same time, it should be remembered that a number of cases of poisoning by its external use have been reported (*Medical Times and Gazette*, Nov. 1856; also *London Pharmaceutical Journal*, 1871). In children it must be used with very great caution; in adults, with a reasonable amount of care, its external use is safe, provided directions be given to have it washed off so soon as any affection of the sight or dryness of the throat is induced.

Having myself no practical knowledge as to the use of atropia in diseases of the eye, I requested Dr. Wm. F. Norris, Clinical Lecturer on Diseases of the Eye, to prepare some remarks upon the subject, and he has very kindly consented to do so, as follows:

The Use of Atropia in Diseases of the Eye.—Pure atropia, from its slight solubility in water, is only applicable where we desire a moderate effect; the sulphate, however, can be dissolved in water in any desired proportion, and, therefore, is generally employed. When a four-grain solution of this salt is dropped into the conjunctival sac of a healthy and emmetropic eye, we find

that in about fifteen minutes the pupil commences to dilate, and that this dilatation rapidly increases, till in from twenty-five to thirty-five minutes it has attained its maximum. The power of accommodation, and consequent ability to read fine print, does not show any marked decrease till twenty-five minutes have elapsed, when the near point commences rapidly to recede from the eye, until in an hour and a half to an hour and forty minutes the power of accommodation is completely annulled, and only objects over twenty feet distant from the eye, or those presenting practically parallel rays, can be distinctly seen. On the second day after the application the power of accommodation begins to return, and increases rapidly up to the sixth day, but is usually not fully regained till from ten to fourteen days; the pupil remains with but little change till the third day, when it rapidly contracts, but has not fully regained its normal state till eleven or twelve days have elapsed.* The mydriatic action of the drug is far more marked from a moderately strong solution applied to the conjunctiva than from its internal use, even when it has been pushed to the production of symptoms of poisoning. Thus applied, it acts on the intraocular nerves and ganglia; and it has been proved by Graefe and Donders that when the aqueous humor of an animal is drawn off and collected after its application, it contains a sufficient amount of the drug to cause dilatation of the eye of another animal when applied to it. These experiments have been abundantly confirmed by later observers, and the rapidity of its action appears to depend on the thickness of the cornea and the age of the subject selected for the experiment. It will be apparent from the foregoing statements that the use of a strong solution of atropia is not to be undertaken without due consideration, inasmuch as it is likely to debar the patient from any satisfactory use of the eyes for a period of from five to ten days. It is, however, invaluable, from its annihilation of the accommodation, where we wish to determine with accuracy the refraction of the eye, and is daily used for this purpose in cases of hypermetropia where the patients find it impossible to relax their accommodation, in astigmatism where it is necessary to determine the exact difference between the two principal meridians of the cornea, and in the rare cases of myopia associated with spasm of the ciliary muscle. To obtain this complete paralysis of the accommodation, a few drops of a four-grain solution should be dropped in the eye, and this repeated after an interval of five minutes. The patient will be ready for examination one and a half hours subsequently. In the vast majority of cases it is entirely unnecessary to dilate the pupil to obtain a satisfactory view of the fundus with the ophthalmoscope; but where this becomes necessary we can often use with advantage a solution of one-twentieth grain in an ounce of water: a drop or two of this will dilate the pupil, without, however, rendering it absolutely immovable, and with scarcely any interference with the accommodation;

* For a more detailed discussion of this subject, see *Donders's Anomalies of Refraction and Accommodation*, p. 584, New Sydenham Society, 1864.

on the next day the pupil is much smaller, and on the third day no trace of its action remains. In cases of suspected cataract the pupil should always be dilated; otherwise we may readily fail to discover the lesion, which frequently first manifests itself in a few faint striæ shooting out from the periphery of the lens. Moreover, it affords us a valuable prognostic point as to the probable success of any operation where the cataract is ripe; for where the iris fails to dilate ad maximum, we may be sure that it is more prone to take on inflammatory action, and more liable to be pressed on by any cortical matter which may remain behind in the eye. Daily experience shows that after the evacuation of the aqueous humor in the operation for cataract, the iris will contract in spite of any previous use of atropia; but as soon as by the closing of the wound the humor reaccumulates and the anterior chamber is re-established, the atropia resumes its sway. It is most useful in all inflammations of the cornea. In phlyctenular keratitis, by its local anæsthetic action on the branches of the trigeminus, it diminishes the photophobia and blepharospasm, and seems to mitigate the intensity of the inflammation by its influence in contracting the ciliary vessels, thus diminishing the supply of nutritive material carried to the cornea. Where an ulcer has perforated the central region of the cornea, and a prolapse of the iris has ensued, the energetic use of atropia often enables the radiating fibres of the iris to detach it from the cornea as soon as the opening has been plugged by lymph, and the anterior chamber restored, thus preventing the formation of anterior synechiæ. Mackenzie long ago called attention to the "healing and anodyne" effect of atropia in ulcers of the cornea occurring in the ophthalmia of new-born children,—a fact since universally recognized, and which holds good equally in the ulcers resulting from other forms of purulent conjunctivitis. It is held by many writers that part at least of this beneficial action is to be ascribed to its diminution of intraocular tension.* In cases of iritis a strong solution of atropia should at the outset be applied repeatedly at short intervals until we have obtained a full dilatation of the pupil, and subsequently sufficiently often to maintain the iris in this condition. We thus place the inflamed tissue at rest, diminish its blood-supply, and prevent the formation of posterior synechiæ, which constitute so frequent a cause of the recurrence of this

* That atropia diminishes intraocular pressure has been asserted by Graefe, Coccius, and many good authorities; and a reference to any manual of diseases of the eye will show that it is an accepted doctrine with most clinical observers of the present day. Other weighty authorities, such as Donders and Stellwag, doubt its ability to produce any such effect. Certainly it fails to reduce intraocular pressure in cases of glaucoma. Numerous attempts to decide the question by physiological experiment have been made by Hensen and Völckers, Wegner, Adamuk, Grünhagen, Dor, etc. The results vary, however, considerably among themselves. Those with the manometer are open to the objection that the cutting open the eyeball for its introduction quite changes the conditions of intraocular pressure and circulation, and that very slight movement of the instrument or eyeball invalidates the results; while those with the tenometer are also unsatisfactory, inasmuch as we have yet failed to obtain any reliable instrument for measuring intraocular tension.

disease by their mechanically-irritating effect: putting a strain on the iris in its every motion, hindering that variation in the diameter of the pupil which normally takes place with every change of convergence and with every variation of the intensity of light. Moreover, extensive synechiæ where the iris is plastered down to the anterior capsule by large patches of lymph tend to produce cataract, by interfering with the nutrition of the lens. Where the synechiæ are narrow and tongue-shaped, they may often be torn through by its action, even after the lapse of years.

Atropia acts more powerfully in iritis after the abstraction of blood,* and occasionally, where there is much exudation, fails to produce its effect till after the constitutional effect of mercury has been obtained. In some individuals the instillation of a strong solution of atropia, by its rapid passage through the tear-passages into the nose and throat, produces symptoms of slight belladonna-poisoning,—viz., flushed face, rapid pulse, dryness of the throat, slight dysphagia; but this may usually be diminished or prevented by gargling the throat with water, by compressing the canaliculi, or by everting the lower punctum lachrymarum. In some individuals it produces a curiously-irritant action on the conjunctiva, known as atropine conjunctivitis; this is sometimes quite severe, calling forth almost erysipelatous symptoms, at other times bringing out a crop of granulations. This is owing to idiosyncrasy, and will then happen with perfectly neutral solutions. It should in such cases be discontinued, and the conjunctivitis which it has called forth combated by weak solutions of alum and sulphate of zinc or other mild astringent. We find, too, in some cases of iritis, especially those occurring in rheumatic patients with posterior synechiæ due to repeated previous attacks, that we not only fail to dilate the pupil, but that atropia acts as an irritant to the eye. We are then obliged to give up its use, and resort to the application of dry warmth and appropriate constitutional remedies. Atropia will occasionally, in cases of chronic glaucoma, precipitate an acute attack, and is, therefore, to be used with due consideration in this disease.†

A great deal has been written about the value of belladonna as a prophylactic in *scarlatina*, and authorities differ very much. For a discussion of the subject I must refer the reader to the treatises of Dr. Geo. B. Wood, of Dr. Stillé, of Dr. Waring, etc. I have never had a decisive opportunity of testing the matter, but have no faith in the efficacy of the remedy. It ought to be tried further, however. The plan practiced has been to dissolve two grains of the extract in a fluidounce of water, and give two drops daily to a child a year old, adding one drop for every year in older children up to twelve years.

* Mackenzie, *Diseases of the Eye*, p. 537, London, 1854. Graefe, *Archiv für Ophthalmologie*, vol. ii. part 2, p. 209 (note).

† Graefe, *Archiv für Ophthalmologie*, vol. xiv. 2, 117. H. Derby, *Trans. Amer. Ophthalm. Soc.*, p. 35, 1868. Wells, *Diseases of the Eye*, p. 517 (2d Amer. ed.).

TOXICOLOGY.—Sufficient has already been said about the general symptoms of belladonna-poisoning. Those which are characteristic are the dryness of the throat, the increased frequency of breathing, the dilated pupils, the red efflorescence on the skin, the rapid pulse, the active talkative delirium, sometimes convulsions, all ending in abolition of function, as shown by stupor, rapid feeble pulse, cold extremities, and paralysis.

If the urine of a patient suffering from belladonna-poisoning be dropped into the eye of the cat or rabbit, it will dilate the pupil; and the diagnostician may avail himself of this test in any doubtful case.

Dr. Morel (*Annales de la Société de Médecin de Gand*, 1873) calls attention to a sort of laryngitis produced by poisonous doses of belladonna, and characterized by pain in the larynx, roughness of voice, and the expectoration of minute, pearly, tough pellets. It was present in the advanced stages of two cases of poisoning under his care.

The minimum fatal doses of the preparations of belladonna are scarcely known. An enema representing eighty grains of the root has produced death in five hours (*Casper's Wochenschrift*, Feb. 1845); but, on the other hand, recovery has occurred after the ingestion of three drachms of the extract (Taylor's *Principles and Practice of Medical Jurisprudence*, London, 1873, p. 432).

A tenth, or even a twentieth, of a grain of atropia will often produce alarming symptoms; yet Dr. Chambers reports (*London Lancet*, 1864) recovery in a child four years old who had taken about two teaspoonfuls of a solution containing a grain of the alkaloid in half an ounce.

After death from belladonna, no characteristic lesions are to be found.

In the treatment of belladonna-poisoning, the first indication is to prevent the absorption of any more of the poison. For this purpose emetics or the stomach-pump should be used. The same difficulties are to be met and the same measures adopted as in opium-narcosis; and, as it is well in that poisoning to exhibit tannic acid freely, because it forms with the alkaloids salts which are soluble with difficulty, so should vegetable astringents be given when an overdose of belladonna has been ingested.

After the stomach has been evacuated, symptoms must be met as they arise. During the first febrile stage, I would not hesitate to use cold bathings; and the moderate use of tartar emetic, or other sedative, is perhaps justifiable. The value of opium in belladonna-poisoning has not, I think, been determined, and its use should only be tentative, although good is to be expected from its judicious employment.

In poisoning by a mydriatic, in order to keep up the respiration and the circulation during the stage of failure of function the same measures should be employed as in opium-poisoning. External stimulation by heat and by mustard, flagellations, etc., artificial respiration, and especially the use of the alternate cold and hot douche, should all be practiced as necessary. I am not aware that physostigma has been used as a counter-poison to atropia in man;

but the experiments of Dr. Fraser have certainly demonstrated that the two drugs are antagonistic within certain limits. As this subject can best be elucidated after the discussion of the action of Calabar bean, the reader is referred to the article upon that medicine for further information. After toxic doses of belladonna, there is very generally complete retention of urine; and as this secretion contains the greater part of the ingested poison, and as reabsorption in the bladder is at least conceivable, the catheter should be used early.

ADMINISTRATION.—Belladonna is never used internally in substance, but in one of the following preparations, or in that of its alkaloid. All the preparations except two (designated below) are made from the leaves.

Tinctura Belladonnæ—*Tincture of Belladonna* (℥ii to Oj). Dose, ten to thirty drops.

Extractum Belladonnæ—*Extract of Belladonna*. A fresh-juice extract. Dose, one-eighth to one-half a grain.

Extractum Belladonnæ Alcoholicum—*Alcoholic Extract of Belladonna*. Dose, one-eighth to one-half a grain.

Extractum Belladonnæ Radicis Fluidum—*Fluid Extract of Belladonna Root*. Dose, one to two minims.

For external use the following preparations are officinal:

Unguentum Belladonnæ—*Belladonna Ointment* (extract grs. lx to ℥i). A soft ointment.

Emplastrum Belladonnæ—*Belladonna Plaster*. This plaster represents in strength its weight of belladonna root.

Both *atropia* and its *sulphate* (*Atropiæ Sulphas*) are officinal. The latter is most commonly used, on account of its solubility in water. The alkaloid or its salt is much preferred to the extracts for the production of mydriasis, on the score of cleanliness, as well as on account of the greater rapidity of its action. To dilate the pupil for ophthalmoscopic purposes, little gelatine wafers containing the one-fiftieth to the one-hundred-and-fiftieth of a grain of the alkaloid are often employed. One-sixtieth of a grain of atropia or its salt, given hypodermically, will generally produce slight dryness of the throat or other indications of its constitutional action. Where rapidity of action is required, this is the best method of administering belladonna.

STRAMONII FOLIA—STRAMONIUM LEAVES.

STRAMONII SEMEN—STRAMONIUM SEED.

The leaves and seeds respectively of the *Datura Stramonium*, or *Jamestown Weed*, a coarse, bushy, annual herb, three or four feet high, growing in waste places both in this country and in Great Britain, and readily distinguished by its large, funnel-shaped, whitish, fetid flowers and its quadrivalve spinescent capsules. The leaves are large, smooth, ovate, irregularly sinuate, with large acute teeth. The seeds are small, brownish black, reniform, with a feebly bitterish narcotic taste.

The active principle is an alkaloid discovered by Geiger and Hesse, and named *Daturia*, but which has been shown by Von Planta to be identical with atropia, possessing the same composition and having similar physical properties.

PHYSIOLOGICAL ACTION.—The physiological evidence as to the identity of *daturia* and *atropia* is in strict accord with the chemical proof. The symptoms of poisoning by stramonium differ in no respect from those of belladonna-poisoning, although Laurent (*De l'Hyoscyamine et de la Daturine*, Thèse, Paris, p. 22, 1870) asserts that irregularity of the heart's action is more marked under the influence of stramonium. The same accelerated pulse, the same elevation of temperature, the same wild delirium, the same increased frequency of respiration, the same widely-dilated pupils, the same red efflorescence on the skin, the same restlessness or convulsions, occur in both cases, and end alike, when the dose has been sufficiently large, in abolition of the functions of circulation, respiration, and innervation: stupor, general paralysis, weak, rapid, thready pulse, threatened asphyxia, constituting the phenomena of the closing scene in poisoning from either narcotic.

The most careful minute investigation of the physiological action of *daturia* which I have met with is that of Charles Laurent, already quoted. In his experiments that observer found that under the microscope the capillaries of the frog's web could be seen to contract after the application of *daturia*, even when the nerves of the limb had been previously severed, and after the hypodermic injection of the alkaloid; that the pulse-rate and arterial tension were both augmented by small doses of the poison, but that by large doses the arterial tension was diminished, although the pulse was still increased in frequency; that when the heart was completely separated from all connection with the central nervous system, *daturia* reduced the number of its beats; that respiration is accelerated by the alkaloid, even after section of the pneumo-gastrics; that by moderate doses of the alkaloid the conducting power neither of the sensory nor the motor nerves is destroyed; that the muscular contractility is not affected; that small doses increase, large ones diminish, intestinal peristalsis.

Elaborate discussion of these facts seems unnecessary. It is seemingly demonstrated, both from a chemical and a physiological point of view, that *daturia* and *atropia* are identical.

THERAPEUTICS.—Stramonium may be used to meet precisely the same indications as belladonna. It has been especially employed in spasmodic *asthma*, chiefly in the form of cigarettes made by rolling up the dried leaves. These are often very efficient when there is no organic disease; but their use requires some caution, as very alarming symptoms, if not fatal results, have been produced by them. In the form of cataplasms, stramonium leaves are often applied with advantage to painful *local inflammations*, *inflamed hemorrhoids*, etc.

ADMINISTRATION.—The preparations of stramonium, especially those of the leaves, vary a good deal in strength; the minimum dose should be used at first, and increased until an effect is produced.

Extractum Stramonii Foliorum—*Extract of Stramonium Leaves*. Dose, one-fourth to one-half a grain.

Extractum Stramonii Seminis—*Extract of Stramonium Seeds*. Dose, one-fourth to one-half a grain.

Tinctura Stramonii—*Tincture of Stramonium* (extract $\frac{3}{4}$ ii to Oj). Dose, ten to twenty minims.

Unguentum Stramonii—*Ointment of Stramonium* (seed $\frac{3}{4}$ i in $\frac{3}{4}$ i).

TOXICOLOGY.—Accidental poisoning, especially of children, by stramonium, is very common. It is scarcely necessary, however, to discuss the subject here, since in all points, as regards both symptoms and treatment, stramonium- and belladonna-poisoning are alike.

HYOSCYAMI FOLIA—HYOSCYAMUS LEAVES.

HYOSCYAMI SEMEN—HYOSCYAMUS SEED.

The leaves and seed respectively of the *Hyoscyamus niger*, a coarse herbaceous biennial, indigenous in England, and naturalized in the Northern United States. The leaves are large, oblong, ovate, deeply sinuated, very hairy. The seeds are minute, grayish, reniform. Like the leaves, they have a faint narcotic odor, and a bitterish acid taste.

In 1821 Peschier announced the existence of an alkaloid in hyoscyamus, but it was not until 1833 that Geiger and Hesse succeeded in obtaining it pure. According to Geiger, when slowly crystallized, *Hyoscyamia* occurs in transparent needles, and in star-shaped or bushy clusters of crystals. Thore states that out of chloroform it crystallizes in tables, out of benzole in fine needles, whilst out of ether and amylic alcohol it separates amorphous. Its chemical constitution has not been determined, and indeed its chemical history is very incomplete, since no distinctive test for it is known. When pure, hyoscyamia is odorless.

PHYSIOLOGICAL ACTION.—Hyoscyamia has a very similar, if not identical, action with atropia. Indeed, at present it seems probable that the active principles of belladonna and hyoscyamus are really one alkaloid. Schroff* asserts that hyoscyamia is decidedly more powerful than atropia, so far at least as concerns its local action on the pupil. On the other hand, Lemattre (*Archives Générales*, 1865) has found atropia much the more active of the two. It is, however, scarcely doubtful that the latter observer had an impure hyoscyamia, and it seems to me that it cannot be at present considered as determined that there is any marked difference in the mydriatic activity of the two drugs.

Both Schroff and Düllenberg* have noticed in man, as the result of the ingestion of hyoscyamus, dryness of the throat and mouth, brief sinking of the pulse-rate, followed by increased frequency, mydriasis, giddiness, muscular weakness, and insecurity of gait.

* Quoted by Husemann, *Die Pflanzenstoffe*, p. 474.

The experiments of Schroff, of Laurent, and especially of Harley (*The Old Vegetable Narcotics*), indicate that hyoseyamus is much more of a hypnotic than is belladonna. Yet Harley's recorded observations would seem to show that, in some individuals, hyoseyamus induces insomnia. According to Harley, the primary sinking of the pulse is much more marked after hyoseyamus than after its sister-narcotic.

Laurent (*De l'Hyoseyamine et de la Daturine*, p. 15) affirms that the delirium caused by hyoseyamus is calm, whilst that by atropia is furious; but on page 19 of his book is recorded a case of henbane-poisoning in which the delirium was "furious." Schroff (*Wochenblatt der Zeitschrift der Gesellsch. der Aerzte zu Wien*, 1865) asserted that pneumonia is a constant and even characteristic lesion of hyoseyamus-poisoning in the rabbit; but Lemattre (*loc. cit.*) has never seen more than little scattered points of hyperæmia; and in Laurent's experience even these have always been absent.

The most elaborate paper I have met with on the physiological action of hyoseyamia is that of Laurent. In his experiments he found that the capillaries of the frog's web contracted after the local application of hyoseyamia, even when the nerves had been previously severed, and also after the hypodermic use of the poison; that small doses augmented both the pulse-rate and the arterial tension, whilst large doses increased the former and diminished the latter; that the direct application of the alkaloid to the heart produced a rapid diminution of the number of its beats; that moderate doses increased the rapidity of respiration even after section of the pneumogastriæ; that upon the nervous system, the muscles, and the intestines, the action of hyoseyamia was the same as that of daturia. Authorities already quoted assert that hyoseyamus is more apt to induce sleep than belladonna. I do not think this has, however, been proven, and at present it seems impossible to point out any established difference in the action of the drugs. Like atropia, hyoseyamia is eliminated by the urine.

THERAPEUTICS.—Hyoseyamus may be used to fulfil any of the indications for which belladonna is employed. Clinical experience appears in a measure to bear out the assertions of various authorities as to the superiority of hyoseyamus as a hypnotic; yet such action of the drug is certainly very uncertain.

TOXICOLOGY.—It is not necessary to make any remarks upon the toxicology of this drug, since the diagnosis and treatment are identical with those of belladonna- and hyoseyamus-poisoning.

ADMINISTRATION.—*Extractum Hyoseyami*—*Extract of Hyoseyamus* (fresh juice). Dose, one to three grains.

Extractum Hyoseyami Alcoholicum—*Alcoholic Extract of Hyoseyamus* (dried leaves). Dose, one to three grains.

Tinctura Hyoseyami—*Tincture of Hyoseyamus* (dried leaves $\frac{3}{4}$ ii to Oj). Dose, half a fluidrachm to two fluidrachms.

CLASS VIII.—ANÆSTHETICS.

THE term *Anæsthetics* is here employed as the name of a group of volatile substances, whose vapor has the power of producing loss of consciousness, preceded by or accompanied with loss of sensibility and diminished muscular action. The medicinal properties of these substances are largely due to their volatility, by virtue of which they are very rapidly absorbed and almost as rapidly eliminated by the mucous membrane of the lungs. As a consequence of this, their action is very easily controlled. That they are taken into the blood, and thereby reach all portions of the system, has been abundantly proven by recent investigations.*

Many of the theories which have been suggested to explain the production of anæsthesia are so groundless that it seems unnecessary to discuss them here. All that are worthy of consideration may, I think, be arranged in three groups, as follows: 1st, those which assert that the symptoms are produced by a partial arrest of oxidation; 2d, those which look upon the anæsthesia as due to precedent physical changes in the blood; 3d, those which teach that the various agents employed act directly upon the various organs and tissues concerned.

As the last of these is the most natural, the burden of proof rests upon the supporters of the theories included in the first two groups.

All the proofs of these latter views which have, as yet, been brought forward, amount to no more than as follows: that in asphyxia the symptoms are similar to those of anæsthesia; that in profound anæsthesia there is an evident lessening of oxidation; and that some anæsthetics probably produce changes in the blood.

The objections to regarding these facts as proving the truth of the theories alluded to are very grave. Thus, it is very well ascertained that the symptoms of asphyxia are only analogous to those of anæsthesia, not identical,† and indeed that anæsthesia as caused by different agents offers different phenomena; also, there is no proof whatever that the lessened oxidation and the blood-changes which are believed to occur when anæsthetics are employed are causes of the nervous symptoms, and not simply coincident phe-

* See especially O. Schmiedeberg, *Inaugural Dissertation*, Dorpat, 1867, *Archiv d. Heilkunde*, viii., 1867.

† See Report of Chloroform Committee, *Medico-Chirurg. Trans.*, vol. xlviii. p. 329.

nomena. It must be insisted on, therefore, that these theories have never been proven. Moreover, positive proof of their incorrectness is not wanting. Thus, in regard to the theory of arrest of oxidation, there are substances, such as nitrite of amyl, which lessen oxidation very remarkably (Richardson, *Medical Times and Gazette*, p. 180, 1868), but are not anæsthetics; and an excess of oxygen in the air does not lessen the rapidity with which anæsthesia is induced.* In regard to the blood-theories, Ludimar Hermann (*Reichert's Archiv für Anatomie*, 1866, p. 27) calls attention to the fact that the anæsthetics produce the same general symptoms in the infusoria, which have no red blood, as in mammals; and Lewisson† has shown that they influence the so-called "salt frogs," which contain little or no blood, precisely as they do the normal frog, from which the conclusion is inevitable that they do not affect the frog by altering the nature of the blood or by inducing asphyxia.

I think, therefore, that the only theory at all compatible with our present knowledge is, that anæsthesia is in most cases due to a direct action of the agent inducing it. At the same time it should be borne in mind that nitrous oxide is probably an asphyxiating agent, and is probably not allied in its action to the true anæsthetics, such as chloroform and ether.

Many, and probably all, of these substances have the power of dissolving the red corpuscles. Ludimar Hermann (*Reichert's Archiv für Anatomie*, 1866) has pointed out a possible connection between this and their action on the nerve-centres. He states that *protagon*, which constitutes the stroma of the red blood-disks, is an important constituent of the nerve-centres. As death would necessarily occur before the protagon could be dissolved out of the nerve-centres,—*i.e.*, before it could be dissolved out of the red corpuscles,—it is evident that no extensive destruction of the latter bodies can occur from the action of an anæsthetic, and the patient survive.

Bile-acids also dissolve protagon, and Hermann states that some experiments he has made seem to indicate that they have anæsthetic properties. Before, however, the ingenious theory of Hermann can be considered fully tested, further investigations must be made.

In 1848 (*Archives Générales*, 2e sér., t. xvi., 1848), Duméril and Demarquay showed that during anæsthesia there is a reduction of temperature. This has been confirmed by Bouisson (*Traité théorique et pratique de la Méthode anesthésique*, Paris, 1850) and by Sulzynski (*Ueber die des Alkohols, Chloroforms und Aethers Einfluss auf den thierischen Organismus*, Inaug. Dissertation, Dorpat, 1865); and Scheinson (*Archiv des Heilkundes*, 1869) has demonstrated that there is no increase in the giving out of heat by the body during anæsthesia, and consequently that the anæsthetics lessen the production of animal heat, no doubt, like alcohol, by checking tissue-metamorphosis.

* See Report of Chloroform Committee, *Medico-Chirurg. Trans.*, vol. xlviii. p. 335.

† See *Chloral* for further details.

The chief purposes for which anæsthetics are used are to *relieve pain* and to *relax spasm*. To meet the first indication, they are employed by surgeons especially; but they are also exceedingly valuable in cases of suffering from disease. It must be borne in mind that their action is very transitory, and is accompanied by more or less disturbance of the general system, and that consequently they are only to be employed when the pain is exceedingly severe and transient. Anæsthetics are frequently used in surgery for the purpose of relaxing spasm, as in cases of *dislocation*, *hernia*, etc. In medicine, they have been employed in various forms of *convulsions*, and are especially valuable in severe *hysterical convulsions*, in *puerperal eclampsia*, and in *spinal convulsions*; in *epilepsy*, they are very rarely called for; in *infantile convulsions*, they may be sparingly used when the convulsion itself threatens life.

In various *spasms of excretory ducts or canals*, and especially during the passage of *calculi*, they act very favorably, both by relieving pain and by producing relaxation. In *asthma*, and in *spasmodic stricture of the œsophagus*, as in all other cases of oft-repeated spasm, they should only be administered to meet temporary indications, as their habitual use is deleterious.


To relieve pain, anæsthetics are used with great propriety during *child-birth*. In natural labor it is not commonly necessary to produce complete anæsthesia. When the full effect of either ether or chloroform is induced, there is almost always a weakening, and very often an abolition, of the uterine contractions. The anæsthetic should be administered in such quantities as to relieve the pain without decidedly interfering with the muscular spasm. In some cases this can be done, in others it is impracticable. I have obtained very advantageous results in some cases by suspending the pains for about half an hour by means of ether, and then entirely withdrawing the anæsthetic. By this treatment the weak, painful, ineffectual efforts of a worn-out, nervous patient may often be converted into regular, successful pains.

I think that the risk of *post-partum hemorrhage* is materially increased by anæsthetics, and therefore habitually give after their use two drachms of the fluid extract of ergot, as soon as the perineum is well distended by the child's head.

NITROGEN MONOXIDE—NITROUS OXIDE. ($\text{NO} - \text{N}_2\text{O}$.)

Nitrous oxide is a colorless, almost inodorous gas, of a sweetish taste. It is a very active supporter of combustion. Water absorbs nearly its own bulk of it. It is made by the distillation of the nitrate of ammonia, which resolves itself into the gas and water. Thus, $\text{NO}_3\text{NH}_4 = 4\text{HO}, \text{OH} + 2\text{NO}$. The accompanying wood-cut represents the best apparatus,* and the following rules embody in a few words all the necessary instructions for the production of the gas. As they are very concise, they should be followed to the letter.

* This apparatus is furnished by Mr. S. S. White, of Philadelphia.



Place the holder in a level position, and fill it with water to within an inch or two of the lower edge of the taper rim.

• Now open all the spigots, especially the one in the top of the receiver, and gently sink it into the water; then close the spigot in the top of the receiver and that on the upper edge of the holder, and arrange the weights that are attached to the receiver.

Now put pure water into the jars—No. 1 two-thirds full; Nos. 2 and 3 within one and a half inches of the top. Into jar No. 2 put a stick of caustic potash.

Arrange the jars in line, and connect them by tubing as shown in the cut. The tube from the retort must connect with the *long* pipe of jar No. 1, and so on. Always be sure that the *long* pipe of each jar is next the retort. When properly arranged, a current of air blown into the tube intended to connect with the retort will cause the water to bubble

in jars Nos. 2 and 3, and the spigot to which No. 3 is attached being open, the receiver will begin to rise.

Put into the retort the quantity of ammonia intended to be used (1 pound will make about 30 gallons of gas). Connect the long pipe of the first jar to the retort by rubber tubing, so that the condensed steam may run into it.

Now apply the heat *gradually*, and *watch* the process, being careful not to overheat the ammonia, which will be known by its violent boiling, instead of the gentle bubbling which indicates the proper temperature. Do not make gas faster than 30 gallons per hour.

The heat should be shut off before the ammonia is quite all decomposed, in order to avoid breaking the retort; and when the gas has ceased to come over, the retort should be disconnected from the tubing, and the receiving spigot closed.

It will be understood that the first jar is merely to receive the drip; the use of the water is simply to keep the jar cool. Some recommend placing this jar into a bucket of cold water.

The inhaling-tube is attached to the spigot at the top of the holder.

The register indicates the number of gallons of gas in the holder, as well as the quantity inhaled.

The water in jars Nos. 1 and 3 should be changed every time the gas is made; that in No. 2 about once a month, when fresh caustic potash should be added. The water in the holder will keep sweet a year or longer if gas is made frequently.

“The various effects of increasing heat upon nitrate of ammonia are shown by the following table:

“At 226° F. fuses perfectly.

“At 302° F. emits white fumes, condensing in drops.

“At 347° F. effervesces slightly.

“At 356° F. boils without decomposition.

“At 437° F. effervesces rapidly.

“At 460° F. begins to evolve gas.

“At 482° F. evolves gas in abundance.

“Above 500° F. *nitric oxide* is given off.

“To determine the temperature, thermometers have been prepared which may be passed through the cork and into the retort, marking the following degrees: 226°, 356°, 460°, 482°, 500° F.

“To obtain the largest amount of gas, the nitrate should first be melted in the retort at a temperature just sufficient (226° to 250° F.). When melted, the heat may be at once carried up to the point of decomposition, 460° F. If a gradually increasing heat is used after the salt is melted, a portion of it will sublime unchanged until the temperature reaches 460° F. The heat should never be allowed to rise above 482° F., for beyond that noxious products are generated. After the gas has begun to come over briskly, the appearance of copious white fumes in the retort is an indication that the heat is too great. The nearer the heat can be kept at the point necessary to generate nitrous oxide, the purer will be the gas.”—*Dental Materia Medica*. By Dr. Jas. W. White. Philadelphia, 1868.

After the gas has commenced to come over freely, it is generally best to lessen the amount of heat applied to the sand-bath.

The gas should be allowed to stand over water for a few hours before using.

It improves rather than deteriorates by age, and in a properly constructed gasoneter may be kept for many weeks, especially if the water between the holder and receiver of the gasometer be covered with a thin film of oil, so as to prevent its yielding to the air the gas which it absorbs, until saturated. In manufacturing the gas, whenever the alkalinity of the potash solution in the jar No. 2 is lessened, all the washing-jars should be emptied and refilled.

PHYSIOLOGICAL ACTION.—Although nitrous oxide has been used enormously as an anæsthetic, and many papers have been written upon it, yet our knowledge of its physiological action is due to three or four observers. When pure nitrous oxide gas is inhaled for from half a minute to three minutes, insensibility is produced, preceded in many cases by decided evidences of excitement. Some persons under its influence will sink quietly away into unconsciousness, but others will become hilarious, erotic, or pugnacious, and be restrained only by force. During the stage of anæsthesia the patient presents the appearances of asphyxia.

It is probable that the paralysis of function invades the different portions of the nervous system in the same order as does that caused by ether; but we have no positive knowledge upon the subject. A single experiment of Dr. Amory (*New York Medical Journal*, Aug. 1870) indicates that upon the conducting power of the motor nerves nitrous oxide exerts very little or no influence.

Neither have we any complete investigations as to the action of nitrous oxide upon the circulation: indeed, we have no studies whatever as to the condition of the circulation before anæsthesia is produced. Dr. Amory, by means of a cerebrometer set into the skull of the dog, has found that during the period of unconsciousness the cerebral pulsations are very decidedly lessened, and finally abolished, although the cerebral pressure is increased. This would seem to prove that the capillary cerebral circulation is checked and finally arrested, but that the amount of blood in the cerebrum is not lessened.

It seems pretty well established that the gas is not capable of yielding its oxygen so as to support life. A taper will burn in it, it is true, but its decomposition is due to the high heat, and at the temperature of the body the compound is a stable one. MM. Jolyet and T. Blanche have found (*Archives de Physiologie*, July, 1873) that seeds will not germinate in it, and that animals (frogs, sparrows, guinea-pigs, rabbits) live no longer in an atmosphere of pure nitrous oxide than in one of nitrogen. Even Dr. Colten, who believes that nitrous oxide produces hyperoxygenation of the blood, states (*The Physiological Action of Nitrous Oxide Gas*, Philadelphia, 1871) that in an atmosphere of the gas a mouse will live only from thirty to sixty seconds, a pigeon one to two minutes, a kitten one to two minutes, a frog thirty to sixty minutes, —all dying of asphyxia. Surely, the absurdity of maintaining that the effects of nitrous oxide gas are due to increased oxidation needs no comment.

The French observers above named affirm that nitrous oxide has no effect upon the system; that the phenomena induced are simply due to deprivation

of oxygen. The series of facts which they have experimentally proven, and upon which their conclusions are based, are: 1. An animal lives no longer in nitrous oxide than in nitrogen; 2. Anaesthesia occurs at the time that the blood of an animal becomes black; 3. Animals breathing an air containing sixty to eighty per cent. of nitrous oxide, and twenty to forty per cent. of oxygen, are unaffected; 4. The analysis of the blood of two dogs yielded the following results:

No. 1. Conscious.		No. 2. Unconscious.	
Carbonic acid	. . . 46 per cent.	Carbonic acid	. . . 36.6 per cent.
Nitrous oxide	. . . 29 per cent.	Nitrous oxide	. . . 34.6 per cent.
Oxygen	 19.7 per cent.	Oxygen	 3.3 per cent.

and other analyses showed that the coma was not developed until the oxygen in the blood was reduced to three or four per cent. It is evident that if the above analyses are, as from their reports they appear to be, reliable, the anaesthesia is not due to the presence of carbonic acid in the liquor sanguinis, since nearly ten per cent. more of that gas was present in the blood of the conscious (No. 1) than in that of the unconscious dog (No. 2), and also that it is more rational to believe the decrease in the oxygen, rather than the slight increase in the amount of the nitrous oxide, made the difference between consciousness and unconsciousness.

In this connection the very careful experiments of Dr. Amory are most interesting. He found that during nitrous oxide narcosis the amount of carbonic acid exhaled from the lungs is only two-thirds of that eliminated before the inhalation, and that immediately after the recovery of consciousness less than one-third the normal amount of carbonic acid is given off.

This certainly is corroborative of the conclusions of the French observers, which have also received confirmation very recently by the experiments of Mr. Elihu Thomson (*Philadelphia Medical Times*, November 15, 1873), who found that animals in an atmosphere of hydrogen and of nitrogen, and also in vacuo, suffered symptoms precisely similar to those caused by the inhalation of pure nitrous oxide; also, that in man the inhalation of pure nitrogen causes the symptoms of nitrous oxide narcosis, except that the sense of exhilaration is absent; finally, that nitrogen, hydrogen, and nitrous oxide, as well as a vacuum, are rendered capable of supporting life if a proportion of oxygen approaching that existing in common air be introduced.

It would seem that this accumulated evidence is sufficient to show that the anaesthesia produced by the inhalation of nitrous oxide is simply asphyxia, and that its administration is simply a neat way of stopping the supply of oxygen, and of thereby inducing asphyxia; yet the marked excitement which sometimes precedes the anaesthesia, and the fact that the gas, even when largely diluted with air, produces in man a feeling of exhilaration, indicate that it is not merely a passive agent.

THERAPEUTICS.—As an anaesthetic, nitrous oxide is chiefly valuable when

short, minor operations are to be performed. The rapidity with which insensibility is induced, and with which it disappears, combines with the general absence of after-effects to enhance the popularity of the gas. When, however, an operation requiring some little time for its performance is intended, ether is far preferable. Nitrous oxide has been administered to many thousands of persons, and until recently no deaths at all attributable to it have occurred. Two have, however, been reported within a short time. In one of these (*Dental Cosmos*, editorial, June, 1872) it is doubtful whether the gas had anything to do with the fatal result, or, indeed, whether it was really administered. In the other instance (*British Journal of Dental Science*, Feb. 1873), death from asphyxia, apparently induced by nitrous oxide, occurred in a healthy man. The gas seems to have been pure, and the operator skillful; but, as no post-mortem was made, the case is enveloped in some uncertainty.

Nitrous oxide is best administered from an india-rubber bag, which should contain at least eight gallons. The mouth-piece should always be made with two valves, so that the expired gas will be thrown out into the air, and not back into the bag.

ETHER—ETHYL OXIDE. ($C_4H_5O - C_4H_{10}O$.)

Ether is a colorless, very volatile liquid, obtained by the dehydration of alcohol by sulphuric acid. It is very inflammable, as is also its vapor, which is two and a half times heavier than air. It is freely soluble in alcohol, and is itself a powerful solvent. Its odor is strong and peculiar; its taste hot. Its specific gravity, when pure, is 0.713, and its boiling-point 95° F. It is official in two forms.

ÆTHER—ETHER.—This ether contains a large percentage of alcohol, and has a specific gravity of 0.750. When shaken with an equal bulk of water, it loses from one-fifth to one-fourth of its volume. A test-tube full of it held in the closed hand should *begin to boil* on the addition of a piece of broken glass. Any specimen which does not conform to these requirements probably contains an overplus of alcohol or of water.

ÆTHER FORTIOR—STRONGER ETHER.—This ether should have a specific gravity of 0.728, should not lose more than one-eighth of its volume when shaken with an equal bulk of water, and in a test-tube held in the closed hand should *boil vigorously* on the addition of broken glass.

PHYSIOLOGICAL ACTION.—The first effects of ether when inhaled are burning in the fauces and a feeling of strangulation, both due to the local impression of the irritant vapor. The first indications of its systemic action are a sense of exhilaration and a lightness in the head, associated with a roaring or buzzing in the ears. These are soon succeeded by a feeling of the immediate surroundings being afar off, and this soon fades into semi-unconsciousness, with visions and illusions. These are of various characters, and are often

accompanied by a species of delirium. Some patients weep, others laugh; some shout, some pray, some rave, and some become exceedingly pugnacious. In rare instances, the dreams become erotic; and cases are on record in which there were distinct evidences of the occurrence of a complete venereal orgasm. In this stage, the patient in most cases may be more or less perfectly aroused. There is very rarely, if ever, anæsthesia before the period of complete unconsciousness.

The second stage of ether-narcosis may be considered to commence with the complete loss of consciousness. In most cases, some degree of muscular rigidity is at first still present, but soon passes off, and the patient lies relaxed and quiet, with slow, regular, automatic respiration.

The occurrence of stertorous respiration, due to a paresis of the muscles of the palate, shows that the stage of muscular paralysis is being reached. It should, except in very rare cases, be the signal for the immediate withdrawal of the anæsthetic.

The face during etherization is reddish; marked pallor or lividity are respectively important indications of failure of the heart's action or of respiration. The stage of excitement generally lasts only a few minutes, but in some cases is prolonged, and in nervous women may pass into a violent fit of hysterics, which soon yields, however, to a persistent use of the anæsthetic. The pulse is quickened and increased in force by ether, and it will often maintain itself during a prolonged narcosis. On the lower animals ether acts precisely as it does on man.

As the functions of the cerebrum are affected before those of the other portions of the nervous system, it is very apparent that the brain is especially sensitive to the narcotic. Flourens (*Comptes-Rendus*, vol. xxiv., 1847, pp. 161, 242, 253, 340) has found that, at a certain stage of etherization, pricking of the anterior or motor nerve-roots causes motor disturbance, although the posterior or sensory portions of the spinal centres are completely insensible. After a more prolonged inhalation, the anterior or motor centres also fail to respond to mechanical irritation, although the functions of the medulla oblongata are being regularly performed, and stimulation of its anterior centres gives rise to motor disturbance, and pricking of its sensory portions even causes manifestations of pain.

If the inhalation of ether be sufficiently prolonged, according to the same authority, the sensory and finally the motor functions of the medulla oblongata are compromised, and death by a paralysis of the respiratory centres ensues.

Longet (*Archives Générales*, 4e sér., tome xiii. p. 374) in part confirms and in part questions the results of Flourens. He states that he has found the sensory functions abolished very early, but has never failed in any stage of the narcosis to get a response from the anterior part of the cord. These apparently different results are simply due to the fact that, whilst Flourens used only mechanical stimuli, Longet employed powerful galvanic currents.

Flourens was substantially correct, and the order of the involvement of the

nerve-centres in man and animals is first the cerebrum, next the sensory centres of the cord, next the motor centres of the cord, next the sensory centres of the medulla oblongata, and finally the motor centres of the medulla oblongata.

No experiments that I know of have been made to determine how far the nerve-trunks are involved in etherization. That they are to a slight extent affected seems, however, probable from the experiments of Longet (*loc. cit.*, p. 382) and of Serres (*Archives Générales*, 4e sér., tome xiii. p. 433). These observers found that the direct application of ether to a nerve produced a paralysis of the sensory fibres of that nerve; so that pinching the nerve below the point of application caused no pain, although voluntary movement was preserved, and galvanization of the nerve-trunk above the point of application induced spasms in the tributary muscles: *i.e.*, the power of conducting an impulse downwards was preserved, that of conducting it upwards was lost. By a longer application of the anaesthetic the function of the efferent as well as of the afferent fibres was abolished, temporarily at first, but, if the application were persisted in, permanently.

Upon the motor system of organic life ether certainly acts, but much less energetically than upon the voluntary system. Thus, after death from ether, the vermicular movements of the intestine, although less active than normal, are very rarely, if ever, entirely absent.

A close, accurate study of the action of ether on the circulation is wanting, but the English Chloroform Committee have shown that it increases very markedly the arterial pressure, and that even in a prolonged ether-narcosis there is no material diminution of this pressure until manifest failure of respiration has taken place.

When death is produced by ether, the heart generally continues to pulsate for a long time after the arrest of respiration.

The rise of arterial pressure during the inhalation of ether is probably due to an increase of the power of the heart, and to a stimulation of the vaso-motor nerves. The only experiments on these points with which I am acquainted are those of Sansom (*Chloroform: its Action and Administration*, p. 92, Phila., 1866). He states that the vessels of the frog's web can be seen to contract during the inhalation of ether, and that this vaso-motor spasm is very permanent, and does not yield to paralysis and passive dilatation until the anaesthesia has almost deepened into death.

It is frequently asserted that ether, when added to blood, coagulates it. A. Schmidt, however, states that the coagulation is due to ozone which has been generated by the ether and is contained in it, since freshly-distilled ether does not coagulate albuminous substances.

The researches of Wittich (*Schmidt's Jahrbücher*, Bd. cxlii. p. 212) and A. Schmidt (*Virchow's Archiv*, vol. xxix., 1864, p. 19) have shown that when ether is added to the blood of horses,* cats, or rats, the red corpuscles

* Schmidt says that sometimes crystallization fails (*loc. cit.*, p. 23) in the blood of the horse.

disappear in a very short time; and, as their stroma cannot be demonstrated by the aid of reagents, this disappearance is without doubt due to its solution. The hæmatin thus set free is dissolved in the serum, but the presence of the ether soon causes it to crystallize out.

There is no proof that these changes in the blood occur to any great extent when ether is inhaled; and the usual rapid recovery from the effects of the anæsthetic indicates that there is no profound alteration of the vital fluid.

An imperfect study by Harley of the effect of ether on the gases contained in drawn blood indicates that ether does not exert much influence upon their proportional amounts. It is, however, quite possible that a more thorough investigation would give a different result.

THERAPEUTICS.—As an anæsthetic, ether does not act with the rapidity and pleasantness of chloroform, but it has the advantage of safety. So dangerous is chloroform, and so safe is ether, that there is no excuse for the use of the former agent under ordinary circumstances.

The reason of the safety of ether is that, unlike chloroform, it never suddenly paralyzes the heart. It may kill by inducing asphyxia, but it does so slowly, and in the great majority of cases after warnings which can be overlooked only through the most reckless carelessness.

Many of the inconveniences which attend the use of ether can be obviated. Thus, in order to prevent the nausea which often follows the anæsthesia, the patient should avoid eating for at least six hours before the inhalation, and should take from one to two ounces of brandy just before commencing the latter process.

Unlike that of chloroform, the vapor of ether should be administered in as concentrated a form as possible. When so inhaled, in most persons, it will produce complete insensibility in from three to eight minutes, provided that no unchanged air be admitted during the time named.*

According to Dr. Snow, when air at 80° F. is saturated with ether, in every hundred parts of the mixture there are seventy-one parts of the vapor. This point is probably never reached in the practical use of ether, but the nearer it is approached the more rapid will be the induction of anæsthesia.

The ordinary method of the administration of ether in Philadelphia is as follows:

Out of stiff paper a cone is made of such size and shape that its base will fit closely over the nose and mouth of the patient. In this cone a napkin, a small towel, or a conical, hollowed-out sponge is to be placed. About an ounce and a half of ether having been poured on the napkin, the cone is to be closely applied to the face of the patient, and *kept there*. If anæsthesia

* Various inhalers have been invented for facilitating the use of ether, but they are very rarely used. Lente's inhaler probably does shorten the time and lessen the amount of ether necessary for the production of anæsthesia: I have, however, never seen it used. For information on these points, see *New York Medical Record*, May 1, 1866, and *New York Medical Journal*, April, 1871.

be not complete in four or five minutes, a second dose of ether should be put on the napkin, care being taken that the removal of the apparatus is only momentary. When patients are fastidious, and a few moments are of no importance, the gradual commencement of the inhalation is much more pleasant, as the first choking sensation is thereby to a great extent avoided.

In using ether at night, its inflammable nature must not be forgotten. I have seen fire flash from a candle through some four or five feet of diffused vapor to the sponge. The light should always be *above* the patient, as the high specific gravity of the vapor of ether causes it to fall towards the floor.

Administered by the mouth, ether has been used with advantage in various forms of *colic*, but is generally inferior to chloroform. When, however, as in some cases of *retrocedent* or *internal gout*, there is with the painful gastric and intestinal spasm a condition bordering on collapse, the stimulant properties of ether make it very valuable.

In sudden *sinking-spells*, either from poison or from natural causes, ether, as a powerful and very quickly-acting stimulant, is often indicated. In some cases of this description it may even be administered by inhalation. Of course, under these circumstances its influence should not be carried nearly to the point of producing anæsthesia.

As an *anthelmintic*, ether has been used by MM. Bourdier and Lortet with success against the *tape-worm*. For this purpose, an ounce and a half may be administered at once, followed in two hours by a full dose of castor oil.

In *hysteria*, *neuralgia*, *nervous headache*, and *spasmodic neuroses*, such as *hiccough* and *asthma*, ether is occasionally employed with benefit.

When ether is swallowed, it produces a sense of strangulation and choking, which seriously interferes with its use. For this reason, it is best given in capsules, or in ice-cold water. Probably large doses are best administered by putting them, mixed with an equal amount of brandy, on finely-cracked ice before drinking.

The dose is from one fluidrachm to a fluidounce.

CHLOROFORMUM—CHLOROFORM.

METHENYL CHLORIDE—TERCHLORIDE OF FORMYL ($C_2HCl_3-CHCl_3$).—This substance, which was discovered in 1831 by Mr. Samuel Guthrie, of Sackett's Harbor, N.Y., is produced by the action of chlorine upon alcohol. It is a colorless, limpid, neutral fluid, which is generally stated to be inflammable, although it can be made to burn with a greenish flame (*Foucault's Chemistry*, Am. ed., 1869, p. 566). Its taste is hot and sweetish; its odor fragrant and peculiar. It is soluble in alcohol and in ether, but when dropped into water it sinks, if pure, as transparent globules without milkiness. The alcoholic solution, when moderately diluted with water, forms an aromatic, sweetish liquid. It is antiseptic, but does not coagulate albumen.

In the U. S. Pharmacopœia it is officinal in two forms.

CHLOROFORMUM VENALE—COMMERCIAL CHLOROFORM.—The crude chloroform of commerce. Sulphuric acid, when shaken with it, after a time separates as a stratum of a dark brownish color. This color is due to a carbonization by the acid of the impurities in the chloroform.

CHLOROFORMUM PURIFICATUM—PURIFIED CHLOROFORM.—The U. S. Pharmacopœia directs that the commercial chloroform shall be purified by the action on it of sulphuric acid and by subsequent neutralization with carbonate of sodium, and distillation in a retort containing quicklime.

The *purified chloroform* should respond to the following tests: Sulphuric acid shaken with it should not be colored; it should not alter litmus blue, even after it has been exposed in a white glass bottle to direct sunlight for ten hours.

PHYSIOLOGICAL ACTION.—Although somewhat of an anæsthetic, chloroform applied locally is a powerful irritant. On the skin it produces redness and burning; if the evaporation be restrained, vesication is induced by it. Taken into the mouth, it causes a burning sensation, and, when swallowed, a sense of warmth in the stomach.

The vapor of chloroform, when inhaled, produces symptoms similar to those induced by ether, except that the choking sensations are absent, and that the stage of excitement is generally, but not always, shorter and less violent than that of etherization.

Dr. Snow (*On Anæsthetics*, London, 1858) divides the chloroform-narcosis into four degrees or stages, but the division adopted by Sabarth (*Das Chloroform*, Würzburg, 1866) and most writers seems more useful. This classification recognizes three stages. In the first of these, the symptoms are similar to those of alcoholic intoxication. This stage is generally very short, but in athletic persons, and especially in those who have been intemperate, it may be very long and very violent, and may persist after loss of consciousness. In drunkards, this excitement at times cannot be overcome without grave danger to life.

During this first stage, although consciousness is not lost, the sensibility is generally blunted, but very rarely is it altogether annulled. Dr. Coleman (Sansom, *Chloroform*, p. 55, Philadelphia, 1866) states, however, that he has extracted his own teeth without pain; and Dr. Snow relates the anecdote of a child who played with his toys during the operation of lithotomy.

During the second stage, which is that of anæsthesia, the consciousness and sensibility are abolished, the muscles are relaxed, and the patient lies perfectly quiet. This is the surgical stage, during which ordinary operations are performed. As already intimated, in some cases the first and second stages are united, so that violent excitement, muscular spasm, and rigidity may coexist with loss of consciousness and of sensibility.

The third stage is one of profound narcosis, with stertorous breathing, intense muscular relaxation, and abolition of ordinary reflex actions. This is

always a condition of danger, and its induction, except under very peculiar circumstances, by chloroform, is absolutely unjustifiable.

The pulse in the first stage of chloroform-narcosis may be quickened, even apparently strengthened; in the second stage it is generally about normal in frequency, but is more or less weakened; in the third stage it may be rapid and weak.

The action of chloroform on the nervous system, like that of ether, is chiefly upon the brain and spinal centres. Carter (*British Medical Journal*, Feb. 1867) found that very decided anæmia of the brain can be seen in animals subjected to its influence after the cerebrum has been laid bare; and accident in man (*Amer. Journ. Med. Sci.*, 1860) has furnished the corroboration of his experiments. Bernstein (*Schmidt's Jahrbücher*, Bd. cxlii. p. 227) has demonstrated that its action on the peripheral nerves is very slight. He found that there was no perceptible difference in the conducting power of the two ischiatic nerves of a frog, chloroformed after one of its iliac arteries had been tied. It is doubtful how far the muscular excitement of the second stage is due to real spinal exaltation and how far it arises from other causes. Bert asserts (*Comptes-Rendus*, t. lxiv., 1867) that the excitement is purely psychical, and that there is during the production of anæsthesia a steady lowering of reflex activity. He rests this assertion upon the fact that, in animals chloroformed after section of the cord, there is no motor disturbance below the point of section,—a fact which certainly demonstrates at least that the muscular excitement and the convulsions are cerebral.

Chloroform at first induces contraction and afterwards dilatation of the pupil. Dogiel (*Reichert's Archiv für Anat.*, 1866) has found that in the rabbit galvanization of the cervical sympathetic does not affect the iris during the stage of contraction, but that when the condition of partial dilatation has been produced the same procedure widens the pupil. He therefore logically concludes that the first contraction is due to stimulation of the oculo-motor nerve-centres, and the subsequent dilatation to paresis of the same.

In some animals the first effect of the inhalation of chloroform upon the circulation is a decrease in the frequency of the heart's action. Dogiel believes that this is due to a stimulation of the inhibitory centres, because he has found that it does not occur after section of the vagi. The after-increase in the rapidity of the pulse appears to be due, at least in part, to paralysis of the inhibitory centres, upon which chloroform seems to act as upon the oculo-motor centres, producing in them at first excessive functional activity, but afterwards functional paralysis.

The very careful experiments of the English Chloroform Committee (*Med.-Chirurg. Trans.*, vol. xlviii. p. 326) proved that after the first half-minute of the inhalation of chloroform there is a progressive lowering of the arterial pressure.

It would, *a priori*, appear probable that this is to some extent due to a vaso-motor paralysis; but Sansom and Harley state that there is a spasm of

the small vessels, which can be readily seen to occur in the web of the frog during chloroformization. Not until the third stage is reached, according to these authors, do the vessels relax into dilatation. If these observations are correct, chloroform probably acts upon the vaso-motor centres as upon those of the vagi.

On the heart itself chloroform undoubtedly exerts a steady, powerful, depressing influence.

As long ago as 1842, Glover (*Edinburgh Medical Journal*, 1842) discovered that chloroform injected into the jugular vein instantly arrests the heart's action and destroys its muscular irritability. This has been confirmed by Gosselin (*Archives Générales*, 1848), Anstie (*Stimulants and Narcotics*, p. 321), and other observers. Moreover, it has been found that the vapor of chloroform exerts a similar influence when locally applied to the exposed heart (*Edinburgh Medical Journal*, 1842). It is, therefore, a demonstrated fact that chloroform applied in a sufficiently concentrated form instantly *destroys the contractile power of the heart-muscle*.

The respirations may at first be rendered slower by chloroform, but after a time are generally quickened, and as the inhalation is persisted in they become more and more shallow, irregular, and distant, and finally cease.

In animals, without doubt, chloroform, when inhaled, usually produces death by abolishing the functional power of the medulla oblongata and thereby arresting respiration. It has been denied that chloroform-vapor ever kills animals in any other way.* This is, however, a mistake.† If the chloroform-vapor be administered in a very dilute form, the heart generally continues to beat for a long time after the cessation of respiration. If the vapor have been less diluted, cardiac and respiratory action may cease almost simultaneously. If the vapor have been administered in a concentrated form, arrest of the heart's beat may precede arrest of respiration.

When death is produced by chloroform in man, it is generally by cardiac paralysis due to a direct action upon the heart-muscle, rarely by asphyxia.

Death by syncope occurs proportionately more frequently in man than in animals, because this form of death is sudden and unexpected. Asphyxia coming on gradually, the threatening symptoms are perceived, and death is averted by prompt measures. In animals, however, the object being to kill, nature is allowed to take its course.

Various other theories besides that adopted here have been brought forward to account for the chloroform deaths. I think they are unnecessary and unproven, although some are ingenious, and it is possible that death may take

* See, especially, report of the French Chloroform Commission. Sabarth, *Das Chloroform*, p. 44.

† Consult Snow, *On Anæsthetics*, p. 111; Anstie, *Stimulants and Narcotics*; Richardson, *Medical Times and Gazette*, 1870; Chloroform Committee, *Medico-Chirurgical Transactions*, vol. xlvii.

place at times in the way they suggest. Want of space prevents my doing more than giving references to the most important papers.*

According to Harley (*Philosophical Transactions*, London, 1865), blood to which as little as five per cent. of chloroform has been added becomes very liquid and of a bright arterial hue. After a time crystals form in it. Boettcher (*Virchow's Archiv*, vol. xxxii. p. 126) was, I believe, the first to study these changes closely. The first alteration noticeable in the red blood-disks is a diminution of their size, which A. Schmidt and F. Schweiger-Seidel (*Berichte d. könig. sachs. Gesellsch. d. Wissensch., math.-phys. Kl.*, 1867, p. 190) assert to be due to contraction, because when blood is treated with water until the red globules disappear, and carbonic acid gas is passed through the liquid until they reappear, on the addition of chloroform the sharply-contoured bodies will be seen to undergo marked contraction. As was first shown by Boettcher (*loc. cit.*, p. 127), and confirmed by Schmidt and Schweiger-Seidel, chloroform alone produces no other alteration than contraction in the red blood-disks. If, however, air be admitted to blood containing chloroform, the corpuscles rapidly disappear, dissolving in the serum, out of which, after a time, hæmatin crystallizes. Both of the authorities quoted believe that the latter changes are due to oxidation.

Boettcher states (*loc. cit.*, p. 129) that chloroform-vapor mixed with air converts enough of the oxygen of the latter into ozone to react with iodinated-starch paper; and Schmidt and Schweiger-Seidel have found that an excess of carbonic acid in the blood interferes with the changes caused by chloroform. From these facts, it seems probable that their opinion as to the nature of the blood-changes is correct.

Harley (*loc. cit.*) has studied the effect of chloroform on the absorption of gases in the blood. He states that when chloroform is added to fresh blood, and the mixture allowed to stand for twenty-four hours, a marked increase takes place in the proportion of oxygen and a lessening in that of carbonic acid. This is in accord with the theory just mentioned; for, after the complete oxidation of the hæmato-globulin brought about by the chloroform, further consumption of oxygen could not occur, and, as it continued to be absorbed from the air, it would accumulate instead of being converted into carbonic acid.

How far, during ordinary narcosis, chloroform produces the changes just described in the blood, is somewhat uncertain; but it would seem very improbable that they occur to any great extent. A very sensitive test of the destruction of the red disks in the body is found in the production of icterus, and icterus never follows anæsthesia.

On the other hand, Husemann (*Schmidt's Jahrbücher*, Bd. cli. p. 84)

* Richardson, *Medical Times and Gazette*, 1870. Andrew H. Smith, *New York Medical Journal*, 1871, p. 46. Consult also J. C. Reeve, *American Journal of the Medical Sciences*, Oct. 1867; Henry M. Gibbons, Jr., *Pacific Medical and Surgical Journal*, June, 1869; Snow, *On Anæsthetics*; and Sanson, *On Chloroform*.

intimates, on what authority I do not know, that after anæsthesia bile-acids appear in the urine; and Bert (*Journal of Anatomy and Physiology*, May, 1870) has found that the oxygen of the blood undergoes during anæsthesia an increase, such as Harley has found to happen when chloroform is added to blood outside of the body.

THERAPEUTICS.—As an anæsthetic, chloroform possesses the advantages of quickness and pleasantness of operation, smallness of dose, and cheapness. These advantages are, however, so outbalanced by the dangers which attend its use, that its employment under ordinary circumstances is unjustifiable. It kills without warning, so suddenly that no forethought or skill or care can guard against the fatal result. It kills the robust, the weak, the well, the diseased, alike, and the previous safe passage through one or more inhalations is no guarantee against its lethal action. Statistics seem to indicate a mortality of about one in three thousand inhalations;* and hundreds of utterly unnecessary deaths have now been produced by the extraordinary persistence in its use by a portion of the profession.

It ought, therefore, never to be employed except under especial circumstances, as in some cases of *puerperal eclampsia* when a speedy action is desired, or in the field during war-time, when the bulkier anæsthetics cannot be transported.

In obstetric cases, chloroform has been used even by those who give the preference to ether in surgery. So far as I know, no death has as yet occurred from chloroform during *parturition*, although alarming effects have been induced. The excitement of child-birth does seem to fortify the system against the deleterious influence of chloroform. But even in these cases I think ether is just as useful as, and much safer than, its sister-anæsthetic.

Various mechanical inventions have been made for the administration of chloroform; but these inhalers do not appear to offer any advantages over the simple napkin, and, at least in this country, are rarely, if ever, used.† A handkerchief or towel may be folded into a bird's-nest shape, and twenty or thirty drops of the anæsthetic be put upon this and then held close to the mouth. Dr. Simpson advises that a towel be laid over the mouth and nose, and the chloroform slowly dropped upon this until anæsthesia is induced. Whatever plan be employed, it is of vital moment that the vapor be well diluted: not more than three and one-half per cent. of it should be contained in the inspired air.

When administered by the mouth in sufficient quantity, chloroform produces symptoms similar to but much more permanent than those which it causes when inhaled. It is, however, very rarely, if ever, used in this way for its constitutional effect.

* See Richardson, *Medical Times and Gazette*, 1870; Henry M. Gibbons, Jr., *Pacific Medical Journal*, June, 1869; Squibb, *On Anæsthetics*, *New York Medical Journal*, April, 1871.

† For a description of these inhalers, see the works of Sansom and Snow.

When chloroform is taken into the stomach, a considerable portion of it is, without doubt, evaporated, so that the intestinal canal becomes filled with the vapor. Chloroform, therefore, when so placed exerts both a local anodyne and a stimulant carminative action. For this reason it is extremely valuable in all cases of *colic*, and it will often even assuage the pain of *colica pictorum*.

Externally, as a rubefacient and anodyne, chloroform is very largely combined with other substances into liniments, which are especially useful in cases of *chronic neuralgic* or *rheumatic pains*.

TOXICOLOGY.—Death may occur at any time during the inhalation of chloroform. In some cases it has seemed to be instantaneous, but generally it is preceded by symptoms such as a change in the expression of the face, which becomes very pale or livid, evident irregularity of the respiration, and failure of the pulse: a shudder, or a violent convulsion, or even a sudden access of maniacal excitement, has been noted in some cases. After death the heart is almost always found relaxed, distended, and its right side filled with blood.

When death is threatened through asphyxia, the alternate dashing of very cold and very hot water upon the face and upper chest is often very efficacious. Artificial respiration should be commenced at once. Faradization of the diaphragm, by pressing one pole firmly against the pit of the stomach and placing the other over the larynx and the root of the neck, has acted very favorably in some cases. Dr. Richardson has shown (*loc. cit.*) that there is a danger of the excitability of the muscle being exhausted by a prolonged use of the galvanic stimulus. The application should not, therefore, be long continued, but should be alternated with artificial respiration. Whenever there is any failure of the heart's action, as is nearly always the case, the body should be laid at an angle of 40°, with the head downwards, so as to favor the passage of arterialized blood to the brain (Dr. E. L. Holmes, *Chicago Medical Journal*, Sept. 1868).

In artificial respiration, act with the patient, and not against him. He will not cease to breathe at once, and wholly. Enjoin silence; watch the first attempt at inspiration, and at the expiration compress the thorax, aiding its elastic reaction, if absolutely necessary, by Sylvester's or some other quiet method. See that the tongue is well forward.

Dry external heat must be vigorously used, and the inspired air should be at from 80° to 85° F., or even higher. Frictions and passive motion, to aid in the circulation, must not be forgotten. Efforts at resuscitation should be kept up for at least two hours.

Poisoning has been produced by the swallowing of chloroform. The symptoms induced have been stupor, with contracted, or, in later stages, dilated, pupils, and a stertorous respiration, which finally becomes very irregular, shallow, and often distant. The amount necessary to destroy life probably varies very much. Recovery has occurred after the ingestion of two ounces (Stillé, *Therapeutics*, vol. ii. p. 107). The treatment consists in the use of the

stomach-pump, of the various ordinary methods of arousing a narcotized patient, especially the alternate cold and hot douche, artificial respiration, and the very cautious use of diffusible stimuli if required. Death may occur during the narcosis, or the patient may survive this, and perish from inflammation of the trachea, œsophagus, and stomach, caused by the local action of the chloroform.

The recognition of chloroform as the probable cause of any given death cannot be based upon the post-mortem appearances. Indeed, the latter are of no value in deciding such a question. The anæsthetic may, however, be recovered by distillation of the lungs and blood within a certain period of time after death. As to the length of this time, so far as I know, no investigations have been made.

Experiments made at the Philadelphia Hospital have proven that persons sound asleep may be chloroformed without their being awakened. Anæsthesia cannot, however, be produced in any one partially awake, or even sleeping lightly, without his knowledge.

Quite a number of professional men have been accused, and some convicted on the charge, of committing rape on females in whom they were inducing anæsthesia. The women, no doubt, believed that they had been violated; but it is very certain that in many, and probably all, of the cases, they mistook the subjective erotic sensations induced by the chloroform or ether, for the real act. The valuelessness of the testimony of persons as to occurrences during the time of their intoxication with anæsthetics should be recognized by law as a governing principle of evidence.

BICHLORIDE OF METHYLENE.

This is a colorless liquid, having a chloroform-like odor, a specific gravity of 1.344, and boiling at 88° F. Its reaction, when pure, is neutral to test-paper. It was introduced to the notice of the profession by Dr. B. W. Richardson (*Medical Times and Gazette*, 1867, p. 478) as an anæsthetic similar to, but more pleasant and possibly safer than, chloroform, and has been pretty extensively used in London. It has never, that I am aware of, been employed in this country. Dr. Richardson, with rare control, expressed no opinion as to the real safety of the new anæsthetic, and subsequent events have justified him. There is no way of knowing how many times it has been employed, but certainly three deaths dependent upon or coincident with its use have been reported (*Medical Times and Gazette*, 1869, ii. p. 524, and *British Medical Journal*, Sept. 1871, and August, 1872). The recorded phenomena in some of these cases indicate that, like chloroform, the bichloride of methylene kills by paralyzing the heart. It is scarcely probable that it will ever come into general use as an anæsthetic.

CLASS IX.—EXCITO-MOTORS.

In this class are included such drugs as increase the reflex activity of the spinal centres, and thereby give rise to disturbance of motility. The only representatives of the class used by the practitioner of medicine are those drugs which contain strychnia as their active principle.

NUX VOMICA—NUX VOMICA.

The seeds of *Strychnos nux vomica*, a middle-sized tree growing in the East Indies, whence the drug enters commerce. They are circular, nearly flat disks, a little less than an inch in diameter, covered with very short, satin-like, grayish hairs; internally they are tough and horny, and are possessed of an intensely bitter taste. They contain two alkaloids,—strychnia and brucia,—existing in combination with an acid, the *igasuric* of Pelletier and Caventou, which, according to Husemann, is probably identical with malic acid.

Brucia, which, unlike strychnia, is not officinal, is readily recognized by the following test. When concentrated nitric acid is added to it, a beautiful scarlet or blood-red color is developed, which becomes yellowish red, and, by warming, yellow; if to this yellow solution somewhat diluted some chloride of tin or sulphuret of ammonium be added, it becomes a beautiful reddish violet. Physiologically and therapeutically this alkaloid is similar to but much weaker than strychnia.

An *extract of nux vomica* (*Extractum Nucis Vomicae*) is officinal. The dose of it is from one-fourth to one-half of a grain, given in pill. The U. S. Pharmacopœia also contains a *tincture* (*Tinctura Nucis Vomicae*, ℞iv-Oj), the dose of which is ten to twenty drops.

STRYCHNIA.

As kept in the shops, strychnia is a grayish-white powder, but by slow crystallization from its alcoholic solution it may be obtained in octahedral or in quadrilateral prisms. It is soluble in about seven thousand parts of cold water; two thousand parts of boiling water; very sparingly soluble in absolute alcohol, ether, and benzine; freely soluble in boiling officinal alcohol, which deposits it on cooling. It is so bitter that it will impart a very intense taste to twenty thousand times its weight of water.

Strychnia yields a very pronounced violet color with many oxidizing reagents. The one most ordinarily employed is a mixture of concentrated sulphuric acid and bichromate of potassium (*Otto's test*). According to Dr. Guy, the test is most delicate if the alkaloid be dissolved in a little concentrated sulphuric acid on a plate, and the bichromate added to it, when a bluish and then violet-purplish color is developed, passing finally into a dirty green. *Davy's test* consists in the substitution of a crystal of red prussiate of potash for the bichromate; *Marchand* uses the peroxide of lead; in this case the sulphuric acid should contain one per cent. of nitric acid. Drs. Vrij and Van der Burg say that these tests are about equally sensitive, and are capable of revealing the one-sixty-thousandth of a grain of the alkaloid. Either the chlorate or the permanganate of potassium may be used instead of the bichromate; and, indeed, Dr. Guy claims that the permanganate is preferable to the latter. If the strychnia be in quantity, it may be dissolved in very dilute sulphuric acid, and solution of bichromate of potassium be added, when golden-red needle-like crystals of the chromate of strychnia will separate. These dissolve, with the production of a beautiful blue color, in concentrated sulphuric acid. F. L. Sounnenschein (*Vierteljahresschrift für Prakt. Pharmacie*, 1871) says that if strychnia be dissolved in a strong solution of the sulphate of sesquioxide of cerium a beautiful color is induced, which generally passes into a cherry-red, and so persists for several days. Dr. Filchol's test (*London Lancet*, April, 1872) consists in the addition of solution of chloride of gold, and the testing of the precipitate by Otto's method. The *physiological test* for strychnia is a very sensitive one. In it a fragment of the suspected extract, dissolved in a little acidulated water, is thrown into the cellular tissue of a small frog, which should afterwards be allowed to swim about freely, so that its unconstrained movements can be watched.

PHYSIOLOGICAL ACTION.—Strychnia acts in the same way upon almost all animals. According to Leube (*Reichert's Archiv für Anatomie*, 1867, p. 630), however, it takes ten times as much to kill chickens as other birds, weight for weight; and among mammals the guinea-pig is very insensitive to it. It has also recently been asserted that on some monkeys it has very little influence (*Boston Medical and Surgical Journal*, 1872). Its local action is that of a slight irritant.

When taken in quantities just sufficient to produce sensible physiological effects, strychnia induces a feeling of restlessness, perhaps accompanied by tremblings in the limbs and some stiffness in the neck and jaws. When a somewhat larger amount has been given, there may be general muscular twitchings and startings, with stiffness and stricture of the throat and chest; formications or other abnormal sensations under the skin may or may not be present.

After poisonous doses the symptoms come on usually in from fifteen to twenty minutes, rarely after the hour, with great suddenness; sometimes the convulsions are preceded by partial spasms of the muscles of the extremities,

but more often the patient is suddenly thrown down by a general tetanic spasm. In this the body is bent backwards and rests upon the heels and the head, in a condition of profound opisthotonos; the legs are rigidly extended and the feet everted; the arms bent and the hands clenched; the eyes staring, wide open; the corners of the mouth often drawn up so as to produce the *risus sardonicus*. The senses are often sharpened, but ringing in the ears and dimness of vision may be induced, if the fits are severe. The face is at first pale, but, if the fit is sufficiently severe and be protracted, it becomes livid from the interference with respiration. Consciousness is not affected, unless when asphyxia becomes so pronounced as to threaten death; in such cases *sometimes* a period of insensibility precedes dissolution, but generally the intellect is clear to the moment of death. The muscles of the jaw are usually the last in the body to be affected, but trismus finally comes on in severe cases. I have seen death occur in this first convulsion in animals, but Tardieu states that he knows of no such instance in man (compare case of Dr. Demme, *Syd. Soc. Year-Book*, 1865-66, p. 441). After a time the paroxysm is at an end, the jaw drops, the muscles relax, and a period of calm comes on, to be succeeded by a second convulsion like the first.

These convulsions are excited by the slightest touch, by a draught or breath of air, even by a loud sound; but a firm grasp or hard rubbing of the muscles is frequently grateful. A slight rigidity is sometimes manifest between the paroxysms, but no marked stiffness.

If the case terminate favorably, the convulsions gradually lessen in intensity, and fade away, leaving the patient exhausted, with a sore, tired feeling in the muscles. The spasms are generally, but not always, very painful. There are often erections of the penis, and the fæces and urine may be passed involuntarily.

After death, post-mortem rigidity is developed very quickly. Autopsies have revealed nothing but the usual congestive lesions of death from asphyxia, and, at times, indications of spinal hyperæmia.

In regard to the method in which strychnia produces the above symptoms, it is obvious that the alkaloid is primarily absorbed; and experimental proof seems so superfluous that I will only mention the fact that Masing has found the strychnia in the blood.

It is very plain that convulsions can be produced by a drug in only five ways: first, they may be epileptiform,—*i.e.*, cerebral; second, they may conceivably be due to a stimulation of the peripheral ends of the motor nerves; third, they may conceivably be caused by irritation of the peripheral ends of the sensory nerves; fourth, they may be muscular,—*i.e.*, due to a direct action on the muscles; fifth, they may be spinal.

That the convulsions of strychnia-poisoning are not cerebral, is proven by the fact which has been frequently noted, and which I have confirmed, that they are not affected by section of the cord, or, at least, are only so far affected as to be more severe in those portions of the body removed from the cerebral influence.

That they are not due to irritation of the peripheral motor nerves and are not muscular is proven by the experiment of Valentin (*Pathologie der Nerven*, p. 327, Leipsic, 1864), who found that the injection of a solution of strychnia through the blood-vessels of the amputated leg of a frog had no influence upon the muscles. A very beautiful experiment of Brown-Séquard (*Comptes-Rendus*, 1849) confirms this, and also demonstrates that the convulsions do not arise from hyper-excitability of the peripheral afferent nerves. The observer last mentioned found that when the spinal cord was cut just below the origin of the nerves supplying the fore legs of a frog, and all the blood-vessels going to the lower section of the cord were also severed so as to isolate the latter, on the exhibition of strychnia convulsions occurred in the anterior part of the body, whilst in the posterior segment quiet and a normal reflex activity were maintained, although the blood was carrying the poison to every part of it except the spinal cord. The philosophy of this is evident. The anterior section of the cord, receiving the poison, gave rise to convulsions; the posterior section, receiving no poison, maintained its usual status.

This experiment of Brown-Séquard has been repeated a great number of times by MM. Martin-Magron and Buisson (*Journal de la Physiologie*, t. iii., 1860, p. 130) with similar results, excepting that in some very rare instances slight convulsions were induced in the posterior portion of the body. These exceptional phenomena appear to have been due simply to a minute portion of the poison reaching the spine by inhibition, since, when by an operative procedure not necessary here to detail (*loc. cit.*, p. 131) the posterior section of the cord was completely isolated, and access of the poison by diffusion rendered impossible, spasms never occurred in the posterior part of the body.

Strychnic convulsions must be spinal, because they do not arise in any of the other possible methods. This conclusion is abundantly confirmed by direct experiment. Thus, Van Deen (*Physiologie de la Moelle épinière*), and also Valentin (*loc. cit.*, p. 329), have shown that when the alkaloid is placed upon the spinal cord, and allowed slowly to diffuse itself, the usual convulsions occur, but are at first confined to those muscles whose nerves have their origin near the point of application, and afterwards spread from muscle to muscle as the poison creeps through the cord. Dr. A. J. Spence (*Edinburgh Medical Journal*, July, 1866) has performed similar experiments, with similar results. He first bisected the apex of a frog's heart so as to allow all the blood to drain from the body, and then, cutting through the cranium, laid a little piece of nuxvomica within it upon the brain so that the poison would diffuse down through the spinal cord. The result was that first the muscles of the throat, then those of the fore legs, and so on in regular order, were affected.*

* Some of the phenomena stated by Dr. Spence to have occurred are at present very difficult to explain. Thus, he noted that, as the poison traveled down the cord, there was a time when irritation of the fore feet caused only spasms in them; later in the experiment, irritation of the front feet caused spasm both of the front and hind feet, although irritation of the latter did not produce other than normal reflex movements;

Claude Bernard (*Leçons sur les Substances toxiques*, Paris) has denied that strychnia produces excitation of the spinal motor centres, because when all of the posterior nerve-roots are cut, no convulsions occur, whereas if a single afferent root be allowed to remain, irritation of its peripheral fibrillæ causes general tetanic spasms. Allowing the truth of his experimental fact, his deduction certainly is not warrantable. The non-occurrence of convulsions may depend upon the fact that the reflex motor ganglionic cells are incapable of originating an impulse, and in strychnia-poisoning are simply in such a condition of over-excitability as renders them exceedingly sensitive to slight irritations, and causes them to respond most energetically to peripheral impulse so feeble as not to be felt in health. That the motor centres are acted upon by strychnia is proven not only by the experiments that have been already detailed, but by the following ingenious one of Van Deen (*Physiologie de la Moelle épinière*). That investigator removed the viscera, vessels, etc., from a frog, so as to leave nothing below the second cervical vertebra but the bones, nerves, and muscles; then, opening the spinal cord in the region of the third vertebra, he cut entirely through the anterior columns of the cord, and finally divided all the tissues, so that the anterior portion of the frog was connected with the posterior solely by the posterior columns of the cord. When one or two drops of a solution of strychnia were placed in the mouth of the prepared batrachian, tetanus, confined to the anterior segment of the body, was developed; and it was also found that whilst irritation of the posterior feet caused in them only ordinary reflex movements, in the front legs tetanic spasms were simultaneously induced.

The action of strychnia upon the motor nerves has been a subject of considerable controversy. That the convulsions occur independently of any such influence, if it exist, has been already shown. After death from strychnia, the functions of the motor nerves are always found to be more or less destroyed, so that galvanization of the nerve-trunk either produces only very feeble contractions in the tributary muscles, or else none at all. Of this fact there can be no doubt; it has been attested on the evidence of personal experiment by many observers, among whom may be mentioned Matteucci (*Traité des Phénomènes électro-physiologiques*, Paris, 1844), Moureau (*Comptes-Rendus de la Société de Biologie*, 1855), M. Ambrosoli (*Gazette Médicale*, 1857, p. 525), M. Wittich (*Bericht über die Fortschritte der Anatomie*, 1857, p. 434), and Kölliker (*Virchow's Archiv*, Bd. x., 1856).

Now, it is evident that this absence of response may be due to loss of functional power in either muscle or nerve. Sometimes the muscle may be at fault; but, as Matteucci (*loc. cit.*) insists, and as has been noted by many observers, not rarely, indeed most generally, in the frog galvanization

later still in the poisoning came a stage when irritation of the front legs was powerless to cause spasm in the hind legs, although irritation of the latter would now cause spasm in the former.

of the nerve fails to elicit response, although the muscle preserves its irritability.

It having been proven that the functional power of the motor nerves is destroyed in strychnia-poisoning, the question arises, Is this destruction a direct action of the poison, or is it simply the exhaustion of over-use, due to the intense activity of the nerve during the stage of spasm?

It cannot be gainsaid that the power of the nerve is lessened by the strain upon it during the convulsions; and Kölliker concludes (*Virchow's Archiv*, Bd. x., 1856) that this is the sole cause of the nerve-paralysis, because when he cut the sciatic nerve of a frog and exhibited strychnia the divided nerve would respond to galvanic stimuli after all functional power had been lost in the nerve whose connection with the centres was intact. Granting the experimental fact, this conclusion of Kölliker is apparently inevitable, since all the nerves suffered equal contact with the poison. The conclusion is, however, opposed to a fact proven by Vulpian (*Archives de Physiologie*, 1870, t. iii. p. 120) and other observers, that an enormous dose of strychnia kills the frog without the induction of spasms by general paralysis, with total loss of power in the nerve-trunks. Evidently in such case the action of the poison on the nerves must be direct.

MM. Martin-Magron and Buisson have investigated (*Journal de la Physiologie*, 1860, t. iii. p. 342) very elaborately this action of strychnia upon the nerves, and the correctness of their experiments and results seems to me scarcely questionable. They found (*loc. cit.*, p. 347) that if the sciatic nerve of a frog were cut and a sufficiently large dose of the strychnia administered, the divided sciatic lost its irritability, although, unless the dose was very large, not so soon as did its fellow. Vulpian (*loc. cit.*, p. 126) has confirmed this, and the opposite result of Kölliker no doubt depended upon his not using sufficiently large doses of the alkaloid or upon his testing the nerve too early. Martin-Magron and Buisson also tied all the tissues of a frog's leg except the nerve, and then, on exhibiting strychnia, found that convulsions ceased in the poisoned much sooner than in the non-poisoned leg, and that at a certain time irritations of the poisoned foot would induce tetanic spasms *only* in the *non-poisoned member*,—proof that the afferent nerve-fibres of the poisoned legs were not affected, and that the motor nerves were paralyzed wherever the poison had access to them; and that to this, not to spinal exhaustion, was due the general paralysis.

M. Vulpian (*loc. cit.*, p. 121) affirms that he has repeated this experiment many times, and always obtained the same result as Martin-Magron and Buisson. It seems to me absolutely to prove that strychnia in very large doses paralyzes the efferent, but not the afferent, nerves; and that the collapse of strychnia-poisoning in the frog is largely due to the affection of the motor trunks, and not to exhaustion of the spinal cord. As Fraser has discovered for atropia, Vulpian (*loc. cit.*, p. 128) has found for strychnia: namely, that after a time,—say from some hours to two days,—if the dose has been of the

right size, the strychnic paralysis passes off, the motor nerves are found to have regained their power, and the convulsions reappear and continue hours or days.

Our knowledge of the action of strychnia upon the circulation is very incomplete.

According to Dr. Sigmund Mayer, Richter (*Zeitschrift f. rationelle Medicin*, 1863, Bd. xviii.) has observed that strychnia causes a very decided rise in the arterial pressure, and also a microscopically visible contraction of the vessels of the frog's web.

Mayer himself has furnished the fullest study which we have (*Medicinische Jahrbücher der k. k. Gesellschaft der Aerzte zu Wien*, 1872, p. 112) of the relation of strychnia to the circulation.

In his experiments, he has found that about the time tetanus commences in strychnized animals an enormous rise of the arterial pressure takes place. That this rise is not due simply to the convulsions, is proven by its occurrence in animals kept quiet by curare. In normal strychnized animals, the pulse-rate increases simultaneously with the arterial pressure; but that the latter is not dependent upon the former is shown by the circumstance that when in the poisoned animal the heart is caused to beat slowly by strong stimulation of the vagi, the arterial pressure still remains far above normal. Mayer confirms the observation of Richter that the small arteries can be seen to contract in strychnia-poisoning, and concludes that the rise of arterial pressure is due to a general vaso-motor spasm.

The centric origin of the spasm is demonstrated by the fact that, after section of the cord high up,—i.e., after separation of the vessels from the vaso-motor centres,—strychnia produces no rise in the blood-pressure, or a very slight one. If these experiments of Mayer be confirmed, he has proven that strychnia acts on the vaso-motor centres as upon the other motor centres.

There is some apparent conflict of testimony in regard to the influence of strychnia upon the inhibitory cardiac nervous system. Dr. Carl Heinemann, who has investigated (*Virchow's Archiv*, Bd. xxxiii. p. 394) at some length the influence of the drug upon the heart of the frog, finds that large doses cause diminished frequency of the cardiac movements, with diastolic pauses. According to his experiments, these phenomena are not due to stimulation of the inhibitory nerves, since they occur after section of the vagi (p. 403); nor are the peripheral vagi paralyzed, since galvanization of one of these nerves causes immediate diastolic cardiac arrest (p. 406).

Mayer (*loc. cit.*) has also found that the peripheral vagus is not paralyzed, since he could suspend the action of the heart in the poisoned animal by galvanization of the par vagum. On the other hand, Martin-Magron and Buisson affirm (*loc. cit.*, p. 352) that, in all of very many experiments, after a greater or less length of time the pneumogastrics lost their power of transmitting an impulse. I see only one way of reconciling these results,—i.e., by the supposition that strychnia does affect the peripheral vagi, but not until very late in the poisoning; and that, perhaps, when small doses are

employed, even death may occur without very decided paralysis of these nerves. That the different results are not due to the use of different animals is evident, since, although Mayer used hounds, Heinemann employed the same animals as did Martin-Magron and Buisson, viz., frogs.

Upon the *blood* strychnia has probably some action. Harley found that blood shaken for twenty-four hours with air contained 11.33 parts of oxygen and 5.96 parts of carbonic acid; whilst blood treated in a precisely similar manner, excepting in the addition of strychnia, yielded 17.80 parts of oxygen and 2.73 parts of carbonic acid. He concludes from this that strychnia arrests oxidation in the body; but the deduction seems to me out of all proportion to the fact.

THERAPEUTICS.—As a vegetable bitter, strychnia is, of course, a tonic, stimulating to a greater or less degree the digestion, besides acting more universally on nerve-power. But it is more than a mere stomachic: clinical experience has shown that it is a most useful tonic when there is general relaxation and loss of nerve-power. A portion of its value arises, it may be, from its action upon the spinal motor nerve-centres; but in all probability it influences other portions of the cord, affecting the vaso-motor centres, and most probably also the trophic centres, if they exist. Be these things as they may, clinical experience has abundantly demonstrated the value of the drug as a tonic in general *functional atony and relaxation*.

The great influence of strychnia upon the function of voluntary motion early led to its use in cases of *paralysis*, often with the result of doing harm rather than good. Its peculiar physiological action being known, it becomes very evident that it can be useful only when the paralysis is dependent upon, or at least accompanied by, a *depressed state* of the *spinal* or other *motor centres*. Whenever there is *inflammation or irritation of these latter*, strychnia may do great injury by increasing such irritation, and must never be employed. Like galvanism in *hemiplegia*, it can do only a very limited amount of good, and should not be exhibited until irritation from the clot has ceased. Andral and others commend it, especially in *lead-paralysis*.

In *amaurosis* it has been very strongly recommended by some and equally condemned by others; but the weight of the testimony seems to me strongly in its favor when the affection depends simply upon retinal atony, especially when there is no decided organic disease of the optic nerve, but only a loss of power from exhaustion or previous injury. It was originally used endermically by Dr. Middlemore in these cases; but in 1872 Prof. Nagel, of Tübingen, introduced the employment of it hypodermically in the supra-orbital region (*Boston Medical and Surgical Journal*, April, 1872), and Prof. J. J. Chisolm claims that he has obtained most brilliant results by following this practice (*American Journal of the Medical Sciences*, Jan. 1872).* In carrying out this plan, one-sixtieth of a grain, rapidly increased to one-thirtieth, should be

* See also paper by C. W. J. Riching, *British Medical Journal*, 1873.

thrown twice a day into the supra-orbital region. When good results follow, they appear to be manifested very soon.

In *dyspepsia* or *constipation* or *diarrhœa*, connected with atony of the visceral muscular coat, strychnia is a very valuable remedy. In various local paralyses, such as *prolapsus of the rectum*, *atonic retention of urine*, *atonic incontinence*, and *loss of voluntary motion* in certain groups of muscles from temporary injury to the supplying nerve, it is useful. In these cases it should be injected into the affected part.

TOXICOLOGY.—Sufficient has already been said in regard to the general symptoms of strychnia-poisoning. It only remains to discuss the diagnosis. This is especially important, because strychnia is frequently used criminally, and because it is not rarely impossible for the chemist to detect it after death. The only disease with which poisoning by it may be readily confounded is tetanus, in its various forms of idiopathic, rheumatic, traumatic, infantile, and hysterical. It has been asserted that in fatal cases the duration of the attack will always distinguish between natural tetanus and that produced by poison. Dr. Louis Starr, however (*Philadelphia Medical Times*, vol. iii. p. 311), reports a case of traumatic tetanus fatal in less than twelve hours after the first appearance of muscular twitchings, and within one hour and a half after the first convulsion.

The following table shows, I think, in as clear and brief a manner as possible the differences* between traumatic or idiopathic tetanus (No. 1), hysterical tetanus (No. 2), and strychnic poisoning (No. 3). The references in column No. 3 are to authorities who affirm that the symptoms there given are peculiar to poisoning.

No. 1.	No. 2.	No. 3.
	Commenced with blindness and weakness.	Begins with exhalation and restlessness, the special senses being usually much sharpened.† Dimness of vision may in some cases be manifested later, after the development of other symptoms; but even then is rare.
Muscular symptoms usually commence with pain and stiffness of the back of the neck, sometimes with slight muscular twitchings; come on gradually.	Muscular symptoms commenced with rigidity of the neck, which gradually "crept over the body," affecting the extremities last.	Muscular symptoms develop very rapidly, commencing in the extremities, or the convulsion, when the dose is large, seizes the whole body simultaneously.‡
Jaw one of the earliest parts affected; rigidly and persistently set.	Jaw rigidly set before a convulsion, and remained so between the paroxysms.	Jaw the last part of the body to be affected; its muscles relax first, and even when during a severe convulsion it is set, it drops as soon as the latter ceases.§

* Column No. 2 is from an actual case. See Trial of Mrs. Wharton, *New York Medical Record*, 1873.

† Taylor, *On Poisons*, p. 683. Wormley, *Micro-Chemistry of Poisons*, p. 536.

‡ Wormley, p. 536. Stillé, *Therapeutics*, vol. ii. p. 148.

§ Taylor, *On Poisons*, pp. 134, 682. *Medical Jurisprudence*, pp. 540, 541, 536. Tardieu, p. 924.

No. 1.	No. 2.	No. 3.
Persistent muscular rigidity, very generally with a greater or less degree of permanent opisthotonos, emprosthotonos, pleurosthotonos, or orthotonos.	Persistent opisthotonos, and intense rigidity between the convulsions; and after the convulsions had ceased the opisthotonos and intense rigidity lasted for hours.	Muscular relaxation (rarely a slight rigidity) between the convulsions, the patient being exhausted and sweating. If recovery occur, the convulsions gradually cease, leaving merely muscular soreness, and sometimes stiffness like that felt after violent exercise.*
Consciousness preserved until near death, as in strychnic poisoning.	Consciousness lost as the second convulsion came on, and lost with every other convulsion, the disturbance of consciousness and motility being simultaneous.	Consciousness always preserved during convulsions, except when the latter become so intense that death is imminent from suffocation, in which case sometimes the patient becomes insensible from asphyxia;† which comes on during the latter part of a convulsion, and is almost a certain precursor of death.
Draughts, loud noises, etc., produce convulsions as in strychnic poisoning.	Desired to be fanned.	The slightest "breath of air" produces a convulsion.‡
May complain bitterly of pain.	Crying-spells, in which he "sobbed violently," and "cried like a child," alternated with the convulsions.	Patient may scream with pain, or may express great apprehensions, but "crying-spells" would appear to be impossible.
Eyes open, rigidly fixed, during the convulsion.	Eyes closed.	Eyes stretched wide open.§
	The spasms in leg must have been partial, as the feet were crossed and toes inverted, which could not happen if all the muscles were involved, because the muscles of eversion, being very much the stronger, would of necessity overcome the antagonistic muscles, and the feet be everted.	Legs stiffly extended, with feet everted, as the spasms affect all the muscles of the leg.

Death from strychnia in man and other mammals mostly occurs in a convulsion, and under these circumstances is undoubtedly due to asphyxia, caused by the unyielding spasmodically-contracted muscles. In frogs, death must occur from other causes, since a frog, as shown by Claude Bernard, will live for days after removal of its lungs, probably by breathing through its skin. The causes of death in the frog are not hard to find when the physiological action of the drug is known. The lymph and true heart (Kölliker, *loc. cit.*; Harley, *London Lancet*, July, 1856) are very much affected, but the chief factor is no doubt paralysis of the motor nerves. In man, death sometimes

* Taylor, *On Poisons*, pp. 134, 136, 682. Wormley, pp. 536, 540, 541. Tardieu, *Sur l'Empoisonnement*, pp. 924, 938, 939. Husemann, *Handbuch der Toxicologie*, p. 168.

† Wormley, p. 536. Taylor, *Medical Jurisprudence*, pp. 331, 332. Wharton and Stillé, *Medical Jurisprudence*, paragraph 757. Tardieu, p. 923. Stillé, *Therapeutics*, p. 148.

‡ Stillé, *Therapeutics*, p. 148.

§ Stillé, *Therapeutics*, p. 148. Wormley, p. 536. Tardieu, p. 924.

|| Tardieu, p. 924; also other authorities, which I have neglected to note, and at present writing have not at hand.

occurs not in a paroxysm, but during relaxation, and probably then is the result not only of the exhaustion following effort, but also of the direct action of the poison on the nerves.

The minimum fatal dose of strychnia is probably something under half a grain; the latter quantity has several times produced death, once in a man in twenty minutes (*Guy's Hospital Reports*, 1865, vol. xi. p. 208); one-third of a grain given at intervals has produced such alarming symptoms as to indicate that in a single dose it might readily destroy life.

On the other hand, Tschepke (*Deutsch. Klinik*, 1861) details a case in which ten grains of the nitrate of strychnia were taken, and in which vomiting did not come on for over two hours, and yet recovery occurred. As the patient's stomach was well filled with berries containing tannic acid, it is probable that but a modicum of the poison was absorbed.

In treating poisoning by strychnia, a chemical antidote should be at once administered, such as tannic acid, or iodine, or one of its soluble salts. As, however, the compounds formed in the stomach by these substances are not permanent, a quick emetic must follow their administration. For the treatment of the symptoms, various substances have been recommended. It is evident that aconite, Calabar bean, tobacco, or their alkaloids, are indicated as more or less complete physiological antidotes.

The evidence brought forward by Dr. Haughton and others is not sufficient to establish clearly the especial value of tobacco; yet to me, from the quickness and completeness of its action, it seems the best of the substances above named.

It is obvious that the use of aconite, or of tobacco, in large doses, is accompanied with grave danger, on account of their influence upon the heart; and we have in bromide of potassium a substance devoid of any such objection, and apparently as complete a physiological antidote to strychnia as are any of the substances above named. The only question is as to whether the bromide has sufficient power and swiftness of action. In a single case,* recovery after the ingestion of three grains of the alkaloid, without vomiting, occurred, under the exhibition of a half-ounce dose of the potassium salt and "its continued use in smaller doses for an hour or so." The symptoms were as intense as consistent with life, but general relaxation was produced in thirty minutes after the ingestion of the counter-poison.

Chloral was stated by Liebreich, its therapeutic discoverer, to be antagonistic to strychnia, and it undoubtedly is so in a measure; but M. Orr (*Gazette Médicale*, July 6, 1872) stated to the French Academy that he had experimentally proven that the dose of chloral which Liebreich had relied on as being mortal to rabbits was very often not so; that the same was true of strychnia; and that consequently the investigation of Liebreich was not to be

* Dr. Cephas L. Bard (*Philadelphia Medical Times*, vol. i.). For cases in which the bromide was given along with other remedies successfully, see *New Remedies*, vol. ii. p. 255.

relied on as proving the respective antidotal powers of the drugs; and, further, that experiments had shown him that if a certainly-fatal dose of chloral were given to a rabbit, the hypodermic injection of strychnia did not affect the result, but that his own researches had not gone far enough to establish the exact relations of the drugs.

Prof. Bennett (*Edinburgh Medical Journal*, vol. xvi. pt. i., 1870, p. 262) has performed a number of experiments which have satisfied him as to the antidotal relations of chloral and strychnia. The only conclusion we can draw from the testimony, and from the known physiological effects of the two alkaloïds, is, I think, that they are in a greater or less degree opposed in their action; but to what extent, and in what ways, we are not yet in a position to state. It is also evident that chloral, like aconite, influences other portions of the body even more intensely than it does the cord; an objection certainly of more or less force to relying upon it in strychnia-poisoning.

On the whole, I should in a case of strychnia poisoning be disposed to rely upon the bromide, rather than any of the other supposed physiological antidotes, combining with it moderate doses of chloral, or the inhalation of ether, or even of chloroform, if necessary. In severe poisoning, half an ounce of the salt might be given at the first dose, and repeated in fifteen or twenty minutes; two-drachm doses being used afterwards if necessary. If nitrite of amyl were at hand, I should certainly try the *cautious* inhalation of it.

In some cases, artificial respiration may be of service. It is evident that the convulsed muscles will often resist such efforts as are usually made to force air into the lungs of man as successfully as they do the unassisted struggles of nature, and Harley states that he has found artificial respiration of no use whatever in animals. On the other hand, Leube (*Archives Anat. et Physiol.*, 1867), in an apparently very careful series of experiments, found that artificial respiration has great influence in saving or prolonging life, according to the amount of the poison ingested. In his experiments, the dose which ordinarily produced convulsions did not do this so long as artificial respiration was kept up; and the "lethal dose" did not kill if artificial respiration was maintained for four hours, although opisthotonos was induced in some cases. Rosenthal, according to Husemann, has obtained similar results, and M. Schiff (*Schmidt's Jahrbücher*, Bd. cxli. p. 43) has, in a series of experiments, corroborated in the main facts the results of the German investigators, although disagreeing with them in minor particulars. He found animals in which *forcible* artificial respiration was maintained survived doses much larger than those ordinarily fatal. The artificial respiration was performed by inserting a canula into the trachea, and filling the lungs by force.

ADMINISTRATION.—As a tonic, strychnia may be given in pill; but when it is desired to push it until its physiological effects are manifested, as in some cases of *paralysis*, it should be always administered as the sulphate in solution, because death has occurred from an irregularity in the solution of the pills in

the alimentary canal, and the consequent simultaneous letting loose of a large amount of the alkaloid. There is no proof of a cumulative action of this alkaloid when given as above directed. Dose, one-thirtieth of a grain, gradually increased *pro re nata*.

Strychniæ Sulphas—*Sulphate of Strychnia*—occurs in minute prismatic crystals. It is officinal on account of its solubility in water. The dose is one-twentieth of a grain.

IGNATIA AMARA.—The seeds of *Strychnos Ignatia*, a tree growing in the Philippine Islands. The *St. Ignatius Beans* are pale brown, about an inch in length, less in breadth, often angular, with three or four faces, covered with a very fine, scarcely visible down. They have been compared by some to an olive in size and appearance.

They contain largely of the igasurates of strychnia and of brucia, and are identical in their therapeutic value with *nux vomica*.

The only officinal preparation is the extract (*Extractum Ignatiæ*), which may be used in doses of one-fourth to one-half grain.

CLASS X.—DEPRESSO-MOTORS.

UNDER this heading are considered certain drugs which are used for the purpose of lessening the activity of the spinal cord. They have, except in this particular, but little in common in their action, and must be studied individually.

PHYSOSTIGMA—CALABAR BEAN.

An irregular, kidney-shaped bean, about an inch in length and three-fourths of an inch wide; the product of the *Physostigma venenosum*, a perennial woody creeper of Calabar, Africa, where the bean has been used by the natives as an ordeal test for criminals, witches, etc., since time immemorial. It contains an alkaloid known as *physostigmia*, or *eserina*. The most reliable tests for the alkaloid are as follows. A watery solution of it or of its salts containing potassa, soda, or lime, on exposure to the air, becomes red, and finally yellow, green, or blue: this is said to occur when only one one-hundred-thousandth part of the alkaloid is in the solution. Chloride of gold throws down from the solution a blue precipitate, out of which the gold is soon reduced. According to Dr. J. B. Edwards (*Medical Times and Gazette*, 1864), with strong sulphuric acid and bichromate of potash *physostigmia* yields a violet color, passing into red. This point needs investigating, in relation to the well-known strychnia test.

The physiological test consists in the placing of a drop of the suspected solution in the eye of a rabbit, when, if *physostigmia* be present, contraction of the pupil will be produced in from eight to twenty minutes.

PHYSIOLOGICAL ACTION.—When an animal receives a small fatal dose of Calabar bean, after a time muscular tremors appear, and almost immediately the animal falls to the ground, or lies down, in a state of perfect muscular flaccidity. The pupils generally contract, and the respirations become slow, irregular, and often stertorous. All reflex actions are almost at once diminished, and this diminution grows greater and greater, until it ends in their complete abolition. So long as the condition of the motor system allows of it, evidences of sensibility are manifested whenever the animal is in any way injured. According to Clementi Papi (*Schmidt's Jahrbücher*, Bd. cxlii. p. 287), the voice is completely lost. The muscular tremors persist during the whole period of paralysis, and, indeed, even after cessation of the respiration.

They vary greatly in intensity, and in some cases are so severe (Fraser) as to simulate general convulsions. As the minutes go by, the rhythm of the respirations becomes more and more affected, and at last death takes place quietly, consciousness being preserved until the last few gasping respirations close the scene. The pupils sometimes, but not always, dilate immediately after death.

After a small lethal dose of the poison, the fatal result is always due to failure of the respiration, and if the body be at once opened the heart is found still beating; indeed, it has been seen to continue to do so for one and a half hours after death (Fraser). If a very large amount of the drug be given, the animal falls almost at once, paralyzed, with only a few muscular twitchings. The pupils contract, and in a very short time the gasping respiration ceases. The heart is now found distended and passive, but often will contract under the stimulation of a galvanic current.

The symptoms induced by the drug in man are completely parallel with those that occur in the lower animals. They are giddiness, lessened heart-action, great muscular weakness, with, in most cases, contraction of the pupil, and sometimes with vomiting, and still more commonly with purging, which may be very free. A pupil of Gubler took 0.15 grain of the sulphate of eserina, and suffered, after a time, nausea, giddiness, and intense muscular weakness, so that he could not stand; three-quarters of an hour afterwards he vomited some of the solution mixed with bile, but his strength did not begin to return for two and a half hours.

The question here naturally arises. To what is the paralysis so prominent in poisoning by Calabar bean due? It is evident that the suspension of reflex action can have only three sources: paralysis of the spinal cord, of the nerve-trunks, or of the muscles. I shall examine the action of the drug upon these organs in inverse order.

The muscular twitchings which have already been spoken of have been mistaken for convulsions by M. Vintschgau (*Sitzungsberichte der Math.-Nat. Classe d. k. Akad. d. Wissenschaften*, Wien, 1867, Bd. lv., Abth. ii., p. 49), who, indeed, concluded that Calabar bean acts like strychnia, because violent convulsive tremors occurred, after injection of a dose of the poison under the skin, in all parts of a frog whose iliaes he had tied. Evidence to be brought forward hereafter, however, shows conclusively that there was some fallacy in this experiment. The fact that the muscular movements continue after death indicates that they are due to a direct action of the drug upon the muscles themselves. This conclusion is thoroughly established by the experiments of Laschkewich (*Virchow's Archiv*, 1866, Bd. xxxv. p. 294), of Fraser (*loc. cit.*), and of Leven and Laborde (*Schmidt's Jahrbücher*, Bd. cxlvi. p. 136). All of these investigators have noted that after death these contractions are increased by exposure to the air and by direct stimulation of the muscles; and Fraser has found that they occur in the frog during life after section of the supplying nerve, and also in a muscle actually cut out of the body. Laschkewich has confirmed the latter fact in the case of warm-blooded animals, and

Leven and Laborde have proven that previous destruction of the lower end of the spinal cord in a guinea-pig does not prevent the development of the muscular twitchings in the hind legs. Although Calabar bean does, therefore, have some direct influence upon the muscles, yet the paralysis produced by it is in no sense the result of this influence, which appears, indeed, to be of an exciting, rather than of a paralyzing, character, since at the time of death the contractility of the muscle is in no wise diminished: on the contrary, Fraser has noted that in poisoning by Calabar bean the supervention of rigor mortis and of loss of functional power in the muscle is very greatly delayed.

Neither is the paralysis caused by the physostigma due to an action on the nerve-trunks, since Dr. Laschkewich, Dr. Vintsehgau (*loc. cit.*, p. 161), and also Dr. Fraser, have found that when the galvanic current is applied to the crural nerve of either cold- or warm-blooded animals rapidly killed with Calabar bean, contractions are freely induced in the tributary muscles.

Indeed, the Scotch investigator, carrying his experiments still further, and using delicate instruments which it is not necessary here to describe, discovered that when the artery going to a hind leg was tied in a frog before the administration of the poison, after a quick death the rate of conduction of impulse was as rapid in the nerve to which the poison had had free access, as in its protected fellow.

Since the abolition of reflex activity has its origin neither in the muscular system nor in the nerve-trunks, it must be spinal. The truth of this conclusion, arrived at by exclusion, has been abundantly demonstrated by direct experiment. Thus, Fraser has found that if in the frog a peripheral nerve be protected by tying its artery and the batrachian be poisoned with Calabar bean, the paralysis in the protected limb occurs *pari passu* with that in the remainder of the body. Again, the same observer divided the spinal cord of a frog, and then cut or tied all the blood-vessels going to the posterior section of it. After this, the animal was poisoned with physostigma, and, whilst the usual symptoms developed themselves in the anterior portion of the body, reflex actions were unaffected in the posterior part. Further, Dr. Fraser has found that when the poison is applied directly to the cord, fibrillary contractions, due probably to a local irritant influence, are induced in the muscles supplied from below the point of application; but in a little while all movements cease, and even galvanization of the cord itself is unable to elicit response. It seems completely established by the evidence which has been brought forward that the most prominent effect of Calabar bean is a *depressant action upon the spinal centres*.

Recently, however, it has been asserted by Papi (*Schmidt's Jahrbücher*, Bd. cxlii. p. 287) that there is in frogs, preceding the stage of depression, one of exaltation of reflex action; and also, on the strength of some experiments upon animals with one-sided section of the optic thalamus, that the cause of loss of voluntary movement in Calabar bean poisoning is paralysis of

the conducting fibres passing from the upper brain to the spinal centres. I have not been able to procure the original paper of Dr. Papi (*Gazz. Lomb.*, 1868), but his conclusions seem to me highly improbable; and it is almost inconceivable that if a stage of reflex hyper-activity really existed it could have been overlooked in the careful experiments of Laschkewich and of Fraser.

Although, as has been demonstrated, the paralysis caused by physostigma is spinal, yet the drug is not without influence upon the nerves, since Dr. Fraser has found that when the blood-vessels of a frog's leg are tied, and the animal slowly poisoned by a small dose of the extract, whilst even many minutes after cessation of respiration both nerves seem equally intact, yet finally a time comes when the nerve of the poisoned leg refuses to react to the galvanic stimulus, although the functional power of the protected nerve, as well as of the muscles, is still perfect. This loss of functional power is probably rather in the termination of the nerve than in the trunk, for Dr. Fraser found that when all the blood-vessels supplying the gastrocnemius muscle were cut in a frog and the animal poisoned, at a certain time irritation of the crural nerve produced spasms of the gastrocnemius alone.

It is to be noted that this perturbation of the peripheral nerves has only been seen in the frog when slowly poisoned, in which case the heart continues to beat long after the cessation of respiration, so that the nerves are, as it were, macerated in a solution of the poison. In warm-blooded animals, nerve-paralysis evidently plays a very unimportant rôle in the production of the death, being, if it exist at all, of so slight intensity as to be imperceptible.

The afferent nerve-fibres probably preserve their function long after the motor fibres have been affected, as was seemingly proven in Dr. Fraser's experiments (*loc. cit.*, p. 19) by tying the vessels in the left leg of a frog which was afterwards poisoned with strychnia, when it was found that reflex movements were excited in the left leg by irritation of the right foot long after irritation of the left foot had ceased to cause movements in the right leg.

Dr. Fraser studied to some extent the effect of a strong solution of the poison when applied locally to a nerve, and found that the afferent fibres were affected before the efferent, and that finally the function of both of them was abolished.

Harley (*loc. cit.*, p. 151) and Papi (*loc. cit.*, p. 287) assert that Calabar bean has little or no influence upon the heart; but they are undoubtedly in error. In poisoning by small doses, the cardiac action of the drug is certainly very subordinate to that upon the nerve-centres; but, as has been shown by Dr. Fraser, when very large doses of the poison are administered, especially if they be injected into the jugular vein, death results from syncope or from consentaneous failure of the cardiac and the respiratory function, and the heart is found arrested in diastole, flaccid, but, according to Dr. Fraser and to Drs. C. Arnstein and P. Sustschinsky (*Untersuchungen aus dem Physiolog. Laboratorium in Würzburg, zweiter Theil*, p. 86), responding, although feebly and uncertainly, to direct stimulation.

When smaller doses of the poison are exhibited, there is slowing of the heart's action, as has been noted by Laschkewich (*loc. cit.*, p. 298), by Fraser (*loc. cit.*, p. 48), by Dr. J. Tachau (*Archiv der Heilkunde*, 1865, p. 70), and by other observers. Although, according to the experiments of Fraser, there is at first a slight fall of the blood-pressure, probably due, as he believes, to diminished pulse-frequency, yet, in spite of the latter, the arterial tension soon recovers itself and remains for a long time much above the normal point, while at the same time the individual cardiac beats are greatly increased in strength (Fraser, Bezold and Götz*).

Finally, the arterial pressure falls far below normal, and the power of the heart is gradually extinguished.

The question as to the exact method in which these changes are wrought is of very difficult answer. The long diastolic pauses and the slow strong beat of the heart suggest at once that a chief action of the drug is upon the inhibitory cardiac nervous system. Tachau, however (*loc. cit.*, p. 172), found that after section of the vagi the poison produces these phenomena in an even more intense degree than in the normal heart. As this has been confirmed by Laschkewich (*loc. cit.*) and by Fraser (*loc. cit.*, p. 49), it must be accepted as a proven fact, especially since Vintsehgau has found (*loc. cit.*, p. 71) that if in the frog the brain and the medulla be destroyed, physostigma still acts in its usual way on the heart. Tachau considers that this demonstrates that the cardiac phenomena of Calabar bean poisoning are not due to an action of the drug upon the inhibitory nerves. Arnstein and Sustschinsky, however, admitting the fact, deny that it warrants the conclusion. Their idea appears to be that it is conceivable that a substance shall so act upon the peripheral inhibitory nerve-endings as to cause them to influence the action of the heart without any other external impulse, and consequently when separated from the nerve-centres. This may be, but, with our present knowledge, seems somewhat doubtful.

In an elaborate series of experiments, Arnstein and Sustschinsky found that the excitability of the peripheral cardiac vagi is increased by Calabar bean. They first tested the effect of graduated galvanic currents applied to the divided vagi in the animal to be experimented with, until the exact strength of the weakest current capable of causing diastolic arrest was demonstrated, and then exhibited the drug and tested the nerves afterwards. They used in these experiments both rabbits and guinea-pigs, and found that, without a single exception, currents much weaker than those which previously were barely effective would, after the poisoning, stop the heart; also, the super-excitability of the peripheral inhibitory nerves was shown by the fact that under the influence of the drug the diastolic arrest continued much longer than normal after the withdrawal of the stimulus from the vagi.

* I have not seen the original paper of these authorities in the *Centralblatt f. Med. Wissenschaft*, 1867, but quote them from the paper of Arnstein and Sustschinsky.

These experiments certainly seem to prove that Calabar bean does cause excitation of the peripheral cardiac apparatus. Arnstein and Sustschinsky further confirm them by other experiments of exceeding interest. They injected into rabbits such amounts of atropia as completely to paralyze the peripheral cardiac vagi, so that the strongest currents when applied to the nerves failed to influence the heart's action, and then *restored functional power* to the pneumogastrics by injections of Calabar bean, so that currents of moderate intensity caused diastolic arrest. These experiments have not, that I am aware of, been repeated, but they have every appearance of being accurate: if they are so, they certainly prove what is claimed for them. In warm-blooded animals, at least, the vagi are not paralyzed during life, as is evinced by the experiments of Arnstein and Sustschinsky (*loc. cit.*, p. 101). Some of the results obtained by Fraser (*loc. cit.*, p. 36) apparently contradict, but in reality accord with them. In the frog there was, indeed, at last loss of functional power in the vagi, but not until very long after the cessation of respiration, after all the nerves of voluntary motion had lost their functional power,—*i.e.*, after death would have occurred in a mammal.

The fact that Calabar bean acts after section of the par vagum, as before stated, indicates that the drug has no influence upon the inhibitory centres,—a conclusion confirmed by Arnstein and Sustschinsky (*loc. cit.*, p. 102), who found that an injection of the drug into a carotid—*i.e.*, into the inhibitory centre—did not cause any marked immediate diminution in the number of the cardiac pulsations. Dr. Fraser's experiments upon the local cardiac application of the drug are seemingly confirmative of the experiments of Arnstein and Sustschinsky, for he found that when the poison was put directly on the heart, or into one of its chambers, it caused a prolonged diastolic pause, followed by contractions, interrupted by pauses, and finally resumption of regular contractions, or else by diastolic arrest, the heart still retaining its power of responding, in an embarrassed manner, to stimuli. After all, however, I cannot help suspecting that we do not yet understand fully the cardiac action of Calabar bean.

As already stated, a very prominent phenomenon in Calabar bean poisoning is rise of the arterial pressure. This, of course, may be of cardiac origin, or it may be due simply to a contraction of the arterioles; or it may arise from a combination of these causes. Since Bezold and Götz (quoted by Arnstein and Sustschinsky, p. 87) found that the arterial pressure still rose under the influence of the drug after section of the spinal cord high up,—*i.e.*, after general vaso-motor paralysis,—the increased force of the circulation must, at least in part, be due to a direct action of the drug on the heart. The fact that the rise is not so great in animals whose cords have been divided as in those uninjured certainly also indicates, though it does not prove, that the increased arterial tension is in part due to vaso-motor spasm.

Dr. Fraser believes that Calabar bean does produce this spasm; but his evidence is certainly insufficient to establish his conclusion. It consists simply

of some experiments upon frogs in which the spinal cord was divided, and, the animal being put on a "frog-plate," the arteries of the web were watched whilst Calabar bean was exhibited. Dr. Fraser believed that under these circumstances the arteries contracted considerably at first, and afterwards dilated. Dr. Harley (*London Practitioner*, 1869, vol. iii. p. 163) states as the result of his studies that Calabar bean can be seen, when applied locally, to cause contraction of the veins, the arteries remaining unaffected; whilst Fraser, in contradiction to this, believes that he has demonstrated that the local application of physostigma produces dilatation of the arteries. It is not necessary here to reiterate my objections to such evidence as this. If, under the circumstances of the first-mentioned experiments of Dr. Fraser, Calabar bean contracts the small vessels, it must be by a peripheric, not centric, action, since the vessels were separated by the division of the cord from the vaso-motor centres. It is plain that this is in direct contradiction to Dr. Fraser's experiments on the local application of the drug. Again, it is contradictory, rather than confirmatory, of that furnished by the study of the blood-pressure. The only logical conclusion seems to me to be that at present we have no proof that Calabar bean acts upon the vaso-motor nervous system. At the same time, we are not in a condition to deny that the drug has some such action.

The most important action of Calabar bean, not yet considered, is upon the pupil. But before discussing this it is necessary to say a few words in regard to the influence of the drug upon the temperature, upon the blood, and upon intestinal movements. The first of these is affected very slightly, but, according to Dr. Fraser's observation, is a little elevated.

I do not believe that physostigma has much action upon the blood. Certainly its influence upon the nervous system is a direct one, since Lewissou (*Reichert's Archiv*, 1870) has found that it acts upon the "salt frog" as upon the normal animal. Fraser, however, states that after death from this poison the blood coagulates slowly and loosely, and the red disks in dogs and in rabbits present various irregularities of outline, among which may be noted a well-marked stellar crenation, but that the respiratory function of the blood is not interfered with.

Intestinal peristalsis is very much increased by the action of Calabar bean (Westermann, *Schmidt's Jahrbücher*, Bd. cxxxviii. p. 290; Papi, *Ibid.*, Bd. cxlii. p. 287; Fraser, *loc. cit.*, p. 57). After poisonous doses, there is at first a stage of exceedingly active movements in the bowels; then spasmodic tetanic contraction of the intestines occurs, so that their calibre is very much diminished; and finally relaxation and dilatation take place. After death the vermicular movements are found very much lessened (Fraser), or altogether abolished (Tachau, *loc. cit.*, p. 73).

The action of Calabar bean upon the intestines appears to be peripheral, due to contact of the poison in the blood with the muscular fibres or the ganglionic nerve-cells in the walls of the bowels. For Westermann (*loc. cit.*, p. 291) found that extirpation of the cardiac ganglion had no effect upon the

action of the drug, but that tying of the mesenteric and of the coeliac arteries, before poisoning, prevented any increase in the peristalsis.

Calabar bean, as is well known, strongly contracts the pupil, both when applied to the eye and when exhibited internally. Evidently, as in the case of atropia, the pupillary action of Calabar bean should be studied from two points of view, the local and the constitutional.

The closeness of the analogy between the pupillary action of atropia and that of Calabar bean is seen in the fact that, like the former, the latter, as shown by the experiments of Vée and Leven (*Comptes-Rendus de la Société de Biologie*, 1865, p. 161) on chickens, does not affect the irides of birds. Thus, analogy would seem to prove that the influence of Calabar bean is directly upon the peripheral nerves of the iris.

Although, then, I am not able to cite any direct experimental proof, yet it seems to me scarcely doubtful that the contraction of the pupil produced by Calabar bean is always a local, peripheral influence, whether the drug be placed in the eye from the outside or be carried to the general circulation. It is evident that the myosis may be caused in one of three methods: by paralysis of the sympathetic fibres, by stimulation of the oculo-motor fibres, or by a conjoint action upon both sets of nerve-endings. In which of these ways the drug acts, we are not yet in a position positively to decide.

It has been held by various authorities that if galvanization of the sympathetic fibres in the neck fail to expand a contracted pupil, the myosis must be due to paralysis of the sympathetic. Evidently, however, this is claiming too much, for, as pointed out by Grünhagen (*Virchow's Archiv*, Bd. xxx. p. 521), it is conceivable that an oculo-motor spasm can exist of such intensity that the antagonistic nerve is unable to dilate the pupil. The question arises very pertinently at this point, What is the fact in regard to Calabar-myosis? Does or does not galvanic stimulation of the cervical sympathetic dilate the pupil? The testimony is somewhat conflicting. Dr. Grünhagen (*loc. cit.*, p. 521) says that dilatation always occurs, although to a slight extent (*in beschränkten Masse*); whilst, on the other hand, Dr. Gustav Engelhardt (*Untersuchungen aus dem Physiologischen Laboratorium in Würzburg*, zweiter Theil, p. 526) has found that galvanization of the cervical sympathetic has no effect upon the contracted pupil. The experiments of Fraser (*loc. cit.*, p. 60), of Bernstein and Dogiel, and of Rosenthal (*Reichert's Archiv*, 1863) reconcile these differences, and, by their accord, conclusively prove that under the maximum influence of Calabar bean the sympathetic is powerless, whilst when contraction is the result of a milder influence of the drug, stimulation of the cervical nerve will cause a certain amount of dilatation. Fraser, and also Engelhardt, have found that if the poles of a battery be applied directly to an iris even most profoundly contracted by physostigma, immediate dilatation occurs. These facts, for reasons stated above, do not absolutely prove, but they certainly render it highly probable, that Calabar bean *paralyzes the*

peripheral sympathetic nerve-fibres in the iris. It is, however, almost equally probable that there is a *consentaneous stimulation of the oculo-motor terminations*; for the myosis caused, like the mydriasis produced by atropia, is an active, not a passive, condition, and is not only much more forcible, but is also much more complete, than that which follows section of the cervical sympathetics.

In regard to the elimination of Calabar bean, I have met with no experiments.

THERAPEUTICS.—The physiological action of Calabar bean has suggested its use in spasmodic affections, in atony of the muscular coats of the bowel, and in various diseases of the eye.

The action of Calabar bean upon the spinal cord very early led to its use in spasmodic affections, and especially in *tetanus*, in which disease it has been more freely employed during the last few years than any other remedy except opium. In the paper of Dr. B. Roemer (*St. Louis Medical and Surgical Journal*, 1873, p. 367) are collected forty-seven cases, of which twenty proved fatal. To these I am able to add the ten whose references are given below,* making in all fifty-seven cases, with thirty-one recoveries and twenty-six deaths,—not a very flattering record.

It is, however, proper to state, as affecting the value of these statistics, that much of the Calabar bean extract which has been offered in the market is practically inert, and in all probability in some of these cases the drug did not have a fair trial; and that when especial care was taken by certain observers better results were achieved, although on so small a scale as to leave the issue in much doubt.†

In *trismus neonatorum*, Calabar bean has been employed with results certainly no more encouraging than those obtained in tetanus. In *chorea* it has also been used with very doubtful advantage.

The physiological action of physostigma upon the unstriated intestinal muscle-fibres has led to its employment in *atony* of the muscular coat of the bowels and other similar organs. Dr. V. Subbotin (*Archiv f. Klin. Med.*, Bd. vi. p. 285, 1869) has used the extract with the happiest results in a case of *chronic bronchial catarrh* with intense *dyspnoea*, believed to be due to weakness of the bronchial muscular fibres; and also in one of apparently “phantom tumor,” with *chronic intestinal dyspepsia* and *catarrh*. In *constipation* dependent upon relaxation, it is also said to be useful.

Calabar bean has also been employed in *strychnia-poisoning*, and a recovery

* **FATAL CASES.**—Fenwick, 1 (*Glasgow Medical Journal*, 1869, p. 300); Franzolin, 1 (*The Doctor*, Oct. 1, 1871); Laborde, 1 (*British Medical Journal*, June, 1872); Valdivieso, 1 (*Philadelphia Medical Times*, vol. i. p. 455); Tyson, 1 (*Ibid.*, p. 418); Johnson, 1 (*Ibid.*, p. 372). **RECOVERIES.**—Fenwick, 1 (*Glasgow Medical Journal*, 1869, p. 300); Newman, 1 (*Medical Examiner*, July, 1869); W. W. Keen, 1 (*Philadelphia Medical Times*, vol. i. p. 195); J. H. Packard, 1 (*Ibid.*, p. 138). All these cases were of the traumatic form of the disease.

† For a favorable record, see Watson, *Glasgow Medical Journal*, N. S., 1869, vol. i. p. 54; consult also *London Practitioner*, Sept. 1869.

obtained after the injection of three grains of the latter alkaloid is reported by Dr. J. W. Keyworth (*Glasgow Medical Journal*, N. S., 1869, i. 54).

In *epilepsy*, some trials have been made of the drug, but its value is very doubtful.

TOXICOLOGY.—So far as I know, Calabar bean has not been used, either in Europe or in this country, with criminal intent. Accidental poisoning has, however, been produced by it. In Liverpool seventy children were poisoned at one time by eating the beans, which had been thrown out with some rubbish (Dr. Cameron, *Medical Times and Gazette*, Oct. 1864, p. 406). Many of the victims vomited spontaneously, and thus relieved themselves. Those brought to the hospital were in a state of extreme prostration, lying perfectly relaxed. They appeared to suffer almost no pain, only some of them saying they had a little "belly-ache." Among some thirteen examined, only one had the pupils contracted. The only child who did not recover was excessively weak, and, crying out suddenly, was dead, as Dr. Cameron thinks, of sudden syncope. The heart was found relaxed and flabby, both sides equally full of blood.

In 1864 Dr. Kleinwächter treated a case of poisoning by an unknown quantity of atropia with Calabar bean, apparently with great benefit. Dr. Bourneville detailed in 1867 some experiments which seemed to show that there is a real antagonism between Calabar bean and the mydriatic, and in 1870 (*Revue Photographique des Hôpitaux*) published five experiments upon guinea-pigs, which were very decisive in that a proven fatal dose of physostigma was given in each case and recovery obtained by the use of non-lethal doses of atropia. In 1869 Prof. Roberts Bartholow, of Cincinnati, on the strength of a few really indecisive experiments, arrived at a conclusion opposite to that of Bourneville. Recently, Dr. Fraser, of Edinburgh, has investigated the subject in so thorough a manner that his essay may serve as a model for those who are desirous of studying questions of antagonisms between poisons. His experiments, three hundred and thirty-one in number, were made chiefly upon rabbits, a few having been upon dogs. He first investigated as to the minimum fatal dose, per pound of the animal, of the preparations used,—the extract of the bean, and the sulphate of eserina. It was ascertained that the minimum lethal dose for rabbits of the extract of physostigma which he employed was 0.4 grain per pound; of the sulphate of eserina, 0.04 grain per pound. Then in sixteen experiments in which recovery followed the administration of a dose of atropia given in combination with a dose of physostigma equal to or in excess of the minimum fatal dose, the animal used was killed long afterwards by a dose of the Calabar bean less than or equal to that from which recovery had occurred under the influence of atropia. In this way a *perfect* demonstration of the power of the counter-poison was obtained.

It was found that the counter-poison acted most efficiently when thrown directly into the veins. Thus, a rabbit weighing three pounds and two ounces

received 1.6 grains of the extract, and five minutes afterwards 0.02 grain of atropia, in a vein, and recovery took place; eight days after this, 1.3 grains of the extract killed the same rabbit in nineteen minutes: in another animal which nine days before had been saved from death after the exhibition of 2 grains of the extract by 0.5 grain of the sulphate of atropia, 1 grain of the extract proved fatal in thirteen minutes.

The next series of experiments was undertaken to ascertain the maximum dose of physostigma that can be successfully antagonized by atropia, and the dose in which the latter should be employed. In all cases the atropia was given five minutes before the Calabar bean. It was found that one-fiftieth of a grain of the mydriatic would successfully antagonize one and a half times, but not twice, the minimum fatal dose of the myotic.

One-fortieth of a grain of the atropia was successful against two to two and a half times the minimum lethal dose of physostigma; three-fiftieths was sufficient for three times the minimum fatal dose. The small size of the required doses of atropia is very noticeable, and at the present point in the investigation a very curious result was obtained. It was found that when three times the minimum fatal dose of the Calabar bean was exhibited, the successful dose of atropia ranged from three-fiftieths of a grain to one grain and a fifth. When three and a half times the fatal dose of the physostigma was exhibited, success was achieved only with doses of atropia of from one-tenth to one-fifth of a grain. Unfortunately, there are only seven experiments bearing upon this point: yet its general accuracy, I think, can scarcely be questioned. When a rabbit received four times the lethal dose of physostigma, the mydriatic was powerless.

No experiments were made to test the value of physostigma in atropia-poisoning. Dr. Fraser states, however, on what grounds I do not know, that the minimum fatal dose of atropia is twenty-one grains; and he found that when one-half the minimum lethal dose of physostigma is given, together with nine and four-fifths grains, or more, of atropia, death results.

In the final series of experiments, the atropia was administered five minutes after the physostigma, and it was found that the largest dose of the latter which could be combated successfully was three times the minimum fatal dose, and the range of the dose of atropia was much less than when it was given before the poisoning. Thus, with three times the minimum lethal dose of the Calabar bean, death occurred when three-twentieths or one-fifth of a grain of the antidote was given, but recovery followed the administration of four-twenty-fifths of a grain.

These researches ought certainly to be carried further, to discover, if possible, why it is, or rather how it is, that atropia acts as the antidote to Calabar bean. It will probably be found that the mydriatic saves life by its stimulant action upon the respiratory centres.

The question naturally arises, How far are the results of these experiments applicable to the treatment of Calabar bean poisoning in man? Without dis-

cussing this at length, I think the following deductions are obvious: first, that atropia ought by all means to be tried in poisoning by physostigma; second, that the doses of it should never be very large, not exceeding in all the tenth of a grain.

The use of atropia should, of course, not cause neglect of such measures of relief as evacuation of the stomach, the external application of dry heat, etc., usual in poisoning by sedative narcotics.

POTASSII BROMIDUM—BROMIDE OF POTASSIUM. (KBr.)

The bromide of potassium is prepared, according to the officinal method, by precipitating freshly-made solution of the bromide of iron by the pure carbonate of potash, filtering, and evaporating the resultant solution. It occurs in milk-white cubic or quadrangularly prismatic crystals of an acrid saline taste, freely soluble in water, and slightly so in alcohol. When it is mixed with starch, and chlorine is added, a yellow color is developed. A bluish tint betrays contamination with an iodide.

PHYSIOLOGICAL ACTION.—Bromide of potassium administered to frogs in minute doses produces as a first result a tetanoid condition, in which there may be very marked opisthotonos. After a short time this stage of muscular excitement gives way to one of great muscular relaxation, and total abolition of reflex actions. Voluntary movements, however, often occur during this period, and the frog which has been lying limp and apparently dead will startle the observer by a sudden vigorous leap. This fact has been so frequently witnessed that there can be no doubt of its truth. It is vouched for by the following observers: J. M. Purser (*Dublin Journal of Medical Sciences*, xlvii. 324, 1869); Lewisky aus Kazan (*Virchow's Archiv*, Bd. xlv. p. 191, 1869); J. V. Laborde (*Archives de Physiologie Normale et Pathologique*, t. i. 423, 1868, and *Comptes-Rendus*, t. lxx., 1867); MM. Damourette and Pelvette (*Bulletin Thérapeutique*, 1867, lxxiii. 249).

Very early in the paralytic stage the respiratory movements are affected, and they gradually grow less until their final arrest. When a very large dose of the bromide is given, death may be induced by paralysis of the heart (Albert Eulenberg and Paul Guttman, *Virchow's Archiv*, xli., 1867); but after a small toxic dose this viscus continues to beat long after the cessation of breathing. If the drug be given by an injection practiced in the vicinity of the heart, sudden cardiac arrest always occurs.

Upon mammals (*Bulletin Thérapeutique*, lxxiii. 256; *Virchow's Archiv*, xli. 97) the bromide acts very much as upon frogs, inducing progressive paralysis, depression of temperature, and death by asphyxia when given in small poisonous doses, and great disturbance of the circulation, with finally diastolic arrest of the heart, when very freely administered.

When a solution of the bromide is applied locally to the heart, it produces instantly marked lessening of its action, and, if in sufficient amount and concentration, even instantaneous diastolic arrest (*Virchow's Archiv*, xli. 101).

Upon the voluntary muscles it acts in a similar manner when similarly applied. If its solution be not too concentrated or abundant, however, the muscle of the frog is first thrown into a tetanic spasm (*Dublin Journal*, xlvii. 325); and Dr. Purser suggests that the tetanic symptoms seen in the frog poisoned by the bromide of potassium are due to this action on the muscles. On the nerve-trunks, and also on the nerve-centres, the bromide acts, when applied locally, as a paralyzing poison (*Bulletin Thérapeutique*, lxxiii. 253, 290; also Dr. Amory, *The Physiological and Therapeutical Action of the Bromide of Potassium*, Boston, 1872, Part II., p. 147).

In general poisoning by hypodermic injection of the bromide, this local action is often very manifest, and paralysis of the part into which the solution has been thrown follows very rapidly upon the injection.

It is, therefore, evident that the bromide of potassium in sufficient quantity is a deadly poison to all the higher animal tissues. To therapeutists, however, the interest centres about its action when exhibited in doses just sufficient to take life, or in still smaller quantities.

The persistence of voluntary movement in the frog after the abolition of reflex actions shows that the influence of the drug is not chiefly exerted upon the cerebral centres of motor impulse, nor upon those cells of the cord which originate movement, but upon either the afferent nerves or those portions of the cord which transmit the impulse from these nerves to the cells presiding immediately over motion. This is confirmed by some experiments of Lewisky, in which it was found that previous separation of the cord from the cerebrum had no influence upon the action of the bromide. Both he and Purser also found that death occurred from small doses before the motor nerve trunks and the muscles had lost their irritability (confirmed by Saisson, *Schmidt's Jahrbücher*, Bd. cxliii. p. 17). This being so, the question arises whether the paralysis be spinal or due to paralysis of the peripheral afferent nerves. There is an apparent conflict in the evidence upon this point. Eulenberg and Guttman (*loc. cit.*, p. 103) found that when access of the poison was prevented to one or more limbs by tying the arteries, reflex actions were abolished in these parts as rapidly as in others. Similar results have been obtained by Lewisky, by Roberts Bartholow (*Bromides: Their Physiological Effects*, Providence, 1871), by Purser* (*loc. cit.*, p. 326), and by Laborde (*loc. cit.*, p. 434). The latter observer has also found that electrical stimulation of a nerve high up will cause violent spasms in the muscles directly supplied by it, although it may be unable to excite the slightest reflex tremor. On the other hand, Damourette and Pelvette assert a contrary result. Unfortunately, they do not give the details of their experiments. They state, however (p. 247), that if the lumbar plexus of vessels be tied before the poisoning, the fore feet lose their reflex activity before the hinder. There are two possible

* From the wording of his memoir, however, it is doubtful whether Purser performed the experiment himself.

methods of reconciling their results with those of the other observers. In some way the operation may have interfered with the circulation in the lower part of the cord, and consequently the poison have reached more freely the upper part of it and acted first upon it. Again, if the injection was, as is very probable, thrown into the anterior portion of the body, the poison may have reached the anterior extremities in so concentrated a form as to have acted, as it were, locally upon their nerves and muscles. The same observers in another portion of their memoir show that the solutions of these salts travel by imbibition; and this and their local action seem to me to be the cause of the differences of experimental results. It seems well established that cutaneous anæsthesia in greater or less degree accompanies the loss of reflex activity; for, as Dr. Purser says, a poisoned animal quite able to jump submits to pinching, pricking, burning, etc., without moving. Eulenberg and Guttmann have noticed the same thing in some rabbits. Damourrette and Pelvette (*loc. cit.*, p. 247) have noticed a condition in which electrical stimulation of a nerve-trunk produced marked reflex action, although no excitement of the skin supplied by the afferent fibres of the nerve was capable of doing this, showing that the extremities of the sensitive nerves are affected before the trunks.

The evidence is, I think, sufficient to prove that bromide of potassium affects all parts of the nervous system of the lower animals, but that the cerebrum, the motor tract of the cord, and the efferent nerves are the last portions to be affected; that the most sensitive to its action is the receptive portion of the cord,—that which receives and transmits reflex impulses,—and next to this, and perhaps almost equally susceptible with it, are the peripheral ends of the afferent nerves.

Upon the cerebrum of the higher animals the bromides undoubtedly exert an influence; but I am not cognizant of any researches sufficiently elaborate to show its extent or nature.

It is well established that large toxic doses of the bromide exert a direct paralyzing action on the heart, lessening both the force and frequency of the beat, and finally causing diastolic arrest. Dr. J. G. Schouten (*Archiv der Heilk.*, xii. 2, p. 97, 1871; *Schmidt's Jahrbücher*, Bd. cliv. p. 11) found that during the slow injection of a two per cent. solution into the vena cava of a rabbit the cardiac systole grew slower, the diastolic pauses longer, and finally the heart stood still, exhibiting only fibrillary contractions of its walls. The same observer is, so far as I know, the only one who has made manometrical studies of the action of small doses of the drug. He found that such amounts of the bromide administered hypodermically or by the stomach always produced increased pulse-frequency with lessened arterial pressure. His experiments were, however, not carried far enough to demonstrate either how these two changes are brought about, or the relations of the drug to the vaso-motor nerves. Much has been predicated upon the theory which asserts that bromide of potassium causes vaso-motor spasm. No decisive proofs have,

however, yet been offered of the truth of this favorite dogma. The evidence so far brought forward is as follows: Lewisky found that if the toes of two frogs—one poisoned, the other not—were cut off, the unpoisoned frog bled much more freely than the other. This experiment has been confirmed by Dr. Amory: it, however, evidently does not prove the existence of vaso-motor spasm, but only that of a lessened activity of the circulation, which may be of cardiac origin.

According to Damourette and Pelvette (*loc. cit.*, p. 249), when the interdigital membrane of the frog is watched during poisoning, there is seen at first very often an exceedingly brief period of increased circulation, but in a very short time the latter becomes much slower. Dr. Meuriot (*L'Étude de la Belladone*, p. 49) asserts that by the aid of the microscope this slowing of the circulation can readily be seen to be due to a contraction of the capillaries, and especially of the small arteries, whose lumen may even be obliterated. Dr. Saisson (*Schmidt's Jahrbücher*, Bd. cxliii. p. 17) also asserts that he has witnessed a similar phenomenon in the tongue of the frog, and Dr. Hammond and Dr. Amory state that they have seen it in the brain of the dog. On the other hand, Dr. Purser (*loc. cit.*) and Dr. F. B. Nunneley (*London Practitioner*, vol. iii. p. 351) assert that the vessels in the frog's web are not affected by bromide of potassium given hypodermically.

My own studies of the action of various poisons upon the vessels of frog's web have yielded such varying and unsatisfactory results as to make me hesitate in accepting evidence of this nature unless otherwise corroborated. In the absence of manometrical studies, I think the most that can be fairly claimed is that our present knowledge renders it somewhat probable that the salt under consideration is capable of producing vaso-motor spasm. The further deduction that the nervous symptoms induced are secondary to and produced by this spasm is wholly gratuitous, unproven, and improbable. The action of the bromide when applied locally to the bared nerve demonstrates that it acts directly upon nerve-tissue. Further, the absolute anæmia of the bloodless "salt frog" produces no such nervous symptoms as does even a non-toxic dose of the bromide; and the direct experiments of Dr. A. Weil (*Reichert's Archiv für Anatomie*, 1871, p. 271) have shown that in the frog the complete abolition of circulation has no effect upon the spinal marrow or upon reflex actions during the first half-hour. The proof is very strong that the drug acts directly upon the nervous tissues.

In warm-blooded animals, toxic doses of the bromide of potassium lower very decidedly the temperature. Although the point cannot be considered at all settled, it is probable that this lowering of temperature is due to a direct checking of tissue-changes. Dr. J. H. Bill (*American Journal of the Medical Sciences*, July, 1868, p. 25) has found a constant decrease in the amount of carbonic acid eliminated after the ingestion of the salt. In his experiments, the daily excretion of urea was not perceptibly affected; but Dr. Rabuteau found it slightly lessened (*Gazette Hebdomadaire*, 1869), as did also Dr. Bartholow (*loc. cit.*, p. 11).

When the blood is charged with the bromide, the salt probably escapes with all the secretions. It has been found by Voisin, Amory, Namias, Bill, etc., in the saliva and in the urine. Amory (Thesis of Dr. H. P. Bowditch, *Boston Medical and Surgical Journal*, Oct. 1868) has also demonstrated its presence in the perspiration. In the body of a man who died whilst taking it, M. Namias (*Comptes-Rendus*, tome lxx. p. 882) found it in all the liquids, as well as in the brain, liver, spinal cord, lungs, etc.

Elimination takes place to a certain extent through the skin, and to some extent through the intestinal mucous membrane also. Dr. Bill (*loc. cit.*, p. 25) always detected it in marked quantities in the fæces of men taking it; and H. Quinke (*Reichert's Archiv für Anatomie*, 1868, xxxv. 158) found that when forty grains of the bromide of sodium were given to dogs with intestinal fistula, two and a half hours afterwards the intestinal juices were free from the bromides, which reappeared in them after from three to six hours. The salt escapes also through the kidneys. The rapidity of elimination seems to vary: thus, Amory recovered one-half of the amount ingested during the first and one-third during the second twenty-four hours, and Mr. Ware (Thesis of Dr. H. P. Bowditch, *loc. cit.*) obtained a little more than half of the amount ingested in the urine of the succeeding thirty-two hours, whilst Bill was not able to get more than one-eighteenth of it during the first day. Dr. Bill has frequently found the bromides in the urine two weeks after the last dose has been exhibited; and Dr. Rabuteau has seen its presence persist under similar circumstances for a month.

No case of acute poisoning with the bromide of potassium has been reported, and I have never seen a single dose of it produce any obvious effect: the results of the continuous employment of large doses of it, however, demonstrate that it acts upon man as upon the lower animals. When it is taken with sufficient freedom to accumulate in the system, a conjunction of phenomena known as *bromism* arises. The cerebral symptoms are a sense of mental weakness, heaviness of intellect, failure of memory, partial aphasia, great somnolence, and depression of spirits. With these there may be decided impairment of the sensibility of the mucous membranes and of the skin, so that titillation of the fauces may be without effect, and, according to Puche, even heat applied to the skin calls forth no complaint, and Huette (*Mémoires de la Soc. Biolog.*, 1850) has seen in some cases absolute anæsthesia of the sclerotic conjunctiva. The sexual function is abolished. There is also very generally fetid breath, and an eruption of acne, which may indeed be very severe. Of course, in any individual case of bromism many of these symptoms may be wanting; but when the use of the remedy is persisted in, they all at last become developed in an intense degree. Professor Edward H. Clarke thus speaks of a case which came under his notice: "The fetid breath becomes nauseous; œdema supervenes on congestion of the uvula and fauces; the whispering voice sinks into aphonia; sexual weakness degenerates into impotence; muscular weakness becomes complete paralysis; reflex, general, and special sensations

disappear; the ears do not hear, nor the eyes see, nor the tongue taste; the expression of hebetude becomes first that of imbecility, then that of idiocy; hallucinations of sight and sound, with or without mania, precede general cerebral indifference, apathy, and paralysis; the respiration, without the ster-tor of opium or alcohol, is easy and slow; the temperature of the body is lowered; as the bromism becomes more profound, the patient lies quiet in bed, unable to move or feel or swallow or speak, with dilated and uncontractile pupils, and scarcely any change of the color of his skin or face."

It is evident that upon man the bromide in large doses acts as a depressant, which, although affecting decidedly the circulation, acts especially upon the nervous system, depressing the cerebral functions, lowering the reflex excitability of the cord, and paralyzing the peripheral ends of the afferent nerves in the same order and degree as it does in the lower animals.

THERAPEUTICS.—The bromide of potassium is employed by the therapist to quiet *cerebral excitement* when not inflammatory in its nature; to lessen over-susceptibility of the spinal centres of reflex action, or of the peripheral afferent nerves which lead to these centres; and to subdue nervous excitement of the genital system. The contra-indication to its use is the existence of a high degree of debility or exhaustion.

There are various forms of *nervous excitement* or unrest, such as sometimes follow excessive intellectual toil, anxiety, and other nervous strain, or occur during convalescence from acute disorder, in which the salt now under consideration is very valuable. The same may be said of some forms of *hysteria*. In some cases of *neuralgia* it undoubtedly affords great relief, but in the majority of cases it fails. It has seemed to me useless in neuralgia dependent upon anæmia or want of power; and my experience agrees with that of Dr. Anstie, that it is especially useful in persons of good nervous power, muscular force, and activity of circulation. As a hypnotic, it is employed in wakefulness from nervous excitement and in *delirium tremens*.

The chief use of the bromides is, however, to lessen reflex activity. It is especially in *epilepsy* that it has attained a most deserved reputation, doing far more good than all other remedies combined, sometimes effecting cures, more commonly ameliorating the symptoms, but occasionally failing entirely. There is no known method of distinguishing before trial with any certainty in what cases it will do good. Trousseau (*Clinical Medicine*, Sydenham Society, London) and Bartholow (*loc. cit.*) both assert, however, that it is least efficient in the mild form of the disorder known as the *petit mal*. The most brilliant results have, as a rule, been obtained in cases of not too long duration in which the fits are frequent and severe. The governing principle in its use is to try it in every case, increasing the dose until a mild degree of bromism is induced, and being guided by the results.

This salt is also often efficacious in other reflex spasmodic neuroses; in the *vomiting of pregnancy* or of uterine disease; in the *convulsions* of children; and, according to Dr. J. T. Rothrock, in preventing the so-called *urethral*

fever induced in very susceptible males by the introduction of the catheter or bougie. The physiological action of the salt seemingly indicates that it is of all known remedies the one best suited for the treatment of *tetanus*. Clinical experience is not as yet sufficient to enable us to come to a definite conclusion, but I have been unable to find a recorded death from the disease after the free exhibition of the bromide, although, according to Dr. Roemer, one or two have occurred.

In any case not less than a half-ounce of the salt should be exhibited in the day, and at night chloral should be used as a hypnotic (see article on Chloral).

The following table contains all the statistical information I have on the subject. Cases not otherwise marked recovered :

REPORTERS.	No.	KIND.	PLACE REPORTED.	RESULTS.
Dr. Bachenal.	1	Traumatic.	London Lancet, Feb. 1869.	
Dr. Derby.	1	"	Boston Med. and Surgical Journal.	Morphia also used.
A. Fergusson.	1	"	Edinburgh Med. Journal, July, 1872.	Chloral also used.
	1	"	U. S. Army Med. Department Circular No. 3, Aug. 1871.	
Dr. Panas.	1	"	Gazette Hebdomadaire, No. 26, 1872.	Opium also used.
Dr. C. L. Bard.	1	"	New Remedies, Jan. 1873.	
Dr. Bruchon.	1	"	Bulletin Thérapeutique, vol. lxxvii. p. 8.	Small bleeding and etherization also used.
Robert Brown.	1	"	Edinburgh Med. and Surg. Journal, vol. xiv., 1869, p. 992.	
Prof. May Figueira.	1	"	Bulletin Thérapeutique, vol. lxxvii. p. 428.	
Ibid.	1	Idiopathic.	Ibid.	
Allen Coultts.	1	"	London Practitioner, April, 1871.	Calabar bean also used.
H. F. Andrews.	1	?	*	?
H. K. Steel.	1	?	*	?
G. Derby.	1	?	*	?
Dr. Bakewell.	1	?	*	?
Mr. Hancock.	1	?	*	?
				Died.† 30 grains of the bromide every four hours, with 20 minims of the tincture of belladonna.

In *strychnia-poisoning*, Dr. Saisson has demonstrated its value by experiments on animals, and Dr. Chas. B. Gillespie (*American Journal of the Medical Sciences*, Oct. 1870) and Dr. C. L. Bard (*Philadelphia Medical Times*, June, 1871) have each reported recovery under its use, without vomiting, after the ingestion of three grains of the alkaloid.

In nervous excitement connected with the *genital function*, the bromides are often of great value. When there is actual inflammatory disease, as in *gonorrhœa*, the drug frequently fails to effect the desired end. When, however, there is no organic lesion of the organs or of their nerve-centres, the continued dose almost always succeeds to a greater or less extent. I have found the remedy effective in cases of semi-impotence from over-irrita-

* All the cases marked with an asterisk are taken from Dr. Roemer's paper (*St. Louis Medical and Surgical Journal*, 1873). I believe they were all traumatic.

† Since the belladonna probably did as much harm as the bromide did good, this case ought to be excluded.

bility of the organs causing emission too soon during attempted sexual congress. There is abundant evidence as to the value of the remedy in *nymphomania*. As an adjuvant to other physical and moral measures of relief, the salt may be used with satisfaction in men suffering from masturbation. In nervous symptoms occurring at the time of the menopause or complicating uterine disease, and in the peculiar train of morbid phenomena arising from the forced suppression of the sexual function in vigorous individuals of either sex to whom circumstances have denied marriage, the bromides have almost "a unique power."

ADMINISTRATION.—I have known half an ounce of the bromide to be taken at once without inducing any serious symptoms; and in severe acute cases, as in tetanus and strychnia-poisoning, it is perfectly safe to administer two-drachm doses at short intervals, as the case may require. Almost all the indications for the use of the bromide are best met by the so-called continuous dose,—i.e., by the administration of so much in the twenty-four hours until an effect is induced. Thus, in epilepsy, half a drachm may be given four times a day, to be increased to one drachm (half an ounce a day) if necessary; although as little of the remedy as will suffice to prevent the recurrence of the fit must be used: yet any amount necessary to do this should be given, unless bromism be produced before the paroxysms are arrested.

The remedy must be given, in a solution freely diluted, after meals. In some cases it causes diarrhœa, which may generally be checked with small doses of opium.

Several of the bromides have been used in medicine as a substitute for the salt of potassium, but none of them have been made officinal. Chief among these substitutes is the *bromide of sodium*, a drug which closely resembles in appearance the bromide of potassium, and has been supposed by Voisin to have closely similar physiological and therapeutic properties. On the other hand, M. J. V. Laborde states (*Robin's Journal de l'Anatomie et de Physiologie*, 1868, p. 560) that in double the toxic dose of the bromide of potassium he has found that it does not produce any characteristic symptoms in the frog, the guinea-pig, or the dog, and leaves the animal perfectly healthy.

By clinicians the drug has been used to a considerable extent. Dr. Meredith Clymer (*New York Medical World*, October, 1871) claims that it will arrest epilepsy without producing the unpleasant cerebral symptoms of bromism. He gives twenty grains three times a day.

Prof. Hammond (*New York Medical Journal*, Dec. 1871) asserts that in epilepsy it is in no wise superior to the potassium salt, but claims that its hypnotic power is much greater.

M. E. Decaisne, as the result of the trial of the drug in twenty-seven cases (*epilepsy, chorea, hysteria*), asserts that its action is the same as that of the potassium salt, except that instead of causing diarrhœa it constipates (*Comptes-Rendus*, No. 17, 1870).

The *bromide of lithium* was, I believe, first employed in medicine by Dr. Gibb (*Report of the British Association for the Advancement of Science*, 1864), who recommended it as gently tonic and sometimes diuretic. He used it in very small doses. Attention was first called to its employment in nervous affections by Dr. S. Weir Mitchell, who stated (*American Journal of the Medical Sciences*, Oct. 1870), as the result of his experience, that when administered to the amount of half a drachm to one drachm daily, it acts in some cases of epilepsy after the bromide of potassium has failed, and is generally efficient in about one-half the dose of that salt; also that its hypnotic action is much more decided. Prof. Clark (*loc. cit.*, p. 111) confirms these observations.

AMMONII BROMIDUM—BROMIDE OF AMMONIUM.

According to the U. S. Pharmacopœia, this salt should be prepared by the precipitation of the freshly-made solution of the bromide of iron by water of ammonia, the desired salt remaining in solution. It may be obtained in colorless crystals, but generally occurs in a granular powder, which becomes yellowish on exposure. It has a saline, pungent taste, is readily soluble in water, sparingly so in alcohol. When mixed with mucilage of starch, if chlorine-water be added it becomes yellowish brown; a blue tint would indicate the presence of iodine.

PHYSIOLOGICAL ACTION.—The physiological action of the bromide of ammonium has not as yet been fully investigated; but our present knowledge indicates that whilst in many points it resembles that of the corresponding salt of potassium, in others it differs essentially from the latter. According to Dr. N. Bistrotff (*Reichert's Archiv für Anatomie*, 1868, p. 723), when two decigrammes are administered to a frog, a period of quietude and lessened irritability is induced, which after fifteen or twenty minutes gives place to violent tetanic convulsion. Later, all excitability is lost, so that even burning calls forth no recognition; the frog lies in whatever position it is placed, the spasms become more violent, and death ensues. Similar phenomena have been witnessed by both Bistrotff and Amory (*The Physiological Action of the Bromide of Potassium and of the Bromide of Ammonium*, Boston, 1872) in the rabbit and guinea-pig, although in one of Dr. Amory's experiments the guinea-pig died without convulsions having been noted. The curious abolition of reflex action and of sensibility consentaneously with the occurrence of violent convulsions was noted frequently, and death seems always to have resulted from asphyxia. In the experience of Dr. Bistrotff, moderate non-fatal doses produced only weakness and uncertain movements in the rabbit.

The bromide of ammonium appears to exert very little influence upon the peripheral motor apparatus. Amory has seen the nerves retain their power of conduction after having been placed in a "strong solution;" and, according to Dr. Bistrotff, muscles retain their irritability after five minutes' soaking in a ten per cent. solution. According to the latter observer, the heart always continues beating after death from the drug, and the heart removed from the

batrachian and laid in a ten per cent. solution did not in any degree lose its normal activity. Even a twenty per cent. solution dropped upon the bared heart produced only a momentary arrest of the ventricular systole.

On the other hand, it is proper to state that Dr. Purser asserts that the heart is soon brought still in diastole in poisoning by this salt, and that the nerves and muscles also lose their irritability sooner than after poisoning by bromide of potassium.

The subject certainly needs further investigation; but it seems to me most probable that the bromide of ammonium exerts less influence upon the muscles than the bromide of potassium, but that in other respects their actions are very similar. The experiments of Amory indicate that the ammonium salt affects temperature and acts on the capillaries in the same way as that of potassium, and that it is also eliminated in a similar manner.

The experiments of Dr. Bistrotff show that in the cat, at least, the bromide of ammonium has no especial influence, as has been claimed, upon the superior laryngeal nerves.

THERAPEUTICS.—The bromide of ammonium has been employed by practitioners for exactly the same purposes as the corresponding potassium salt, but certainly has not made its way into general use. Dr. Brown-Séquard commends highly the combination of it and its sister-salt in *epilepsy* as being much superior to either of them alone. He gives sixty grains of the potassium with thirty of the ammonium salt, and claims especially immunity from the disagreeable symptoms of bromism. On the other hand, Echeverria (*On Epilepsy*, p. 316) asserts that the combination is in no wise superior to the bromide of potassium. Prof. Clark, however (*loc. cit.*, p. 106), from an experience in other than epileptic neuroses, is inclined to coincide with Dr. Brown-Séquard.

The disease which the bromide of ammonium has been most used in is *pertussis*. Dr. J. M. Da Costa commends it highly in *acute rheumatism* (*Pennsylvania Hospital Reports*, vol. ii.): a drachm to a drachm and a half of it was given, well diluted, in the twenty-four hours.

CHLORAL HYDRATE—HYDRATE OF CHLORAL.

Chloral, which is itself not used in medicine, is an oily liquid which at the ordinary temperatures gives off pungent fumes, and which is manufactured by the action of chloral on alcohol. United with water, this oily liquid is converted into a hydrate.

*Chloral Hydrate** is a volatile, crystalline solid, of a hot, burning taste, insoluble in cold chloroform, but very soluble in water, ether, and alcohol. It occurs generally as transparent, colorless tablets, but sometimes in acicular or even in rhomboid crystals. The compound of chloral and alcohol,—*Chloral*

* There is a chloral hydrate containing a very small proportion of water, which is insoluble in the latter menstruum. (*Gmelin's Handbook*.)

Alcoholate.—which resembles closely the hydrate, can be distinguished at once by its insolubility in water and its solubility in cold chloroform.

If an alkali be added to a solution of chloral hydrate, it breaks up into formic acid and chloroform, which, when water has been the solvent, at once separates in the form of oily drops.

PHYSIOLOGICAL ACTION.—When chloral is given to man or other mammals in moderate doses, the most prominent result in the great majority of instances is a quiet sleep, as closely allied as possible to natural sleep. The subject can readily be aroused from the lighter degrees of this, waking to full consciousness, but soon dropping off again when left quiet. The pulse is in this degree of action not affected, or is rendered a little slower; the pupil is contracted, but becomes normal so soon as the subject is awakened; the respiration is deep, full, and regular. When larger amounts are given, the sleep is much deeper, and may pass into profound coma; the respirations fall in number; the pulse is weakened and rendered slower, but may become rapid and irregular if the dose has been toxic; the temperature is reduced; the muscular system is relaxed, and both sensibility and reflex action are diminished. If a fatal dose has been given, all these symptoms are intensified: with coma, intense muscular relaxation, weak, thready pulse, and a pupil contracted at first, but afterwards dilated, the animal gradually sinks into death, paralyzed and anaesthetic.

The immediate cause of death is generally a paralytic arrest of respiration; but in many cases there appears to be a simultaneous arrest of the cardiac action, and it is very possible that fatal syncope may at times occur. At post-mortem examination, congestion of the meninges and substance of the brain and cord, and of the lungs, is commonly found. The blood is thought by Richardson (*Medical Times and Gazette*, Sept. 4, 1870) to coagulate less firmly than when normal.

The most constant and prominent of all the symptoms produced by moderate doses of chloral is sleep: this is without doubt due to a direct action of the drug upon the cerebrum. In most cases, as already stated, it is quiet, but sometimes is restless, and in man has even occasionally been wildly delirious; although it is somewhat uncertain whether the latter condition may not have been due to impurities in the drug. It seems to be well established that in the milder degrees of this sleep there is no anaesthesia. Demarquay (*Bulletin Thérapeutique*, lxxvii. p. 307) claims that hyperaesthesia very commonly follows the exhibition of small doses. Bouchut (*New York Medical Gazette*, Dec. 1870) found that it does occur, although somewhat rarely. Dieulafoy and Krishaber (*American Journal of the Medical Sciences*, Jan. 1870), Giovanni and Ranzoli (*Schmidt's Jahrbücher*, Bd. cli.), confirm this, whilst Liebreich and Labbé* deny it, and Hammarsten, who has noticed such hyperaesthesia, is inclined to think it apparent rather than real. It appears to me most probable

* He has noticed it, however, in a single case, confined to the ears. (*Archives Générales*, 1870, tome xvi. p. 338.)

that its existence must be acknowledged either as a constant or an occasional phenomenon, for Rajewsky (*Schmidt's Jahrbücher*, Bd. cli.), as the result of apparently very elaborate experimentation, recognizes it, and also states that there is in frogs a corresponding period of over-excitability of the reflex centres. The last observer mentions that in rabbits he has noticed a glowing heat borne without much complaint, when pinching would produce violent outeries. There can be no doubt, however, that in *very large* doses chloral produces anaesthesia; but, unless the amount employed be so great as to be almost toxic, this anaesthesia is in most cases very trifling.

The paralysis and loss of reflex excitability induced by chloral are not muscular in their origin, for Labbé has found that after death the muscles respond perfectly to galvanism. Both Labbé (*loc. cit.*) and Rajewsky (*loc. cit.*) have found that the motor nerves are in no wise affected by large or even fatal doses of chloral, which must therefore act upon the spinal cord to produce the paralytic phenomena. The experiments of Rajewsky have afforded positive confirmation of the conclusion arrived at by this process of exclusion; for he found that in the latter stages of chloral-poisoning direct irritation of the spinal cord gave rise to much less severe spasms than in the unpoisoned animal. Before this paralytic stage is reached, as already stated, Rajewsky affirms that in the frog there is a period of increased reflex activity, and that at this time stimulation of the spinal ganglia shows that they are more susceptible than normal. According to the observer last named, these phenomena occur just as freely after destruction of Setschenow's centre in the frog as before, and are therefore independent of it.

According to Demarquay, when chloral has been administered to animals there is evident enlargement and engorgement of all their blood-vessels; and Rajewsky* states that he has found sinking of the blood-pressure in rabbits from small as well as large doses of the drug. On the other side, Labbé (*loc. cit.*, p. 341) asserts that the rabbit's ear grows pale after the injection of a very feeble dose. In man, Bouchut has obtained sphygmographic traces, which he thinks indicate a primary increased arterial tension. Nancias, of Venice, has found the tension normal, but Anstie and Andrews (*American Journal of Insanity*, July, 1871) confirm the results of Bouchut when small doses are employed. After very large doses, according to both Andrews and Da Costa (*Amer. Jour. Med. Sci.*, April, 1870), the tracings indicate very much lessened arterial pressure. Whatever may be the effect of small doses (and the point needs further investigation), it seems well made out that, both in man and animals, by large doses the blood-pressure is very much lessened, probably in part owing to vaso-motor paralysis, but in largest measure to an action of the drug on the heart. Various observers state that there is in man and the lower

* The work of Rajewsky was done under Professor Rosenthal, and the results, without the experiments, were published as an inaugural thesis. Of course, half of its value is lost for want of the experimental records, and the subject needs re-investigation. I am acquainted with the pamphlet only through *Schmidt's Jahrbücher*.

animals at this time slowing of the pulse, which, according to Rajewsky, is altogether independent of the inhibitory nerves in the frog and rabbit, occurring equally after as before their section. When toxic doses have been employed, the heart, after numerous pauses, is finally arrested in diastole. This arrest appears not to be muscular in its origin, for both Rajewsky and Labbée state that galvanic stimulation will produce a single full beat; and the latter author affirms that when the cord has been previously cut in the frog, the heart will continue to beat for hours. Further, Labbée states that when powdered chloral is placed upon the frog's heart freely, there results a marked slowing, but no arrest of its action. It would seem, therefore, most probable that the chloral influences the heart through the centres at the base of the brain.

In poisoning in man, the pulse has towards the last been very feeble, generally rapid and irregular, and, even in some cases in which recovery has occurred, it has been altogether absent for a time.

In full doses, chloral lessens the number of respirations per minute, causing them to become slow and full; and when toxic doses are taken their action becomes more and more marked, until finally the rhythm is very much affected, and the respiration grows very irregular, and sometimes very rapid and shallow, until finally it ceases. As these phenomena occur equally after section of the vagi (Rajewsky), the influence of chloral must be exerted upon the respiratory centre at the base of the brain.

A most remarkable action of chloral is upon the temperature; and in this point all observers are in accord with Dr. Richardson, of London, who has seen it fall 6° F. in a rabbit which recovered. Bouchut has noticed a fall of 2° (C.?) in an infant, and Da Costa and other observers have noticed lesser reductions of temperature in man. Hammarsten has found that the fall of temperature is very rapid, 6° C. in an hour, and that it is dependent upon diminished production of heat, since it occurs equally in animals well wrapped up and laid in a warm place.

Locally applied, chloral is an irritant; and probably in virtue of this it sometimes, when given by the mouth, causes vomiting, or even purging.

The physiological action of chloral may be summed up as follows. Upon the cerebrum it acts as a most powerful and certain hypnotic; in full doses it acts as an intense depressant upon the centres at the base of the brain, and upon the spinal cord, causing slowing and weakness of the heart's action, probably vaso-motor paralysis, slowing of the respiration, and muscular weakness, with a certain amount of anaesthesia; in fatal doses it causes death generally by arresting, through paralysis of the nerve-centres, first respiration, and finally the heart in diastole. Its action in very small doses is uncertain, but there is considerable evidence to indicate that it irritates or stimulates the spinal and the cardiac, and even the vaso-motor, centres. On the vagi and on the motor nerve trunks it has no marked influence.

The conversion of chloral by alkalis in solution into chloroform and formic acid first suggested its use in medicine to Liebreich (*Wiener Medicinische*

Wochenschrift, August, 1869); and the theory that its action is really due to chloroform generated by the alkalinity of the blood has been received with favor by Personne and other writers. The evidence by which this theory is to be disproven or established is twofold in its nature,—i.e., chemical and physiological.

Personne (*Journal de Pharmacie et de Chimie*, 1870), by distilling the blood of animals poisoned by chloral at 40° C., a temperature decidedly above that of the body, obtained chloroform; and his results have been confirmed by Pellogio (*Schmidt's Jahrbücher*, Bd. cli. p. 89) and other chemists, so that I think their correctness is not to be questioned. They are, however, not decisive, for it is very possible that the chloroform may be formed during the distillation, owing to the comparatively high heat employed. This evidence is, then, to be thrown out, especially as Hammarsten (*Ibid.*, Bd. cli.) has found that if chloral be mixed with fresh blood and streams of carbonic acid be forced through it, no chloroform can be detected, but if the mixture be heated the latter can be obtained in abundance. Further, in dogs deeply poisoned with chloral the same observer examined the expired air and a portion of the blood without finding chloroform, although when a clyster of the latter substance was administered it could be detected in a few minutes, in both the blood and the breath. In non-chloralized animals to which similar enemata were given, the chloroform could be detected in the breath before anæsthesia was induced. Rajewsky (*Ibid.*, Bd. cli.) has confirmed these results, as has also Amory (*New York Medical Journal*, 1870): so that it may be considered settled that chloral mixed with blood at ordinary temperatures remains unaltered, and that in the most severe poisoning no chloroform can be detected either in the blood or the expired air.

Hammarsten examined the blood and expired air unsuccessfully, for chloral, in dogs poisoned with the drug; but Amory (*loc. cit.*, p. 616) has obtained from the blood acicular crystals, evidently of chloral. It is still uncertain what becomes of the substance, whether it is finally destroyed in the organism or eliminated unchanged. All attempts to find it in the breath or the secretions have so far yielded negative results only; but this may be because there is no known delicate test.

To my mind, in this case the physiological evidence is in strict accord with the chemical facts, for the symptoms induced by chloral are only analogous to those caused by chloroform, *not identical*. As was pointed out by Dr. S. Weir Mitchell, and as any one can readily verify, if proportionate doses of chloral and chloroform be given to animals the sleep induced by the former is not only much more prolonged, but also more intense; further, the anæsthetic effect of chloroform is much more intense than that of chloral. Moreover, Djurberg (*Schmidt's Jahrbücher*, Bd. cli. p. 84) has shown that, whilst after chloroform-poisoning biliary coloring-matters appear in the urine, after chloral-poisoning none can be detected, and that when chloral is added to blood outside of the body no destruction of the red disks occurs.

As was first shown by Cohnheim, if the abdominal vein of a frog be opened and salt water be injected until almost all the blood is washed out and the circulatory system is filled with the foreign fluid, the batrachian will live from one to three days in apparently perfect health. Both Rajewsky (*loc. cit.*, p. 91) and Lewisson (*Reichert's Archiv für Anatomie und Physiologie*, 1870, p. 346) have found that upon these "salt frogs," with the circulating fluid completely neutral, the chloral acts precisely as in the normal frog.

I think the evidence which has been adduced completely disproves the chloroform-theory, and forces assent to the proposition that chloral acts directly upon the organism.

THERAPEUTICS.—The results of the clinical use of chloral are in strict accord with its known physiological action. The indication which it most usefully meets is *to induce sleep*. The more purely nervous the wakefulness is, the more successful is this remedy. When from functional over-excitement of the brain due to excessive mental strain, or from anxiety or other kindred cause, the patient cannot sleep, chloral is by far our most valuable hypnotic. On the other hand, when severe pain causes wakefulness, chloral is of very little value,—at least in doses which I think safe. Sometimes even in these cases sleep will come, but it will be a restless, troubled sleep, with moaning or other indications of suffering; and it may be that the patient on awaking will complain that he has suffered more whilst sleeping than when awake.

In the *sleeplessness* occurring at times during *convalescence* from acute disease, chloral is very efficacious. In the early stages of *fevers* it is sometimes of advantage; Dr. Russell (*Glasgow Medical Journal*, Feb. 1860) recommends it especially in the *wild delirium of typhus* in its earlier stages. In advanced fever-cases, when the symptoms are gravely adynamic, I conceive that the use of chloral would be very perilous. In *delirium tremens* it often induces sleep readily, but not rarely it fails, even in large dose. In the sleeplessness of acute puerperal or non-puerperal *mania*, there is abundant testimony to the value of chloral.

On the whole, as a pure *hypnotic* chloral is indeed unequalled, and may be used in all cases when no contra-indication exists.

The second indication to meet which chloral may be employed is *to relax spasm*. For this purpose it has been used with advantage in *puerperal* and *uræmic convulsions*. It must be remembered that in many of these cases, although next to chloroform the best palliative, it is only a palliative, and must only be used to quiet the nervous disturbance until other remedies can have time to act.

In *tetanus* it has been claimed that chloral is *the* remedy. Dr. Jos. R. Beek (*St. Louis Medical and Surgical Journal*, June, 1872) has collected thirty-six cases, with twenty-one recoveries, of the traumatic form of the disease, in which it constituted the whole or the major part of the treatment.

References* are given below to eighteen cases in addition; so that the figures stand, thirty recoveries and twenty-four deaths. These results do not seem to warrant the high estimate which has been set upon the value of chloral in tetanus. Very recently, however, Dr. Macnamara (*London Practitioner*, November, 1874) has been employing the remedy in a different way than usual. Believing that it exerts very little control over the spasms, he has not used it for such purpose, but has employed it simply as a hypnotic, giving the patient forty grains of it at bedtime, and only when the temperature rises above 101° F. a single dose of thirty grains in the morning. No other medicine is given, but the patient is made to swallow four ounces of milk with brandy every four hours, one egg being mixed with the milk morning, noon, and evening. In this way he asserts that out of twenty consecutive cases (all traumatic?) occurring in natives of India he has saved seventeen. I think the conclusion from the evidence is irresistible that chloral in tetanus should be used only as a hypnotic, and as such only in the evening.

In *trismus nascentium* Dr. Widenhofer recommends it very highly. He says that in the Children's Hospital he formerly lost all his cases, but that by its use he has saved six out of twelve. He gives it in one- and two-grain doses by the mouth, or, when the spasms prevent, double the quantity by the rectum.

In *chorea* chloral has been used sometimes with great advantage, more often, I think, with the result of simply diminishing temporarily the choreic movements, and sometimes without any effect. In my own experience the movements have generally returned with unabated violence so soon as its exhibition was suspended.

As a nocturnal quietant and hypnotic, it would appear to offer very great advantages in cases of *acute chorea* in which speedy death is threatened from the incessant and violent movements; also in cases complicated with fractures, where a temporary lull is of importance.

In the *convulsions* of children it has been employed with apparent good; in *cramps* in pregnant women it has been commended by Dr. Morgenstern (*Wiener Medicin. Presse*, Nov. 1871); in *singultus*, by Dr. Leavitt (*American Journal of the Medical Sciences*, April, 1871); in the spasmodic *nocturnal enuresis* of children, also, it may often be used with advantage (Dr. J.

* Fergusson (*Edinburgh Medical Journal*, July, 1871), one case: recovery.

Porta (*Schmidt's Jahrbücher*, Bd. cli. p. 110), two cases: death.

Watson (*Lancet*, 1870), one case: recovery.

Bartlett, one case: recovery.

May, one case: recovery.

Ballantyne, one case: recovery.

Cushing (*Pacific Medical and Surgical Journal*, March 1, 1872), one case: recovery.

Itard (*Schmidt's Jahrbücher*, Bd. cli.), one case: death.

Macnamara (*The Indian Medical Gazette*, April, 1871), seven cases: six deaths.

Lovegrove (*British Medical Journal*, Nov. 1872, p. 493), one case: recovery.

Herndon (*Atlantic Medical and Surgical Journal*, 1873, p. 69), one case: recovery.

B. Bradbury, *British Medical Journal*, April, 1871); in *laryngismus stridulus* (Dr. Rehn, *Jahrbuch für Kinderkrankheiten*, 1871), and other spasmodic affections; in *spermatorrhœa* at bedtime it is of service (Gascoyne, *British Medical Journal*, 1872).

In *whooping-cough* it would really seem to be of very great value, as has been attested by Drs. Adams (*Lancet*, i., 1870), Murchison (*Ibid.*, ii., 1870), Rigden (*Practitioner*, xxvii., 1870), Waterhouse (*Ibid.*, December, 1870), and various French observers. Small doses (two to eight grains), repeated at regular intervals during the day, are often very efficacious. Another plan, especially useful when the paroxysms are very severe at night, is to give a full dose at bedtime. When there is a tendency to bronchitis and pneumonia, chloral must be used with care in these cases, as in large doses it favors congestion of the lungs: yet Murchison saw a very threatening case, complicated with bronchitis and pneumonia, greatly benefited by its exhibition.

In *asthma* it has sometimes been of use, but more often it has failed.

The third indication for which chloral has been used is *to relieve pain*. That it will do so when given in very large doses there can be no doubt; but, unless the dose be so large as to be dangerous, my experience of chloral is that it is of very little value as an anæsthetic. Its powers in this direction are incomparably less than those of opium, and its habitual use is probably attended with very grave dangers.

Theoretically, chloral might be of use to *reduce temperature*. Its other active properties will very probably completely interfere with its use for this purpose in the vast majority of such cases, especially as, in order to check the development of animal heat, the dose must be very large. When, however, there is a high sthenic state of the system, it might be tried with caution; but clinical experience is almost entirely wanting. I know of but a single case (*Medical Times and Gazette*, Nov. 1869) reported: in this its use is said to have given satisfaction.

TOXICOLOGY.—That chloral is a dangerous agent, capable of destroying life, is attested by a number of published cases; but this is true of other drugs; and the practical point to be determined is, Does it ever act out of proportion to the amount ingested? or, in other words, does the ordinary therapeutic dose ever become toxic, and does it ever act in a cumulative, unexpected manner? Abroad, it has very commonly been prescribed in half-drachm and even drachm doses, and in the vast majority of cases without any bad results. That thirty grains is not a safe dose is shown, however, by the case of Dr. Reynolds (*London Practitioner*, March, 1870), in which forty-five grains caused most alarming symptoms; by that of Dr. Watson (*Medical and Surgical Reporter*, Jan. 27, 1871), in which eighty grains, given in ten-grain doses spread over thirty-six hours, nearly proved fatal; and especially by a number of cases recorded by Dr. H. W. Fuller (*London Lancet*, March, 1871), in some of which very alarming symptoms followed the exhibition of thirty grains, and in one death in a healthy young woman of thirty. Dr. W. H. Lathrop (*Year-*

Book of Therapeutics and Pharmacy, 1872, p. 254) records the case of a man previously healthy, but suffering from delirium tremens, who took sixty grains between 12 and 1 P.M., at 2.30 P.M. twenty grains more, and at 3 P.M., no effect being manifest, twenty grains more. His physicians then left him sleepless and complaining only of a slight paralysis of the lower extremities; and almost in a moment he was dead. This case, and others, less striking, but similar, would seem to indicate that chloral, if given rapidly, accumulates in the system, and finally kills, very possibly, by a sudden paralysis of the heart. I think the practical deduction from these facts is that twenty grains is the highest safe dose of the remedy; that this amount should not be repeated oftener than once an hour, and, after sixty grains have been taken, not for some hours, unless in very urgent cases, as acute tetanus or violent chorea threatening speedy dissolution. On the other hand, recovery has been reported by Dr. Eshleman (*Philadelphia Medical Times*, Oct. 1870) after the ingestion of four hundred and sixty grains.

The treatment of chloral-poisoning is identical with that of opium-poisoning, consisting in the free use of alcoholic and external stimulants, such as sinapisms, dry heat, frictions, flagellations, etc., to maintain the circulation, and of shaking, walking, application of galvanism, cold douches, etc., to keep up the respiration. Artificial respiration should always be resorted to before natural respiration altogether fails; and Clemens (*Schmidt's Jahrbücher*, Bd. cli. p. 99) has found that animals asphyxiated by chloral may often be at once aroused by the inhalation of oxygen. Hypodermic injections of strychnia have been recommended on uncertain theoretical grounds, but are of doubtful value. Atropia seems to me a much more rational remedy.

Considerable attention has been given both in this country and abroad to the subject of chronic chloral-poisoning; and, whilst some affections have been erroneously attributed to the drug, there seems to be no doubt that its long-continued use often does produce serious symptoms.

Cases of chronic chloral-poisoning are divisible into two or three groups, which are, however, really artificial, as is shown by the occurrence of cases belonging to two or even three of the groups. The first of these includes those patients in whom the respiration is chiefly affected. The dyspnoea may be slight, and may only be felt at times, as after exertion or after meals; but it may be constant and alarming. Cases of this character are reported by Jastrowitz, by Schule, and by Ludwig Kern (*Allgem. Zeitschrift für Psychiatrie*, 1872; *London Practitioner*). In one instance (Prof. N. R. Smith, *Boston Medical and Surgical Journal*, 1871), death from bronchial effusion is believed to have been caused by chloral.

In the second group of cases, eruptions of the skin are the chief manifestations of the toxæmia. In the mildest of these there is no distinct rash, only the occasional appearance of transient red blotches on the face or neck. But a very extraordinary tendency exists towards the production of a rash or discoloration at the slightest cause, so that drinking a glass of wine will produce

an intense, even livid, erythematous redness of the face. In other instances there is marked erythema (Schule, *Allgem. Zeitschrift für Psychiatric*, xxviii.), occurring first in spots upon the face, but extending downwards to the trunk, becoming more and more general, and showing a marked tendency to follow the nerve-trunks. This erythema is seemingly due to vaso-motor weakness, and consequently is allied to other more urgent symptoms seen in chloral toxæmia. Sometimes it invades the mucous membranes, which become red, swollen, and œdematous; and if the glands are involved, as in a case reported by Dr. Chapman (*London Lancet*, 1871), the result may be serious. A deeper implication of the vaso-motor and cardiac nervous system was probably the cause of the general œdema, profound weakness, and failure of heart-action in the case recorded by Prof. N. R. Smith (*loc. cit.*). Prof. Smith also calls attention to desquamation of the cuticle and ulcerations about the nails as being present in these cases.

In the third group of cases, petechiæ, ecchymoses, ulcerations, and even high fever and other pyæmic symptoms, are asserted to have been produced by the continuous use of chloral. It seems to me, however, very doubtful whether the drug really was the cause of the symptoms which have been recorded by Crichton Brown, by Monkton, and by Kern.

ADMINISTRATION.—Sufficient has been said as to the dose of chloral. It is best given diluted with a weak syrup.

CROTON-CHORAL HYDRATE, which has very recently been brought forward as a remedy by Oscar Liebreich, is formed by the action of chlorine gas upon aldehyde. It crystallizes in small glittering tables,* and is soluble with difficulty in water. Although its name would seem to signify it, yet croton-chloral has no relation with croton oil. Chemically, it is a chlorated aldehyde of crotonic acid. According to Liebreich (*British Medical Journal*, Dec. 20, 1873), a drachm of this substance, dissolved in water, and introduced into the stomach, produces in the course of from fifteen to twenty minutes a deep sleep, accompanied by anæsthesia of the head. Whilst the eyeball has lost its irritability, and the trigeminus nerve shows no reaction whatever on being irritated, the tone of the muscles remains unaltered. The effect upon the pulse and respiration is also stated to be much less than that produced by equivalent doses of chloral hydrate. I have not met with any extended experiments upon animals with this drug, but the authority already quoted states that it produces in them deep sleep, trigeminal anæsthesia, and, if in sufficient dose, death by arrest of respiration. The circulation, he affirms, is kept up with great tenacity, and, even if cardiac action, as well as respiration, has ceased, artificial respiration is able to restore the action of the

* For a detailed account of physical and chemical characters, see a paper by E. Schering, *Neues Repertor. für Pharm.*, Bd. xxi. Heft 5, 1872, which I have abstracted into *New Remedies*, vol. ii.

heart immediately, and the life of the animal may thus be saved. Immense doses of croton-chloral produce cardiac paralysis.

Croton-chloral, when subjected to the influence of an alkali, first forms allyl-chloroform, a trichlorated body, which is rapidly decomposed into a bichlorated substance called bichlorallylene. Prof. Liebreich believes, but does not prove, that it is these products of decomposition which act upon the animal.

Croton-chloral has been highly praised by Liebreich for its powers of relieving neuralgias and other painful affections of the trigeminus. He states that it will afford relief even in severe *tic douloureux*, but is, unfortunately, only palliative. His statements have been confirmed by Dr. J. W. Legg (*London Lancet*, 1873) and by Dr. Benson Baker (*British Medical Journal*, Oct. 1873). In a single very severe case of centric tic under my own care, ten grains of the drug have given very decided temporary relief, and compelled sleep. It is usually administered in doses of from ten to twenty grains, in syrup.

AMyli NITRITUM—NITRITE OF AMYL.

This non-official drug was discovered by the French chemist Balard (*Annales de Chimie et de Phys.*, xii.) in 1844, and the attention of physiologists was called to it in 1859 by Guthrie; but it was not until 1865 that Dr. Richardson, of London, introduced it to the notice of the profession. Since that time it has been a good deal used; but it has not been made official. It is a yellowish, oily, very volatile liquid, of a very penetrating, persistent, fruity odor. It is prepared by the action of nitric acid on amylic alcohol, or, as it is commonly called, fusel oil.

PHYSIOLOGICAL ACTION.—Nitrite of amyl can be absorbed by any surface except the skin, but, on account of its volatility, has hitherto been used in man only by inhalation. Owing to the physical property just alluded to, its action is extraordinarily quick and very transient, it being absorbed and eliminated with great rapidity.

The most prominent symptoms induced when it is inhaled by a man in moderate quantities, are a sense of great fulness and distension of the head, amounting at last to severe pain, and accompanied by intense flushing of the face, a deep, labored respiration, and an exceedingly rapid, violent action of the heart. The succession of these phenomena is so rapid that often they seem to be simultaneous; but it is said that the cardiac disturbance is sometimes very distinctly manifest before the other symptoms.

Beyond the point already described the action of the nitrite has rarely been carried in man. In one or two cases I have known alarming prostration to supervene.

In the lower animals the first stage of the action is like that just described in man. After this the breathing becomes violently hurried and panting, progressive muscular weakness and diminution of reflex activity ensue, and finally death from failure of respiration,—sensation and consciousness being preserved almost to the last.

A very peculiar symptom is that a long time before death the arterial blood becomes of almost the same color as the venous blood. Convulsions are sometimes present; but in my experience more often the animal is exceedingly quiet throughout the poisoning.

Elaborate experimental studies of the action of the nitrite of amyl upon the circulation in animals have been made by Dr. T. Lauder-Brunton (*Journal of Anatomy and Physiology*, vol. v.,—*Berichte der Math.-Phys. Classe d. k. sachs. Gesellschaft Wissensch.*, 1869), by myself (*American Journal of the Medical Sciences*, July, 1871), and by Dr. Amez-Droz (*Archives de Physiologie Normale et Pathologique*, Sept. 1873, p. 467). The results are so uniform and in such accord that they must be accepted as proven facts. It has been found by all three observers that, although the pulse is very much increased in frequency sometimes from the very beginning, the arterial pressure is diminished, and finally is reduced almost to zero, and that the fall of pressure occurs equally after section of the vagi as at other times. As the number of heart-beats in the uninjured animal is increased rather than diminished, whilst the strength of the individual beat is not perceptibly lessened, it is evident that, at least in the early stages of the poisoning, the diminution of arterial tension is not cardiac in origin, but must be due to dilatation of the capillaries.

This conclusion is confirmed by an experiment of Brunton, who found that if the descending aorta was tied high up, no perceptible fall of pressure was produced by the inhalation of the amyl salt until very late in the poisoning, when the heart itself was acted upon by the drug; also by the fact noted by Dr. Amez-Droz, that the arterioles, but not the veins, of the rabbit's ear and of the frog's web can be seen to dilate when the salt is inhaled.* An interesting question which here arises is, whether the dilatation is centric, due to an action on the vaso-motor nerve-centres, or peripheric, due to a direct action upon the muscular coat of the arterioles. It must be peripheric, it cannot be centric, since, according to the experiments of Brunton and myself, it occurs when, by division of the cord, the arterioles are separated from the vaso-motor centres. It must be due to a direct paralyzing action of the drug upon the coats of the arterioles,—a conclusion confirmed by our knowledge of the local action of the nitrite upon muscular tissue. That dilatation of the vessels takes place in man as well as in the lower animals, is shown by the flushing of the face, as well as by the enlargement of the retinal vessels, noted by Dr. Chas. Aldridge (*West Riding Lunatic Reports*, vol. i. p. 187). In the latter stages of the poisoning, another factor enters into the causes of lessened blood-pressure, the

* A noteworthy fact asserted by Amez-Droz is that after a long period of dilatation the vessels contract again, whether the inhalation be continued or not. I think the explanation of this is simply that, owing to the volatility of the nitrite, it soon all escapes from the dossil of lint on which it is placed for inhalation: an explanation strongly confirmed by a statement of Dr. Amez-Droz, that in these cases a new inhalation was followed by dilatation as before.

heart-force itself becoming largely extinguished by a direct action of the poison on the muscle.

I have found (*loc. cit.*), as the result of numerous experiments made in the ordinary methods, that the diminution of reflex activity and of voluntary motion which undoubtedly occurs in toxæmia from the agent now under consideration is chiefly spinal in its origin; since after death the nerves and muscles preserve, though in an impaired condition, their functional power. On the motor centres of the cord the nitrite acts as a direct and powerful depressant, at the same time that it exerts a similar but much less pronounced action on the nerves and muscles, decreasing, but not destroying, their functional life. The diminution of reflex activity is never preceded by a stage of functional excitement. In some animals convulsions do occur, especially when the drug is administered by inhalation; but they are in all probability cerebral, not spinal, and due to the asphyxiating influence of the poison. Over the sensory nerves and centres nitrite of amyl has but little power. They are among the last portions of the body to be affected, sensation being intact until near death, so that the drug is in no sense an anæsthetic. The cause of death appears to be due to the failure of power in the cord or respiratory centres higher up.

A very interesting phenomenon of nitrite of amyl poisoning, discovered by Dr. F. A. Hoffmann, is the appearance of sugar in the urine. He found (*Reichert's Archiv*, 1872, p. 747) that in the rabbit a hypodermic injection of 0.111 to 0.113 gramme of the drug is enough to cause diabetes. If twice this amount of the amyl salt is used, the sugar becomes very abundant in the urine, and continues to be present for from twelve to thirty hours. Concurrently with the elimination of sugar, there is a great increase in the amount of the urine.

A study of the local action of the drug shows that it causes a progressive loss of functional power in every highly-organized tissue with which it comes in contact. Nerve-centres, peripheral nerves, muscles of organic and voluntary life, all succumb to it alike. If such contact be not continued too long, the tissue may recover even after a total suppression of its function,—a proof that the nitrite exerts no destructive chemical or devitalizing influence upon the tissues, such as that of sulphuric acid or veratria.

Nitrite of amyl, in whatever way exhibited, reduces most remarkably animal temperature. I have seen a pigeon perfectly conscious, although its temperature had been brought down by this agent some 13° F. This influence is as marked in fever as in the normal condition of the animal, and is independent of the nerve-centres, occurring after section of the cord, and even after death in those cases in which post-mortem rise or continuance of high temperature normally takes place. I have also experimentally determined that it is associated with diminished excretion of carbonic acid. It must therefore be due to a direct arrest or check of tissue-changes or oxidation within, or without, the blood.

The vapors of the nitrite have a very marked influence over oxidation outside of the body, as is shown by many facts, of which it is only necessary here to cite the extinguishment of glowing phosphorus by a few drops of the amyl salt diffused through the jar. It cannot be doubted that within the economy the same thing occurs. If, however, the arrest of oxidation was complete, instant death from suffocation would result. The true explanation of the symptoms evidently lies in diminution, not destruction, of oxidation.

When an animal inhales the nitrite of amyl, the arterial and venous blood soon become of a nearly uniform hue, which resembles somewhat that of normal venous blood, but is quite distinct from it, having a chocolate tint. In my investigations as to the physiological action of the drug, I was unable to find any cause for this change of color; but Dr. Arthur Gamgee (*Philosophical Transactions*, 1868, p. 589) has made a masterly examination into the subject, and thrown much light upon it. He finds that the spectrum of blood treated with nitrite of amyl changes simultaneously with the color. The two sharply-defined absorption-bands of the oxyhæmoglobin become fainter and fainter, and entirely disappear, unless the stratum of examined blood be exceedingly thick, when faint indications of them remain apparent. At the same time new bands appear, precisely resembling those of acid hæmatin. If ammonia be added to the chocolate blood, the color changes back to a blood-red again, and simultaneously the spectrum-lines regain their normal position. If, however, an amount of phosphoric acid precisely equivalent to the amount of ammonia added, and therefore just sufficient to neutralize it, be placed in the blood, the chocolate color reappears, and with it the changes of the spectrum. Moreover, if a reducing agent, such as sulphide of ammonium, be added to the chocolate blood, it is able to deoxidize the oxyhæmoglobin, but, before doing so, evidently removes it from union with the nitrite, since the new bands disappear and those of oxyhæmoglobin reappear in the spectrum before the lines of the reduced hæmoglobin manifest themselves. These facts are, seemingly, explicable only on the supposition that the nitrite makes with the oxyhæmoglobin a compound which is so unstable as to be broken up by ammonia and by reducing agents.

Dr. Gamgee next pushed his researches to find whether the respiratory function of the blood was interfered with by the nitrite,—whether the power of the hæmoglobin to absorb and yield up oxygen was destroyed. By a series of experiments, in which blood was shaken with an accurately-measured amount of air, he found that, whilst normal blood absorbed a large percentage of oxygen, blood to which the nitrite had been added failed to take up an appreciable portion of the gas, even if it (the blood) had been previously saturated with carbonic acid gas. As is well known, carbonic oxide gas has a remarkable power of expelling the loose oxygen of the blood; but it was found that upon blood to which the nitrite had been added it had no influence; and, conversely, that on the florid blood of carbonic oxide saturation the amyl salt had no effect. It was also found that the air-

pump was powerless to draw oxygen out of the blood containing the salt, although the oxygen still existed in the nitrite blood, since reducing agents restored the oxyhæmoglobin spectrum even in blood from which all access to the air was prevented. These facts are in complete accord with those previously spoken of, as showing that the nitrite of amyl unites with the oxyhæmoglobin, to form a new compound. Dr. Gamgee finally furnished the last ocular demonstration by obtaining the compound in crystalline form, by means of a process which it is not necessary to detail here. This nitrite-oxyhæmoglobin has no power of absorbing oxygen, as follows from experiments previously detailed, and as Dr. Gamgee also determined by direct experiment.

These researches of Dr. Gamgee have not, that I am aware of, been confirmed, but I think may be accepted as accurate. They do not, however, prove that nitrite of amyl entering into the blood-vessel at once paralyzes the hæmoglobin of the blood-corpuscle and checks all oxidation. As already stated, the experiments of Dr. Gamgee showed conclusively that this new compound yields up its oxygen to reducing agents. Further, the doctor found that when the nitrite blood was brought into contact with prepared guaiacum-paper it still ozonized it, though not so actively as normal. It is evident that the blood-corpuscles retain to a greater or less degree their power of yielding up ozone to bodies desiring it; that they are capable of exerting at least this portion of their respiratory function; further, where this oxygen is given and the oxyhæmoglobin changed into hæmoglobin, so far as our present knowledge goes, the hæmoglobin must absorb more oxygen before it can reunite with the nitrite. Evidently, then, absorption of oxygen must take place; evidently the blood-corpuscles must perform their respiratory function; but evidently also they are greatly crippled and impaired in the rapidity and ease of its performance. Hæmic respiration is, in other words, greatly interfered with, but not abolished. •

The accord of the results of this chemical investigation with those arrived at by a purely physiological study of the drug is very striking and very beautiful, both teaching the same thing,—lessened, but not absolutely arrested, oxidation.

Having ascertained the existence of diminished oxidation in poisoning by nitrite of amyl, the temptation is very strong to attribute all the symptoms produced by it to this arrest. I do not, however, think that this *post hoc propter hoc* argument is justifiable, for the following reasons: In the first place, the nitrate of potassium and other nitrates, according to Dr. Gamgee, act in the same manner upon the blood, yet the symptoms caused by them are very different from those caused by nitrite of amyl. In the second place, when arrest of oxidation is caused by deprivation of oxygen (see article on Nitrous Oxide) the symptoms are very different, the brain and consciousness being always affected before the centres of reflex action, whereas under the influence of the nitrite of amyl the contrary occurs. In the third place,

other substances, such as toxic doses of alcohol, check oxidation, but do not cause the same symptoms as does the drug under consideration. The obvious inference seems to me to be that nitrite of amyl acts directly upon the nerve-centres, independent of its influence on the blood.

Because, in the first stage of its action on man, there is very marked functional excitement of the heart, it has been held that the nitrite of amyl is a motor-stimulant: a most erroneous idea. The cardiac excitement seems not to be directly owing to the drug, but rather to be reflex in its nature. When the nitrite is taken into the lungs it instantly arrests or diminishes oxidation, and a thrill of impending suffocation runs through the system, in obedience to which the respiratory and circulatory organs gather up and exert to the utmost their forces. The central impulse sent to the cardiac and respiratory muscles is at first much more than sufficient to overcome any direct action of the nitrite upon them; but the inhalation being persisted in, the impulse is constantly growing weaker and the direct influence of the drug stronger, so that there soon comes a time when the reverse is true, and the heart's power is more or less nearly extinguished.

THERAPEUTICS.—The results of the clinical use of the nitrite of amyl are in accord with what has been said of its physiological properties. Its administration in *angina pectoris* appears to have been first suggested to Dr. Brauton (*London Clinical Society's Reports*, vol. iii. ; *Lancet*, July 27, 1867) by the sphygmographic tracings giving evidence of arterial spasm in a case of that disorder. The pathology of these cases of heart-pang is not definitely made out, and so it seems useless to speculate how the nitrite acts in many cases;* but there is now abundant evidence of its value in relieving almost instantly agony which has resisted all other treatment. This appears also true whether valvular disease or merely functional disorder exists. In cases of advanced fatty degeneration, or of very great dilatation of the heart, I think its use would be attended with danger, owing to its effect upon the heart-muscle.

Its physiological action would indicate that it should be of service in all cases of spasm of the capillaries, of the bronchial tubes, and of the muscular system generally. Accordingly, Dr. Oskar Berger (*Allgem. Medicin. Central Zeitung*, May, 1871) has used it with very good effect in a case of *migraine* with evident capillary contraction. In *asthma* my own experience of several cases coincides with that of others, that it will often instantly arrest the paroxysm,† especially in those instances in which there are no secondary lesions, such as emphysema and dilated heart.

The convulsion of *epilepsy* is, according to the present theory, due to a vaso-motor spasm at the base of the brain, to correct which the amyl salt would seem to be indicated, as it also is by the mere existence of the convulsion. In advanced stages of the paroxysm it must, however, be used with very

* Case of failure, see *London Lancet*, August, 1867.

† For a remarkable case, see *British Medical Journal*, Sept. 30, 1873.

great caution, if at all, on account of the obscuring of its early effects by the symptoms of the disease. When there is a notable interval between the aura and the convulsion, it would appear that the latter could be entirely prevented; at least such a case is reported by Dr. S. Weir Mitchell (*Philada. Med. Times*, 1872, vol. ii. p. 262), who directed with entire success the patient to carry a small vial containing a few drops of the drug, and to inhale from this whenever the aura was felt.

Dr. Wm. F. Jenks reports a case of *convulsions* occurring immediately after delivery, in which the paroxysms were instantly arrested by the inhalation of a single drop of the nitrite as they were coming on. Its use, however, was followed by relaxation of the firmly-contracted uterus and very alarming hemorrhage, a result to be expected from the known physiological action of the drug (*Philada. Med. Times*, 1872, vol. ii. p. 404).

In *tetanus* the nitrite of amyl would seem to be indicated as a spinal sedative, and as controlling excessive tissue-changes and consequent rise of temperature. It has been used, so far as I know, in only two cases (*London Lancet*, 1871), both of which recovered, one with the amyl salt alone, and the other with it and chloral.

Although its sedative action would seem to contra-indicate its use, yet, as the most powerful known drug for checking the development of animal heat, I conceive a cautious use might be made of it in the so-called "*cerebral rheumatism*" where death is threatened from excessive temperature.

The physiological action of the remedy would very strongly indicate it as an antidote in *strychnia*-poisoning. No case of its use for this purpose in man has come to my knowledge, but the experiments of Dr. St. Clair Gray (*Glasgow Medical Journal*, p. 188, 1871) on rabbits yielded very favorable results, although they were not sufficiently elaborate to decide the matter. In two rabbits to each of which ten drops of the nitrite and half a grain of strychnia were given together subcutaneously, no decided symptoms whatever were induced; whilst one-quarter grain of the alkaloid alone frequently caused death in a single convulsion.

ADMINISTRATION.—As already stated, the only method of administration hitherto employed is by inhalation, from one to three or five drops being placed on a handkerchief and held near the mouth or nose, the handkerchief being removed so soon as a sense of fullness of the head is experienced. Used with care, the nitrite, although a very rapidly-acting and powerful agent, seems to be safe, since I have never seen either in man or animals any sudden or unexpected action,—any influence apparently out of proportion to the amount given. It must be borne in mind that the symptoms generally increase in intensity for a minute or so after the withdrawal of the drug.

LOBELIA.

The leaves and tops of the indigenous herb *Lobelia inflata*. The dried plant has a slight irritating odor and a taste at first scarcely perceptible, after-

wards burning, acrid, and attended with a flow of saliva. The active principle is *Lobelia*, a yellowish liquid alkaloid, discovered by Prof. Procter, of this city. *Lobelic* acid, a fixed and a volatile oil, gum, chlorophyl, etc., are also present in the drug.

PHYSIOLOGICAL ACTION.—*Lobelia* appears to have the same influence upon the lower animals as upon man. At least Prof. Procter found a grain of the alkaloid to produce in a cat violent emesis, with intense prostration. In man, the symptoms induced by it when freely administered are nausea, soon followed by violent vomiting, accompanied with intense prostration, as is shown by feeble pulse, cold sweats, pale skin, and great muscular relaxation. Purging may or may not occur. Numerous cases of fatal poisoning by it have been recorded. The symptoms are those above mentioned, intensified; in some cases vomiting does not occur, and it is especially under these circumstances that fatal effects have been noted. Burning in the fauces and œsophagus, epigastric distress, in addition to the intense prostration, bordering upon collapse and finally merging into complete collapse, with coma, stupor, muscular tremblings, and in some cases convulsions, precede the fatal termination.

The exact method in which *lobelia* affects the system has not been made out, but it resembles very closely tobacco in its effects, and probably acts both as a cardiac and a spinal depressant.

TOXICOLOGY.—The symptoms of *lobelia*-poisoning have been sufficiently described. The treatment should consist in washing out the stomach with plenteous draughts of a warm solution of tannic acid, in the free exhibition of opium and of alcoholic and ammoniacal stimulants, and the use of external stimulation by dry heat, frictions, mustard, etc., precisely as in poisoning from *veratrum viride*.

THERAPEUTICS.—*Lobelia* has been used as an emetic; but its depressing effects are so severe as to forbid such employment of it. It has also been employed to relax spasm in various affections, as in pertussis, tetanus, epilepsy, chorea, convulsions, but has been superseded by more efficient and less dangerous remedies. It is often useful in spasmodic *asthma* or in acute *bronchitis*, with bronchial spasm, and appears to be expectorant as well as anti-spasmodic. An infusion (℥i to Oj) has been strongly recommended as a local application in the eczema produced by the *rhus toxicodendron* or "poison-vine." The powder is very rarely used; the dose as an emetic is twenty to thirty grains. The official preparations are as follows:

Tinctura Lobeliæ—*Tincture of Lobelia* (℥ii to Oj). Dose, as an expectorant, gtt. x to xx; in the paroxysm of *asthma*, f℥i to f℥ii every half-hour until nausea is induced.

Acetum Lobeliæ—*Vinegar of Lobelia* (℥ii to Oj). Dose, same as that of the tincture.

CONII FOLIA—CONIUM LEAVES.

The leaves of *Conium maculatum*.

CONII FRUCTUS—CONIUM FRUIT.

The full-grown fruit gathered while yet green and carefully dried.

Conium is an umbelliferous plant, native of Europe, and naturalized in the United States. The dried leaves have a strong heavy odor, increased by the addition of an alkali, and resembling somewhat that of mice. They are bi- or tripinnate, and very much incised. The fruits are one to two lines long, roundish ovate, striated, with five crenated ribs on the outer sides of the easily-separable halves; the odor is that of the leaves. The active principle is *coniin*, a yellowish, oily, liquid alkaloid, highly volatile, of a strong odor similar to that of the urine of mice, a very acrid taste, freely soluble in alcohol and ether, slightly so in water, with which it forms a hydrate. It coagulates albumen; when exposed to the air it undergoes decomposition, becoming first brown, afterwards resinous; heat accelerates the change.

PHYSIOLOGICAL ACTION.—The chief symptom produced in man by conia when taken in doses just large enough to impress decidedly the system is great muscular weakness or languor, with some disorder of vision and giddiness. On attempting to walk, the patient suffers from a feeling as though his feet were made of lead, and he staggers or falls from the refusal of his knees to support him. There is an intense desire to lie quiet in the horizontal position, and, as the eyelids are especially affected, the eyes are kept shut. In some subjects these symptoms are preceded or accompanied by burning in the mouth or fauces, nausea, and even vomiting, besides heat of head, often with a sense of weight or pressure, or even severe frontal pain. The disorder of vision is apparently due in great part to a sluggishness and finally to a paralysis of accommodation. The experiments of Poehlmann (quoted by Husemann, *Die Pflanzenstoffe*, p. 269) show that very grave symptoms may be induced and yet the pupil remain natural; but sooner or later, as the drug gains power over the system, it probably always dilates. The pulse is first diminished, afterwards increased in frequency. In decided poisoning by conium, the symptoms are probably simply those already mentioned, intensified. I have met with accounts of but two fatal cases of such character. In one, that of the mistress of Dr. Hermann Jahn, killed in a few minutes by from ten to fifteen drops of the alkaloid (quoted by Husemann, *Die Pflanzenstoffe*, p. 269), violent palpitation of the heart is said to have been a prominent symptom. The other case, reported by Dr. Bennett (*Edinburgh Med. and Surg. Journal*, 1845), is that of a man fatally poisoned by eating the fresh plant for "greens." The prominent symptom in his case was universal paralysis, with total failure of voluntary movement and of the voice before consciousness was lost. Convulsive movements were present very late in the case. Sensation appeared not to be lost until death was at hand.

In animals conia produces symptoms parallel with those observed in man, and it probably acts similarly upon all vertebrates. In some animals, however, the convulsions are more prominent than in man; in frogs they are rarely if ever present; in birds they are occasionally so; in mammals they are more frequent,—thus, Ihmsen saw them in twelve out of twenty-three experiments; they are chiefly clonic, but tonic spasms do occur in the hind legs. As in man the legs are usually affected before the arm, so in quadrupeds the hind extremities are usually paralyzed first. Sensibility is maintained to the last. The respiration is generally much affected, and the heart continues to beat after its cessation.

Conia is certainly absorbed, as it has been found by Orfila in the spleen, kidneys, and lungs of poisoned animals, and it is probably eliminated from the system. Its volatility makes its escape through the lungs probable, and it has been detected by Zaleski and Draggendorff in the urine.

All observers agree that the chief symptom produced by conia (*i.e.*, the paralysis) is not due to any direct influence upon the muscles, which, indeed, preserve perfectly their contractility up to death. In 1856, Kölliker (*Virchow's Archiv*, Bd. x. p. 228) announced that the failure of motion in conia-poisoning is due to a direct action of the alkaloid upon the afferent or motor nerves. He first experimentally found that in frogs killed by the drug the application of the galvanic current to a nerve fails to induce contractions in the tributary muscles. He then tied the aorta in such a way as to cut off the supply of blood to the hind extremities, and found that after voluntary motion had ceased in the fore legs, and even after galvanic stimulation of the anterior nerves had lost its influence upon the muscles directly supplied by those nerves, irritation of the same anterior nerves produced reflex contractions in the hind legs, showing that the anterior *afferent* nerves and the spinal cord still retained functional activity after the loss of it in all those afferent nerves reached by the poison. After repeating these experiments a number of times, he drew the conclusion already given.

His experimental results have been confirmed by Funke (*Berichte über die Verhandlungen der k. sachs. Gesellschaft der Wissenschaften zu Leipzig*, Bd. xi. p. 23, 1859), by Guttmann (*Berlin. Klin. Wochenschrift*, 1868, quoted by Husemann), and by MM. Pelvette and Martin-Damourette (*Gazette Méd.*, 1870, quoted in *Archives Générales*, 6e sér., t. xvi. p. 88). The latter observers extended the series by severing in a frog all the tissues at the upper part of the thigh except the nerve, and found that when a batrachian so prepared was poisoned with conia, after the paralysis was complete in all portions of the body to which the poison had access,—after stimulations of the poisoned nerves were powerless to excite contraction in the tributary muscles,—the leg which had been protected from the action of the conia upon it responded not only to irritations applied to its nerve, but also to stimuli placed upon distant portions of the body. These same observers also noted that when conia and strychnia were given simultaneously to a frog from one of whose

sciatic nerves the circulation (*i.e.*, direct access of the poison) was cut off in either of the manners spoken of, they produced by their conjoint action a commingling of paralysis in all other parts of the body with violent tetanic spasms in the protected leg,—a commingling explainable only on the supposition that the conia paralyzed all the motor nerves to which it had access through the circulation. As Verigo (*Schmidt's Jahrbücher*, Bd. cxlix. p. 16) has also experimentally demonstrated this action of conia upon the afferent or motor nerves, it must, I think, be considered settled that conia is a paralyzer of these nerves. It also evidently follows from the various experiments which have been detailed that it is not the trunks but the extreme peripheral ends of the motor nerves which are primarily affected.

What its influence upon the spinal cord may be, or whether, indeed, it have any influence, must yet be considered undetermined. Dr. Verigo (*Schmidt's Jahrbücher*, Bd. cxlix. p. 16) asserts that it acts very forcibly upon it as a depressant, and MM. Pelvette and Martin-Damourette (*Archives Générales*, 6e sér., t. vi. p. 89) say it acts as an excitant. I have not had access to the original papers of these investigators, and am unable, therefore, to offer judgment upon their evidence.

Apparently upon the foundation of a single, very crude, and inconclusive experiment, Dr. Harley has recently advanced the opinion that conia affects chiefly the corpora striata, and the other centres at the base of the brain, supposed to mediate between the will and the spinal cord. As Dr. Harley brings forward no *proof* of the truth of his theory, it is contrary to the plan of this work to enter into any discussion of it. It should be stated, however, that Drs. A. D. Davidson and D. Dyce Brown, in support of the views of Harley, cite an experiment of their own, in which one femoral artery of a young cat was tied, and yet the limb lost power as rapidly as the others when conia was given. This single experiment is opposed by so many as to make it almost certain that some fallacy underlies the result. It may be that the alkaloid was carried to the nerve by the collateral circulation.

Upon the sensitive nerves conia in therapeutic doses has no influence, and even lethal amounts of it scarcely impress them. The pupil is generally dilated by it, but both Von Praag (*Journ. für Pharmakodynamik*, Heft i. p. 31) and Verigo assert that the phenomenon is not constant, at least in animals. The ptosis of conia-poisoning indicates that the dilatation of the pupil is due to oculo-motor paralysis. The known action of the drug upon nerve-trunks indicates that this paralysis is peripheral, a conclusion corroborated by the experiments of Dr. I. Hoppe (*Die Nervenwirkungen der Heilmittel*, Heft i., Leipzig, 1855), who found that when conia was dropped into the eye of an animal it caused at first contraction, apparently due to the intense irritation, and afterwards dilatation, of the pupil.

Lethal doses of conium cause a decided lowering of temperature (Verigo, Von Praag, and others). Although Harley failed to find the conia in the urine of animals poisoned by it, yet it is eliminated by the kidneys, in whose

secretion Zaleski and Draggendorff have found abundance of it in the first twelve hours of the poisoning, and traces of it for two days and a half.

Upon the circulation conia seems to exert very little influence.

When locally applied in a concentrated condition, conia probably is fatal to all the more highly organized tissues. Certainly Christison (*Edinburgh Philosophical Transactions*, vol. xiii.) proved this to be so in regard to the muscles which are not influenced by it when taken internally. Upon the mucous membranes it acts as an intense irritant.

THERAPEUTICS.—The paralytic action of conia naturally would suggest its use in spasmodic affections; and accordingly it has been tried in *chorea*, in *paralysis agitans*, in *whooping-cough*, and other diseases of similar nature. Although it seems not to have met with continued favor, and is but little used, when life is threatened by the mere convulsive actions, as it will suspend these for the time being, it may be employed. If Dr. Harley's views as to its physiological action be correct, it ought to be especially useful in all motor disturbance connected with irritation at the base of the brain. Clinical proof is, however, nearly as scarce as physiological in this matter.

In the treatment of the insane, conia is very much used by some alienists.

Conium has also been employed to relieve pain, but it is evident that it can do this in an indirect manner only. As a deobstruent and alterative it has been very largely used, both locally and internally, in *cancerous* and other *tumors*, in *chronic glandular enlargements*, *swollen joints*, and in various *chronic ulcerations*.

It has also been employed to arrest the secretion of milk.

These various uses of conia as an alterative certainly have no definite physiological basis, but they appear to be justified to some extent by clinical experience.

ADMINISTRATION.—One of the great practical drawbacks to the use of this drug is the uncertainty of its preparations; none of them can be relied upon. Of the leaves the U. S. Pharmacopœia recognizes two extracts, the fresh juice (*Extractum Conii*) and the alcoholic (*Extractum Conii Alcoholicum*).—dose, one to two grains, increased until some symptoms are induced; also a tincture (*Tinctura Conii*, ℥ii to Oj).—dose, $\text{f}\overline{3}\text{ss}$ to $\text{f}\overline{3}\text{i}$. The official *fluid extract* is made of the fruit (*Extractum Conii Fructus Fluidum*). Dose, ℥j to ℥ii . Of these preparations, the last appears to be the only one which can be at all relied on.

The English *Succus Conii*, so praised by some writers, I have known to be used by the ounce without effect.

This variability has its origin first in the varying amount of the active principle in the drug; secondly, in the great volatility of this principle; and thirdly, in the proneness of the alkaloid to undergo spontaneous decomposition, even when kept under the most favorable circumstances, and to a much greater extent when exposed to light and air.

The alkaloid, on the whole, would probably be the best form in which to

use the remedy, but it is not officinal; abroad, however, it has been used to a considerable extent, and has generally given satisfaction.

The dose is one-twentieth to one-twelfth of a grain, which may be dissolved in alcohol.

TOXICOLOGY.—Sufficient has been said about the symptoms caused by conia. After death from it no distinctive lesions are to be found, only the usual indications of death from asphyxia. The treatment consists in the immediate evacuation of the stomach and the exhibition of tannic acid,—the tannate formed is, however, probably more or less poisonous,—with the use of external heat and of internal stimulants; artificial respiration should be steadily maintained so long as there is the faintest indication of cardiac action. No physiological antidote is known; but it is possible that atropia might be of service by aiding to maintain the respiration.

CLASS XI.—ALTERATIVES.

THERE are employed by practitioners of medicine, to affect certain diseases most intimately connected with the processes of nutrition, various substances which do not, at least in the doses commonly employed, produce any very obvious symptoms. These drugs may perhaps neither stimulate nor depress, so far as can be perceived, any function of the body; their action may be silent and imperceptible, their mode of influence may be unknown, but their therapeutic effects are among the most assured of clinical facts. It is to medicines of this character that the name of *Alteratives* has been applied, because when administered they seem simply to alter morbid processes.

Speculation has been rife as to the mode in which alteratives influence the body; and as the accepted pathology has been humoralistic or otherwise, so has it been strenuously argued that they act upon the vital fluid or upon the solids of the body. The term "purifying of the blood" has been especially applied to their action, and is sufficiently suggestive of their function as viewed from the pathological stand-point of the old humoralist.

What we know of the action of these medicines at present amounts to this, that they modify the nutritive processes of the body. As the *physiologist* has scarcely learned the alphabet of that part of his science which treats of the general nutrition, having no knowledge as to what is the real dominant force in the nutritive processes; as he is unable to tell with certainty in what organs the universal vital pabulum, the blood, is formed, much less to point out the method of its formation and the laws which govern its development; as the *pathologist* is completely baffled in attempting to find the essence, as it were, of the morbid processes which are successfully met by alteratives; as he cannot point out to us what perverted functions underlie these diseases as their basis,—why should the *therapeutist* be expected to explain the rationale of his treatment? The empirical facts of the clinical pathologist are met by the empirical facts of the clinical therapeutist. It is absurd to gaze into mid-air for the crowning spire before the foundation-stones of the temple are laid.

To deny, as has been done, the existence or value of medicines of this class because we cannot tell why mercury relieves syphilis or why iodide of potassium cures rheumatism, is as absurd as to deny the existence of the syphilitic and the rheumatic dyscrasia because we do not know their ultimate nature. Let us be content, until more light comes, to hold fast to the clinical

facts, each believing for himself, if he choose, that alteratives alter nutrition by affecting the functions of the blood-making organs, or, if he prefer, that they act by impressing the cells of the body directly; or, what is even more philosophical, each holding his mind free from belief, an unoccupied tablet on which the truth may readily be inscribed when it is discovered.

ARSENICUM—ARSENIO. (As.)

Although *Arsenic* is the officinal title for the metal which forms the base of arsenious acid, yet the term has come into such universal use as a synonym of *white arsenic* or *arsenious acid* that I shall here so employ it.

Arsenicum is a brittle metal, of a steel-gray color and a brilliant lustre when freshly broken, but becoming dull and dark on exposure. Its odor when volatilized is that of garlic.

ACIDUM ARSENIOSUM, *Arsenious Acid*, *White Arsenic*, *Arsenic* (AsO_3 — As_2O_3), as first prepared by sublimation from the ores, is in transparent masses, but on keeping becomes milk-white externally. It is soluble in water, has a vitreous fracture, is odorless, of a faint sweetish taste, and volatilizes without fusion “at a temperature not exceeding 400°F .” When it is put upon red-hot iron it emits a garlicky odor, owing to its being reduced to a metallic state and to the volatilization of the metal thus formed.

PHYSIOLOGICAL ACTION.—When applied to any part in a concentrated form, arsenic is a very active escharotic, and even when very much diluted it is a severe irritant. When a single dose of just sufficient size to be felt is ingested, colicky pains, diarrhoea, and perhaps nausea result. After a very large toxic dose, in from one-quarter to three-quarters of an hour an intense burning pain is felt in the œsophagus and stomach, soon spreading to the whole belly, and often accompanied by a sense of constriction at the throat, and an acrid, metallic taste. In a very short time violent vomiting and purging come on. The matters rejected are at first mucous, and variously colored by the contents of the primæ viæ; but they soon become bilious, and often yellowish or greenish, and finally serous, with mucoid flakes and a greater or less amount of blood.

The symptoms above detailed increase in intensity, and to them are soon added others of different nature. The thirst is excessive; the urine suppressed; the extremities icy cold; the pulse small, feeble, and frequent; the respiration very much embarrassed, painful from the abdominal tenderness, rapid, labored; the surface dark and cyanosed; violent cramps add their torture to the other affections; exhaustion deepens into collapse, convulsions or coma ensue, and death occurs in from five to twenty hours.

In another set of cases, when the dose has been smaller or the subject less susceptible, the termination is not reached so soon. After symptoms similar to but less violent than those just described have lasted from a few hours to one or two days, a remission occurs; the purging and vomiting grow less fre-

quent, or perhaps intermit; even the abdominal tenderness may in great measure disappear, but the persistent thirst, cold extremities, and suppressed urine show that the danger is not overpast, and after a time the case puts on a more alarming aspect. The belly becomes very tumid, the abdominal pain more severe, difficulty of respiration develops itself, the face is swollen and cyanosed, nervous symptoms, tremblings, cramps, and convulsions appear, and finally an icy coldness pervades the frame, and death occurs in from two to six days. The mind is generally clear to the last.

If the patient survive long enough, an eruption very frequently appears, sometimes as early as the second, sometimes not until the fifth, day. Its character is various; thus, it may be petechial, urticaria-like, papular, vesicular, or pustular (*Histoire des Éruptions arsénicales*, par Dr. Imbert-Gourbeyre, *Moniteur des Hôpitaux*, 1857).

Such are the ordinary symptoms of arsenical poisoning; but anomalous cases are not rare. Sometimes profound and rapid collapse, without abdominal pain, has occurred; in other cases heavy sleep, deepening into coma, is said to have been the most marked symptom. Sometimes arsenical poisoning resembles most closely cholera, and has been mistaken for it not only in life, but also after death, on the post-mortem table (*Virchow's Archiv*, 1870, Bd. 1. p. 456).

When arsenical poisoning is not fatal, the convalescence is very apt to be slow, and interrupted by various disorders. Prominent among these are affections of the alimentary canal, due to the structural changes produced by the poison. Nervous symptoms also are not rare. They are chiefly neuralgic pains, anaesthesia, and paralysis. The latter affects preferably the lower extremities, commencing and remaining longest in them; does not select the exterior muscles, and is almost always accompanied by anaesthesia, or at least by numbness and formication, and by coldness of the extremities (Dr. Leroy d'Etoilles, *Gazette Hebdomadaire*, 1857, iv.).

The most obvious lesions found after death from acute poisoning by arsenic are in the stomach and bowels, even when the poison has found entrance through other channels into the system. The gastric mucous membrane is usually swollen, maculated with patches of a deep crimson or, more commonly, brownish-red color, and is often softened and covered with a diphtheritic exudation, but is rarely ulcerated. Perforation is exceedingly uncommon. The mucous membrane of the upper part of the small intestine, and sometimes of the whole of it, is in a condition similar to that of the stomach. In some cases the lesions resemble very closely those of cholera, as was first pointed out by Prof. Virchow (*Virchow's Archiv*, Bd. xlvii.). In the microscopic examination of a cadaver whose bowels were filled with a "rice-water" fluid, that observer found in the intestinal contents epithelial flakes, and the fungus described by Klob as peculiar to, and, indeed, the cause of, cholera. The epithelial cells of the mucous membrane were choked with granules, and many of them in an advanced stage of fatty degeneration; the interstitial tissue was full of large round granulated cells; the solitary glands and Peyer's patches

were very much swollen. These facts have been confirmed by Dr. Höffmann (*Virchow's Archiv*, Bd. l. p. 456).

It can scarcely be doubted that in arsenical poisoning there is a widespread granular or fatty degeneration of the tissues. The evidence as to man is not so complete as is desirable, but M. Karajan (*Tardieu, Sur l'Empoisonnement*, p. 335) reports in one case, which had been mistaken during life for acute atrophy of the liver, that the viscus was atrophied and of a characteristic fatty color; and Fr. Grohl and Fr. Mosler (*Virchow's Archiv*, Bd. xxxiv. p. 213) have recorded the case of a boy poisoned with arsenic, in whom they found fatty or granular metamorphosis of the glands and epithelium of the stomach and intestines, of the cardiac muscle, of the diaphragm, and, to a slight extent, of some of the voluntary muscles. In this case the kidneys were also profoundly affected; the cortical tubes were opaque and finely granular, and their epithelial cells could not be isolated.

The absolute demonstration of the degeneration produced by arsenic was, however, made by Dr. Saikowsky (*Virchow's Archiv*, Bd. xxxiv. p. 77), of Moscow, who was also the first to point it out. In his numerous experiments upon rabbits he found that when the animals were poisoned by a small dose of arsenic, so as to live from three to six days, the liver was much enlarged and very fatty,—indeed, contained more fat than the “phosphorus-liver.” On microscopical examination, the cells on the exterior of each acinus were seen to be natural; those in the centre in the most advanced stages of degeneration. The kidneys were similarly affected,—their tubes choked up with fat-globules, their epithelium almost completely destroyed. The muscles of the heart and diaphragm were almost equally compromised. Dr. Saikowsky also noted that early in both arsenical and antimonial poisoning the glycogenic function of the liver is abolished.*

The action of arsenic when given in poisonous doses to frogs has been studied by Dr. W. Sklarek, of Berlin (*Reichert's Archiv*, 1866). He found that the arseniates of potash and of soda had exactly the same effect as the arsenic itself. Within five minutes after the injection of one-fourth to two c.c. of a two per cent. solution, all voluntary movement ceases in the frog; but when the animal is laid upon his back he struggles very actively to recover his position. At this time, however, all sensibility to chemical and mechanical irritants is lost, cutting, burning, or corroding failing to elicit any response. That the motor system is not at fault is shown by the active movements when the frog is placed upon his back, as well as by the results of electrical stimulation of the nerves. The paralysis or quietness must be due to an abolition of sensation. That this is spinal, and not peripheral, is proven by the circumstance that tying the iliac artery upon one side before the administration of the poison has no effect in preserving sensibility in the protected leg. The

* For a spectroscopic study of the effect of arsenic upon the coloring-matter of the blood, see *Centralblatt*, 1868, p. 609. It is interesting here to note that arsenic, antimony, phosphorus, and ammonia act very similarly, if not identically, upon the blood.

only explanation of the struggles of the frog to recover his position after poisoning is to be found in his being influenced through vision, or else in the theory that the muscular sense is distinct from that of common sensibility and is not affected by arsenic.

Upon the heart both of the frog and the cat Dr. Sklarek found that arsenic exerts a very powerful influence, lessening the rapidity and force of the beat, and finally arresting the contraction. That this cardiac action of arsenic is direct is shown by the fact that the phenomena produced by the application of arsenic to the heart cut out of the body are the same as those seen in the arsenical poisoning. The arrest is never instantaneous, but always preceded by slowing of the beat; and after movement has ceased, galvanic or mechanical irritation will cause imperfect systolic movements.

In no case did Dr. Sklarek observe any signs of functional excitement preceding the development of the cardiac or motor paralysis.

As arsenic is never used in medicine for an immediate effect, the chief interest to the therapist centres around its action when administered in small repeated doses. Minute quantities of it may be given for a long time without producing any perceptible effect, unless it be a sharpening of the appetite, due to the local action on the stomach. When the dose is increased, more active manifestations of gastric irritation may appear, such as loss of appetite, nausea, abdominal pain or uneasiness, diarrhœa, and perhaps sympathetic headache. By the use of frequent small doses, these symptoms may be generally avoided, and what may be termed the constitutional action of arsenic be obtained. The first sign of this is generally a puffiness about the eyes, at first visible only in the early mornings, but soon increasing into decided œdema, which after a time may lose its local character and the patient be involved in general anasarca. This anasarca, as was, I believe, first pointed out by Dr. S. Weir Mitchell (*New York Medical Journal*, vol. i.), may or may not be preceded or accompanied by the presence of albumen and of tube-casts in the urine. Beyond the production of the symptoms spoken of, arsenic should never be employed in medicine.

Unfortunately, owing to the frequent use of the metal in the arts, chronic poisoning with it is by no means uncommon.

Although the symptoms vary a good deal and are often obscure, yet in almost every case they are such as should at once awaken suspicion. They have been summed up by Prof. Taylor as follows: "Dryness and irritation of the throat, irritation of the mucous membranes of the eyes and nostrils, dry cough, languor, headache, loss of appetite, nausea, colicky pains, numbness, cramp, irritability of the bowels, attended with mucous discharges, great prostration of strength, a feverish condition, and wasting of the body." It is very evident that the symptoms of irritation of the respiratory mucous membrane are largely, if not entirely, due to the local action of the arsenic, since the poison finds access to the system through the respiratory organs. The constitutional troubles most uniformly present in these cases are weakness and

emaciation, often accompanied by more decided nervous manifestations than the picture drawn by Dr. Taylor would suggest; great depression of spirits and irritability of disposition, sleeplessness, muscular tremors, vertigo, and even convulsions and paralysis, are very common.

Dr. Kirchgässer, as the result of very large experience, asserts that the most characteristic phenomena of chronic arsenicism are a brown pigment deposit in the skin of the face, inflammatory affection of the eyelids, and the disturbances of sensibility and motion, which affect most frequently the lower extremities, together with scalding during urination. Out of eight cases, he found arsenic in the urine in six (*Berlin. Centralblatt*, p. 574, 1868).

Opposed to this every-day experience is the asserted "arsenic-eating" of the peasants of Styria. Originally affirmed by Vogt (*Lehrbuch der Pharmacodynamik*, dritte Aufl., Bd. i.), the existence of this practice has been especially asserted by Tschudi (quoted by Prof. Stillé), and more recently by Chs. Heisch (*Pharmaceutical Journal and Transactions*, vol. i., 2d series, 1859 and 1860, p. 556). It is stated by some that the arsenic is taken by the young girls to beautify their complexion and to enhance their charms; by others, that the object to be attained is protection against arsenical fumes by those engaged in the manufacture of the metal, and the increase of the powers of endurance and of the "wind" in huntsmen and others who do a great deal of mountain-climbing. The habit is said not to be detrimental to life. Indeed, the toxiphagi are asserted to be remarkably long-lived people. In regard to the dose, three grains is said to be taken as a commencement, and to be increased to thirty!

We have not here space for a thorough discussion as to the asserted arsenic-eating. A reviewer in the *British and Foreign Medico-Chirurgical Review* (vol. xxix. p. 144) sums up very elaborately the evidence to the date of his article, 1862. He shows, I think conclusively, that, whilst a good deal of evidence as to the existence of the habit had, up to the date mentioned, been brought forward, yet it was mostly hearsay, and far from being sufficient to overcome the inherent improbability of the practice. Recently, however, very strong evidence has certainly been educed. According to the *Boston Medical and Surgical Journal* (p. 434, 1873),* Dr. MacLagan not only saw how well-known arsenic-eaters take their dose of several grains, but had some of their urine collected, from which the drug was subsequently recovered by a chemist.

An unsigned (editorial?) communication in the *Edinburgh Medical and Surgical Journal* (vol. xvi., 1871, p. 569) further asserts that a royal commission has examined into the subject, and their report indicates that the practice exists, but that it has been grossly exaggerated. They affirm that arsenic-eating is practiced chiefly in the northern and northwestern parts of

* The *Boston Medical and Surgical Journal* refers to the *Edinburgh Medical and Surgical Journal*, Sept. 1867, for Dr. MacLagan's article. No such paper is at that reference, and I have been unable to find it elsewhere.

Styria; that the white arsenic is preferred, the yellow commercial article being sometimes taken, the native red arsenic, or orpiment, very rarely; that the commencing dose is about 0.22 of a grain, which is very slowly increased up to 0.62 grain avoirdupois. The "ratsbane-eaters" almost all belong to the lower classes, and are said to be generally strong and healthy persons, courageous, pugnacious, and of strong sexual disposition.

When arsenic is administered in small repeated doses, it may act as a tonic, by slightly irritating the stomach and thereby provoking an appetite; and in certain cachexias it does increase the muscular strength and the general vigor. I do not, however, see any reason for believing it to be a tonic in the usual sense of the term: the amelioration, the increase of strength and blood in the cases alluded to seem to be due to an indirect action of the drug, to a removal or overcoming of the morbid agent on the disease, and a consequent allowing of the recuperative powers of the system to assert themselves. Indeed, all that we know of the effect of arsenic upon the system seems to throw only enough light upon its therapeutic action to enable us to class it as an alterative—a modifier of nutrition.

There is abundant proof of its absorption, and no less of its elimination. Indeed, the latter appears to take place with exceptional rapidity. Thus, MM. Flandin and Danger (Husemann, *Toxicologie*, p. 823) failed, three days after the last dose, to detect it in the bodies of animals to which fifteen grains had been given daily; and in a child killed in two days by an arsenical pigment, none of the metal could be found in the body (*British and Foreign Med.-Chir. Review*, 1870, vol. xiv.). In the great majority of instances, however, there is no trouble in finding arsenic in the bodies of those poisoned by it, and Steinhäuser reports a case in which it was found in the remnants of a corpse that had been buried for twenty-two years (*Berlin. Centralblatt*, 1868, p. 160). The principal channel of escape is the kidneys, but elimination also takes place through the mucous membrane of the alimentary canal, through the skin, and even in the saliva and tears. M. Chatin has found it in the serosity of a blister, Bergeron and Lemaître in the sweat (*British and Foreign Med.-Chir. Review*, vol. xlviii., 1871), and Taylor (*Guy's Hospital Reports*, vol. xi., 3d series, 1864, p. 227) in the contents of the stomach of a child poisoned by its application to the scalp.

THERAPEUTICS.—Our knowledge of the value of arsenic in disease rests solely upon clinical observation, which has abundantly established its use in certain very diverse affections. Chief among these is *chronic malarial dyscrasia*. No one would at present think of employing it in acute *remittent fever*, or even in acute *intermittent*, unless under very peculiar circumstances. It is in those cases which have resisted quinia, in which the paroxysms have become irregular, returning at long or irregular intervals, and in which the anæmia and the general nutritive disturbance are even more prominent than the febrile disorder, that arsenic is especially valuable. In these cases it should be administered with sufficient boldness, very generally in conjunction

with iron. Prof. Geo. B. Wood recommends that the first doses should be as large as the system will endure, so as to make a decided impression at once. When the acute-paroxysms are frequent, it is perhaps well to employ this plan, but when it is rather the cachexia than the active disorder that is to be combated, it is preferable to commence with small doses and to increase them until some constitutional symptom is produced.

In ordinary *intermittents*, after the paroxysms have been broken up by quinine it is very well to place the patient upon a preparation of arsenic, iron, and quinine, as a prophylactic against their return.

When, in ordinary *intermittent* fever, for any cause quinia cannot be administered, arsenic may be employed. In these cases, as already intimated, the first doses should be large, so as to make an immediate impression; from five to ten minims of Fowler's solution, properly diluted, may be given every two or three hours until some decided symptom is produced. When the stomach refuses the remedy, it has been recommended by Boudin to give it by the rectum, which he affirms will often bear even a grain of the acid. Not more than a third of this amount should, however, be used as a commencing dose.

In *malarial intermittent neuralgia*, arsenic may be employed as a very useful adjuvant to the antiperiodic alkaloids.

Having had very little experience myself in the use of arsenic in *skin disease*, Dr. Louis A. Duhring, Clinical Lecturer on Diseases of the Skin at the University Hospital, has kindly furnished me with the following:

As is well known, arsenic has long been used and held in high esteem as a remedy in the treatment of cutaneous diseases. It is proper to state, however, that at the present day there exists a great diversity of opinion concerning its actual value as a therapeutic agent against this class of diseases. Certain dermatologists claim to derive marked good from its employment in quite a large number of affections, while others of equal experience are inclined to place but little reliance upon its curative powers. Without entering at all into this discussion, it may, I think, be unhesitatingly said that it is a remedy of real worth and service in certain cutaneous conditions; but the cases must be selected if we would expect satisfactory results.

To say that arsenic is of use in diseases of the skin viewed collectively, is an assertion so vague and meaningless as to be of no practical value whatever. It is indeed a grand error to think that it is of service in all cases. In reality it is only in a comparatively few conditions that we can look for improvement following its administration. Not only is it necessary to specify the disease, but it is even proper to designate the particular stage, if we would employ the remedy successfully.

Arsenic exerts its influence chiefly upon the rete and epidermis. Hence it is found that diseases affecting the more superficial parts of the skin are most amenable to its influence. It possesses little or no effect upon the diseases which have their seat in the deeper structures of the skin. It has no effect upon infiltrations of the corium.

The action of arsenic upon the skin is slow, weeks and months being requisite to produce the desired result. Improvement once obtained, it is expedient to allow the patient to continue its use for some weeks after all symptoms of disease have disappeared.

Arsenic should never be employed in the acute, inflammatory stage of any disease of the skin. It should never be prescribed when there is great heat, burning, intense itching, or rapid cell-change. It not only is of no benefit at this stage, but is positively injurious. It tends to augment the activity of the morbid process. It stimulates the rete into action when rest is most needed. Its administration then, in whatsoever case, should be withheld until the acute symptoms have completely subsided.

Of the many dissimilar diseases in which arsenic has been employed, both with and without reputed success, no mention need here be made. Views upon this subject are so conflicting as to render an analysis of them out of place upon the present occasion. It will suffice to refer to those which it is generally conceded are more or less favorably influenced.

It is unquestionably of service in *psoriasis*. But it is not of benefit in every case, nor should it be prescribed for all forms, or in all stages, of this disease. Where the process is very active, and attended with intense hyperæmia, it only increases the already inflammatory condition. The more active the cell-proliferation, the less probability is there of its being of benefit. On the other hand, the more indolent and sluggish the disease, the greater the chances for a rapid improvement. Let it be withheld until the process has fairly settled in its career.

It is of use in certain varieties of *eczema squamosum*, especially in those cases where the affection has always been of a squamous character; also in cases where the true elementary lesions which tend to distinguish an *eczema* are but ill defined; and, finally, in examples where the disease is unusually superficial, and where there is only slight infiltration of the skin. Certain forms of persistent, localized papular and abortive vesicular *eczema*, as, for example, of the fingers, also often yield readily to arsenic. But, viewing this subject of the value of arsenic in *eczema* in a broad light, it may be stated that as a rule its employment will be found unsatisfactory. Over the great mass of cases it seems to possess no controlling power.

Pemphigus is frequently influenced, and at times permanently relieved, by its use. And here, as in other conditions, the older the process the more likely are we to obtain favorable results. In examples of the rare disease *lichen ruber*, it is employed with great advantage.

As in certain forms of *eczema* it is administered with good result, so likewise in several varieties of *acne* it proves of service. Taken for a considerable time, several months, it exerts a marked influence upon the indolent, papular form of *acne* of the nose and surrounding parts, especially where there is an accompanying *rosacea*.

The two preparations of arsenic which it is advisable to employ are arsenious

acid and liquor potassæ arsenitis. The latter will be found the more desirable form for ordinary use. Arsenious acid is given in pill form, usually combined with black pepper and powdered liquorice, constituting the compound known as the Asiatic pill, which may be prescribed in various strengths, suitable to the case. The following is the formula somewhat modified: Arsenious acid, two grains; black pepper and glycyrrhiza powder, of each thirty-two grains, with a sufficient quantity of mucilage; mix, and divide into thirty-two pills. S.—One to be taken three times a day, directly after meals. The strength of the pill may be increased or diminished according to symptoms.

The liquor potassæ arsenitis is best given combined with a bitter tincture or infusion; it is also well borne combined with the wine of iron. When prescribed in this way there is less likelihood of gastric and intestinal derangement. The mode of ordering the solution pure, and directing so many drops to be taken at each dose, is objectionable for many reasons, which it is unnecessary to refer to. The average dose which will be found to be most suitable to the majority of individuals is three minims. Four, five, and six minims will often be tolerated, and occasionally even larger doses; but take one hundred patients, and it will be found that the greater number cannot take more than three or four minims for any length of time without derangement of the system. Arsenic should always be taken either with the food or directly afterwards.

Nervous diseases.—In certain nervous affections arsenic acts very favorably, in some unknown way. It is especially in *chorea* that it has acquired a deserved reputation. In this affection iron and other tonics are generally indicated, and may be given consentaneously with the arsenic. It is best, however, to administer the latter separately, as the dose must be steadily increased until œdema or other manifestations betray its decided action.

Arsenic has also been recommended in *whooping-cough*, but is at present very rarely, if ever, used. In ordinary *non-malarial neuralgia* it may be tried, and is sometimes serviceable; in simple *gastralgia*, or *gastric neuralgia*, it has been especially recommended.

Arsenic is employed sometimes with advantage in *asthma*, and may be given by the stomach, but is perhaps more generally useful when inhaled. The following formula has been long employed in the Philadelphia Hospital. The prepared paper is rolled into cigarettes, one of which is smoked two or three times a day, until relief is afforded or some giddiness produced:

CHARTÆ ARSENICALIS COMPOSITÆ (*Compound Arsenical Paper*). R.—Belladonnæ fol., gr. xvi; hyoseyami fol., stramonii fol., āā gr. xlviii; extr. opii, gr. iv; tabaci, gr. lxxx; aquæ, Oj; M. ft. sol. et add. potas. nit., gr. clx; potas. arsenit., gr. ccxx. Saturate bibulous paper, and dry for use.

Rheumatic affections.—Arsenic is of value in those forms of chronic rheumatism in which iodide of potassium is commonly employed. It is often advantageous to alternate, administering one of these alteratives for three or four weeks, and then the other for the same length of time. In *rheumatic*

gout, or *rheumatoid arthritis*, it has been highly extolled, but in my experience has furnished no better results than other remedies. It should, however, always be tried in this most obstinate disorder.

TOXICOLOGY.—Sufficient has already been said concerning the symptoms of both acute and chronic arsenical poisoning. No mention has, however, been made of the peculiar local affections produced in the hands of those artisans who work with the preparations of arsenic. Ulcers about the roots of the nails are generally the first trouble in these cases, but after a time eczematous or papular eruptions appear, and even subdermal crysipelatous inflammation is developed. Very commonly to these local symptoms are added, after a time, the usual phenomena of chronic arsenical poisoning.

In the arts, preparations of arsenic are largely used* as pigments; and, excepting the manufacturers of arsenic, it is almost exclusively those who are accidentally exposed to the deleterious influence of these pigments who suffer from chronic arsenical poisoning.

The poisonous colors are all of them some shade of green, are very cheap, and are remarkable for the purity of tone and their permanence under exposure to light, and are, therefore, much used by paper-makers. Scheele's green—the arsenite of copper—contains fifty-five per cent., by weight, of arsenious acid; and Schweinfurt green—the aceto-arsenite—fifty-eight per cent. Paper coated with them has been largely used not only as hangings, but even as wrappings for confectionery and other edibles. Sweetmeats have been colored with them; pasteboard boxes, artificial flowers, tarlatan dresses, shelves in groceries, walls of dwellings, toys of children, and various other articles, have been made the vehicles of death, so that hundreds of cases of poisoning more or less severe have resulted from the use of these pigments, which ought to be banished by the strictest laws.

Possibly, in these cases, the arsenic sometimes finds access to the system in the form of vapor; but more commonly it is the minute dust, which is separated mechanically and diffused through the room. Not rarely the poison has been taken directly into the stomach, especially by children.†

The fatal dose of arsenic varies very much. Dr. W. C. Jackson (*Amer. Jour. of the Med. Sciences*, July, 1858) reports a case of recovery, under the early use of emetics, after an estimated dose of two ounces had been taken; and Dr. E. D. Mackenzie gives an account (*Indian Medical Gazette*, 1872) of a man who swallowed an unknown quantity of arsenic in lumps, and received no treatment for sixteen hours, yet recovered after passing per anum one hundred and five grains of arsenic in two masses.

On the other hand, death has resulted from the use of very small amounts. Dr. Taylor asserts that the smallest fatal dose hitherto recorded is two grains.

* It is said that from five hundred to seven hundred tons were manufactured in 1862 in England alone. (*Mass. Report*, p. 20.)

† For an excellent report upon this subject, see *Third Annual Report of the State Board of Health of Massachusetts*, Boston, Jan. 1872.

Dr. Lachèse (*Ann. d'Hyg. et de Méd. Légale*, 1834, 1e sér., t. xvii.) affirms that six milligrammes (0.09 gr.) will produce decided but not serious symptoms, and that from one to three centigrammes (0.154 to 0.462 gr.) are poisonous, and from five to ten centigrammes (0.77 to 1.54 gr.) fatal. Tardieu places the minimum lethal dose at ten to fifteen centigrammes (1.54 to 2.31 gr.). The escapes from death after the ingestion of large amounts of arsenic have, without doubt, depended upon its being, as in the cases above narrated, in an insoluble form. The effects of the arsenical solutions, such as Fowler's, are more rapid and severe than those of the solid drug.

As arsenic in large doses generally induces vomiting, it is very rarely necessary in poisoning to evacuate the stomach by artificial means. If free emesis, however, have not occurred, a prompt emetic, such as mustard or sulphate of zinc, should be at once exhibited, and very generally the stomach should be well washed out by large draughts of warm water, with salt, if necessary for the return of the water. With the emetic, or sooner, if possible, the antidote should be administered.

The substance whose antidotal value has been most thoroughly tested and assured by clinical experience is the *freshly precipitated sesquioxide of iron*, which forms with arsenious acid a very insoluble compound. The antidote must be freshly prepared, and must be given in great excess: according to the experiments of Messrs. T. and H. Smith, of Edinburgh, at least eight grains of the iron being required for the conversion of one grain of the arsenious acid. In practice, any of the sesqui solutions of iron—that of the chloride being generally preferred, as most readily procured—should be neutralized by the carbonate of soda, and a portion of the precipitate given at once, stirred up in *hot* water. The remainder of the antidote, having been hastily washed by emptying on to a piece of muslin or a filter, pouring water on it and allowing it to drain, should be administered very freely; indeed, indefinitely, as it is entirely harmless.

Recently, H. Köhler, of Halle (*British and Foreign Med.-Chir. Review*, vol. xlv., 1870, p. 538), has made a very elaborate series of chemical, physiological, and clinical experiments upon the comparative antidotal values of the *saccharated oxide* of iron and the freshly-precipitated sesquioxide. His results indicate that the former preparation is the better; but, as the efficiency of the sesquioxide has been so frequently proven at the bedside, further testimony is desirable before it is superseded, especially since the other ferric preparation is not officinal with us, and is not so readily prepared on the spur of the moment as its fellow. *Magnesia*, freshly calcined or freshly precipitated from a solution of its salts, is an antidote of some avail in arsenical poisoning, but is decidedly less efficient than the oxide of iron.

After the emetic has acted in a case of arsenical poisoning, and while the antidote is being given, castor oil should be administered, for the purpose of expelling the poison from the bowels. The further treatment should be directed by general principles: demulcent drinks, opium, stimulants, dry external

heat, and rubbing, being employed as called for by the symptoms. When there is a tendency to suppression of urine, very large draughts of water containing sweet spirits of nitre should be given as frequently as the stomach will bear them.

The chief indications in *chronic arsenical poisoning* are to remove the patient from the exposure and to treat symptoms as they arise. Although I do not know of any clinical records bearing upon the subject, it might be well to exhibit the iodide of potassium, with the hope of hastening the elimination of the poison.

ADMINISTRATION.—The commencing dose of arsenic is one-twentieth of a grain, which should be given in pill *after* meals, and be slowly increased until a perceptible influence, or the desired therapeutic effect, is obtained. The following are the only official preparations:

Liquor Potassii Arsenitis—*Solution of Arsenite of Potassium.*—*Fowler's Solution* contains four grains of arsenious acid to the ounce, is nearly colorless, odorless, with a very faint taste of the compound spirits of lavender, which it contains. It is an excellent preparation, which may be substituted for solid arsenic. The average commencement dose is five drops in a wineglassful of water, to be increased and used with the same precautions as arsenic.

Sodii Arseniatis—*Arseniate of Sodium.*—This salt occurs in transparent, slightly efflorescent, soluble crystals, and is solely used in making the *Liquor Sodii Arseniatis*. The *Solution of Arseniate of Sodium* (grs. lxiv to Oj) may be used instead of *Fowler's Solution*, in similar doses.

Arsenici Iodidi—*Iodide of Arsenic.*—This is an orange-red, crystalline solid, wholly soluble in water and entirely volatilized by heat. It has been used as an alterative, and also as an external application in certain diseases of the skin, especially *lupus* and *chronic tubercular affections*. Iodide of arsenic enters into *Donovan's Solution*.

Liquor Arsenici Chloridi—*Solution of Chloride of Arsenic.*—This preparation is of the same strength, and has the same therapeutic properties, as *Fowler's Solution*. It is, however, a little more irritant than that preparation. The doses are the same.

HYDRARGYRUM—MERCURY. (Hg.)

PHYSIOLOGICAL ACTION.—When a mild, unirritating preparation of mercury is introduced into the system so as to produce constitutional effects, the first symptoms of its action are to be looked for in the mouth. In the mildest degree these symptoms consist of a slight fetor of the breath, and some soreness of the teeth when knocked forcibly together or struck with a key. Mercurial fetor is peculiar, and is generally the first indication that the drug is affecting the system, and is sooner or later accompanied by a disagreeable metallic taste. If the use of the mercury be persisted in, the gums become swollen, soft, and spongy, bleeding on very slight abrasion, and there is a

decided increase in the secretion of saliva. Beyond this point the therapist is never justified in carrying the use of the drug. If it is done, the local symptoms in the mouth increase in severity, the gums become swollen, inflamed, very vascular, and marked by a dark-red line at the junction of the teeth; the tongue is also swollen, sometimes enormously, protruding from the mouth, whose closure it may entirely prevent; the teeth are loosened in their sockets; the saliva is enormously increased in quantity and altered in quality, forming great, ropy, viscid masses, which pour over the thickened lips; the parotid glands, and even the submaxillary, are very much enlarged, and tender. Sometimes, before salivation occurs, slight systemic erethism, marked by a quickened pulse and general restlessness, may be present; but when the mouth-symptoms are severe, very generally there is a distinct febrile reaction of a low type.

In some cases of mercurialization the stomatitis has been very intense: loss of the teeth, extensive ulceration of the soft parts, and even necrosis of the jaw-bones, have occurred, and death from exhaustion resulted, or the patient struggled through to recovery, seamed and disfigured for life. In these cases passive hemorrhages often recur again and again, and, it may be, contribute largely to a fatal result. During severe ptyalism emaciation goes on rapidly, and seems to affect especially imperfectly organized tissues, so that exudations very generally rapidly disappear. The disturbance of nutrition is further shown in some cases by the occurrence of ulcers upon the extremities. The blood suffers very decidedly, becoming more fluid and watery than normal, and having its power of coagulation impaired. According to the researches of Dr. Wright, its solid constituents are notably diminished, including albumen, fibrin, and the red corpuscles, and it contains a large quantity of a fetid, fatty material.

Sometimes the influence of mercury falls almost exclusively upon the nervous system, and produces a peculiar train of paralytic phenomena. This occurs chiefly, if not exclusively, when it, as vapor, finds entrance to the blood through the lungs, and is most frequently seen in those who work in the metal. It is generally the result of long exposure; but that it may be produced in a very short time is proven by the case, related by Dr. Christison, of two barometer-makers who slept one night in a room containing a pot of mercury upon a stove. One was severely salivated, the other was affected with a shaking palsy which lasted all his life. According to Dr. Sigmond (*Mercury, Blue Pill, and Calomel; Their Use and Abuse*, London, 1840), the attack of mercurial palsy, which is sometimes sudden, sometimes gradual, begins with unsteadiness and shaking of the extremities, and of the muscles of the face, which movements interfere with walking, speaking, or chewing; the tremors become frequent, nay, almost constant; "every action is performed by starts." If the exposure be continued, sleeplessness, loss of memory, and death terminate the scene. A peculiar brownish hue of the whole body, and dry skin, generally accompany the disease. In its first attack it may be mis-

taken for St. Vitus's dance; in its latter stages for delirium tremens. According to Noël Guéneau de Mussy (*Gazette des Hôpitaux*, 1868), these two forms are rather distinct varieties than different stages of mercurial tremors. In the latter the affection simulates *paralysis agitans* in its shaking movements; in the former the motions are violent, and occur independently of the will of the patient, even when he is lying quietly in bed.

In other cases, neuralgic pains are a prominent result of mercurial exposure; and sometimes epilepsy is produced, or the intellect is especially affected, and insanity, most frequently of a melancholic type, results. According to Dr. Sigmond, the peculiar paralysis of lead-poisoning, including the drop-wrist, has been known to follow persistent mercurial inunctions.

In some cases, exposure to the vapor of mercury, or even its persistent medicinal use, has resulted in the production of a state of the system somewhat resembling scurvy, characterized by great anæmia, emaciation, and general loss of power, with loss of the hair, aching pains in the bones and joints, œdema, fetid breath, diarrhœa, and, generally, disordered secretions. This is the so-called *mercurial cachexia*.

As already stated, the salivary glands are especially sensitive to the constitutional effects of mercury, and there is some reason for believing that the pancreas, which resembles them in structure, is also obnoxious to the drug. Thus, in a case related by Dr. Copeland, a woman after excessive salivation experienced deep-seated epigastric pain and heat, with nausea, thirst, and fever, and voided thin stools containing liquid resembling salivary fluid. At the post-mortem the pancreas was found weighing four ounces, red, congested, and with its duct dilated. In regard to the action of mercury upon the liver, see article on Calomel as a purgative.

That mercury is absorbed there is abundant proof. That it is eliminated by the secretions is also very evident. Thus, it has been found in the blood,* in the urine,† in the serum of ulcers,‡ in the saliva,§ in the feces,|| in the pus from ulcers, in the seminal fluid,¶ indeed, in every conceivable secretion and in every tissue. Heller, of Vienna,** has found it in the aborted fetuses of salivated women, and Mayençon and Bergeret in the urine of a baby whose nurse was taking calomel.

It is a matter of much interest to know how rapidly mercury is eliminated, and whether, when given internally, it accumulates in the system. The recent researches of Mayençon and Bergeret (*loc. cit.*) throw some light upon this point as well as upon the rapidity of absorption. They found that

* Eld and Buchner, quoted by Prof. Stillé.

† Cantu, Jourda, Andouard, quoted by Prof. Stillé.

‡ Foureroy, quoted by Prof. Stillé.

§ Gmelin, *Bull. de Thérap.*, xiii.; Byanon, quoted by Mayençon and Bergeret; Saikowsky, *Virchow's Archiv*, xxxvii. 347; Oesterlen, quoted by Prof. Stillé.

|| Saikowsky, *loc. cit.*, p. 347.

¶ Mayençon and Bergeret, *Robin's Journal de l'Anatomie*, 1873.

** Quoted by Prof. Stillé.

if one centigramme of corrosive sublimate was given hypodermically to a dog, the urine for the next twenty-four hours contained mercury, afterwards none. When a centigramme was given daily for ten or twelve days, the urine contained mercury for four or five days after the cessation of medication. In their last series of experiments, rabbits received the drug, and were killed at different intervals: in half an hour the metal could be found in all the tissues, the liver and kidneys containing most of it; in four days, or even in a shorter time, mercury given in a single dose was all eliminated, and could not be found in the tissues. This would seem to prove that mercury given in a single dose does not remain in the system; but the experiments of Mayençon and his colleague, confirmed by clinical experience, show that when the drug is administered repeatedly elimination ceases before it is all discharged.

Thus, forty-eight hours after the cessation of a mercurial course, when the urine of one of the investigators was free from the metal, the iodide of potassium was exhibited; and the urine of the next twenty-four hours contained an abundance of mercury, which continued to be present in diminishing quantities for seventy-two hours. The chief channel of escape seems to be the kidneys; but it is very certain that, at least in some cases, the drug is freely excreted by the salivary glands as well as by the intestines.

In regard to the constitutional action of mercury we know but little, except that in some way it influences nutrition.* It certainly increases the secretions, probably of all the glandular organs, and the likelihood is that it does this by a direct action upon the cells. It has been shown by Saikowsky that mercury will cause in the rabbit diabetes, as well as a deposit of the phosphate of lime in the kidneys, and in the dog fatty degeneration of the renal epithelium. These facts, and the emaciation, the perverted functions of nerves and glandular tissue, the various skin-eruptions, the altered blood of chronic hydrargyria, all point to a profound impression, affecting the nutrition of every part of the organism.

The researches of Dr. Hermann von Boeck (*Schmidt's Jahrbücher*, Bd. cxlv. p. 142) would seem to show that mercury causes no especial waste or destruction of the nitrogenous compounds of the body.

The observer mentioned analyzed the feces and urine of a man before, during, and after the exhibition of mercury, taking proper precautions to assure uniformity as to diet and exercise. There was a slight but not a notable increase in the amount of nitrogen in the two excretions during the mercurial period.

THERAPEUTICS.—The use of mercury in affections of the liver and of the alimentary canal are fully discussed in another portion of this work; and, although the drug has been used for almost innumerable purposes in times

* It is well to give a reference here to an article by A. Polotebenow, of St. Petersburg, although the results obtained do not seem to require discussion in a work like the present. It is on the effect of mixing large quantities of corrosive sublimate albuminate with blood outside of the body, and may be found in *Virchow's Archiv*, 1864, Bd. xxi. p. 35.

past, it seems here only necessary to speak of its action as an antiphlogistic and as an antisyphilitic.

Antiphlogistic action.—The use of mercury in inflammation originated towards the close of the last century with a Dr. Robert Hamilton, and soon became universal in England and America. It is a matter of regret that no sufficient analysis of the blood of pyralized persons has been made to determine exactly what are the changes produced in the vital fluid by mercury. The indications are, however, very strong that chief among them is a lessening of the amount of fibrin. As is well known, increase of the hæmic fibrin is one of the most characteristic effects of inflammation: consequently, theory, so far from being opposed to the antiphlogistic use of calomel, affords at least some grounds for the belief that there is more or less of antagonism between the processes of mercurialization and of inflammation.

All important evidence as to the antiphlogistic value of medicines of this class at present available is clinical, and even of this it seems impossible to find much that is very exact and of such nature as to exclude possible fallacies. It is the enormous mass of testimony that overrides the chance of fallacies. It is the general judgment of the profession, founded upon the thousand daily-observed bedside facts, that indorses the use of mercury as an antiphlogistic. In other words, our knowledge of the value of mercurials in inflammation at present is clinical rather than experimental, empirical rather than scientific, but it seems scarcely possible that it is not true.

There is one inflammatory affection, *iritis*, which, from its anatomical relations, is completely visible at all stages; and the effects of the drug upon its processes have been noted from day to day hundreds of times. Oculists are, I believe, agreed that when there is a marked tendency towards the exudation of lymph in this disease, mercury is to be exhibited until pyalism is induced.

Of all inflammations, those of the *serous membranes* seem to be most allied to *iritis*; and it is exactly in the condition above spoken of, where there is a tendency to fibrinous exudations in *pleuritis*, *peritonitis*, and *pericarditis*, that mercury is so constantly employed with so good an effect. In parenchymatous inflammations, especially in *pneumonia* and in *hepatitis*, mercury is used with asserted advantage by many practitioners, but its value is certainly more questionable than in serous inflammations. Calomel is useful in severe *laryngitis*, and especially in the *pseudo-membranous* variety, when the type is sthenic; and no time should be lost in bringing the system under its influence.

The extent of its power to arrest the course of *endocarditis* is certainly an open question; but as it is extremely important, if possible, in this disease, to prevent exudation, and as mercury is the most efficient known agent for effecting this, it should be administered freely and at once. If the disease be of rheumatic origin, the alkalies may be administered conjointly with the mercurial.

In whatever disease a mercurial is administered as an antiphlogistic, it should be given during the stage of exudation, and to facilitate the absorption of the newly-organized lymph after it has ceased to be thrown out.

In the majority of cases, mercury given for its constitutional effects should be combined with opium, to prevent its acting on the bowels.

Calomel should not be used in *adynamic inflammations*, or where the exudation is serous rather than fibrinous. In *puerperal peritonitis* it has been strongly advocated by some, and as strongly condemned by others, simply because there are two varieties of the disease, the *sporadic* or *sthenic*, and the *epidemic* or *asthenic*; and in the one both bleeding and calomel are strongly indicated, whilst in the other they are effective only for evil.

Mercury as an Antisyphilitic.—It was formerly believed that syphilis could not be cured without the use of mercury; but latterly there has arisen a school of syphilographers who assert that the drug is not only not necessary, but is at all stages and in all cases of the disease most injurious; that the worst symptoms of the disease are due not to the constitutional affection, but to the remedy given for its relief.

The great bulk of the profession occupies a middle ground between these extremes, holding the opinion that whilst mercury is not absolutely essential for the relief of syphilis, it is yet in many cases of the utmost value when judiciously used.

The justice of this position cannot, I think, be rightly questioned: the universal verdict in its favor is too fixed and definite; so that the important point now is to determine at what stages, and under what conditions, the remedy is advisable, and what is the best method of its application.

Syphilis is ordinarily, and with sufficient accuracy for practical purposes, divided into three stages, the primary, the secondary, and the tertiary. According to the teachings of the dualists, there are two varieties, or rather species, of venereal ulcer, the hard and the soft chancre, or the true chancre and the chancroid, the infecting and the non-infecting sore. When a venereal ulcer offers the characteristic of the latter of these, mercury should never be exhibited. So long as there is a doubt as to the nature of the primary sore, the remedy should be withheld; but when there is distinct induration, and the inguinal glands begin to be involved, it should be given.

It is proper to state that some high authorities deny the expediency of giving the mercurial until the appearance of distinct secondaries. They affirm that mercury is powerless to prevent the occurrence of these phenomena; that the good which it does is found in its hastening their completion, and consequently that it should be reserved until the second stage of the disorder.

In *tertiary* syphilis, mercury is to be used cautiously, and only in combination with the iodide of potassium.

In the *hereditary* syphilis of infants, a decided mercurial impression offers the best chance of relief.

In the primary or secondary stages of syphilis, whatever form of mercury

be employed, it should be so administered as to exhibit signs of its constitutional action upon the mouth. It is not necessary to pyralize the patient severely, or indeed at all, the proper course consisting in the steady maintenance of the slightest possible distinct soreness of the gums. There are various methods by which this may be done. That most frequently employed, because most convenient, is the administration of small doses of calomel or blue pill by the mouth: from one-quarter to one-half a grain of calomel, or twice as much of the blue mass, combined, if necessary, with opium, to prevent its action upon the bowels, may be given three times a day, and increased if required. Instead of the internal use of the mercurial, the system may be brought under its influence by inunctions with the unguentum hydrargyri. When this is done, the skin should be well cleansed and softened by frequent bathing, and then a drachm of the ointment may be rubbed into the inside of the thighs, legs, and the popliteal spaces, in such a way that the application be not made to any spot oftener than every other day.

It is commonly advised in English works to employ the armpits; and, as the skin is there exceedingly thin and the absorbents very numerous, mercury is without doubt more rapidly taken up at that place than at any other part of the body. When, however, mercury is applied to any hairy surface, it very commonly in a short time induces a troublesome eruption, due to inflammation about the hair-follicles. The eruption appears anywhere on the skin if the mercurial ointment be applied too freely; and, in order to avoid this inconvenience, in Germany the following plan is adopted (Dr. H. Zeissl, *Lehrbuch der Syphilis*, Theil ii. p. 349): The patient, having been prepared by thorough warm bathing, and having received about half a drachm of the ointment, is directed to place it in the hollow of the hand and to rub the two hands together until the ointment is equally diffused; then to apply it forcibly and slowly to the part directed until almost all of the salve has disappeared, having been rubbed into the skin. In most cases the mercurial is applied daily; but in very susceptible persons only every third day. A regular order is maintained in the application, as follows:

1st day. Inner side of both upper arms.

2d day. Inner side of both thighs.

3d day. Inner side of both forearms.

4th day. Inner side of both legs.

5th day. Upon both groins.

6th day. Upon the back.

7th day. Recommence the series.

The advantage claimed for inunction is that the digestion is less apt to be disturbed than when the drug is exhibited by the mouth; the disadvantages are the greater or less publicity which it entails, the trouble which it involves, and its apparent dirtiness. In private practice it is rarely used except in the case of infants, when the mercurial ointment is rubbed into the abdomen and armpits, or often simply smeared upon the flannel roller or binder which

usually envelops the body. The mercurialization of the nurse, with the object of affecting the child, is unjustifiable, unless the nurse and nursing are alike diseased: indeed, to allow a syphilitic child to feed at the breast of a healthy woman is extremely reprehensible.

Mercury is sometimes administered in secondary syphilis in the form of fumigations. The patient is placed upon a chair, and surrounded by a large blanket, or, better, india-rubber cloth, so arranged as to fit tightly around his neck above, and below to encompass the chair. The mercurial preparation is placed upon a metal plate, heated by a spirit-lamp, beneath the chair, and the fumes are allowed to fill the space around the patient inside of the blanket. The heat produced generally causes the patient to sweat profusely, and in from fifteen minutes to half an hour the lamp should be withdrawn, and the patient allowed to cool off, and after a time be put to bed and wrapped up in blankets, with the deposit of mercury still adhering to his skin. The fumigation may be practiced every other night, or at longer intervals, and is believed by some to be especially useful in cases of secondary skin-eruptions. Calomel, black oxide, and cinnabar are the preparations generally used. When the last is employed, care must be exercised that the patient do not breathe the fumes.

In advanced secondaries, the iodides of mercury, given by the mouth, are to be preferred. They should never be pushed to the point of producing salivation.

In tertiary syphilis, mercury should rarely, if ever, be exhibited by itself, but the combination of the corrosive sublimate and the iodide is often more efficient than the latter drug alone. In most cases not more than one-twelfth of a grain of the bichloride should be given, three times a day.

ADMINISTRATION.—The following preparations contain metallic mercury :

Unguentum Hydrargyri.—*Blue, or Mercurial, Ointment* is made by triturating two ounces of mercury with one ounce each of suet and lard, until the metal is extinguished,—i.e., until a portion of the mass rubbed upon a piece of paper exhibits no globules under a magnifying power of four diameters. Mercurial ointment is soft, of a bluish color, becoming darker by age. When frequently rubbed upon the same part, it not rarely produces a disagreeable eruption. It is used to make a constitutional impression, and also locally as a resolvent in cases of enlarged *indurated glands*.

Emplastrum Hydrargyri.—*Mercurial Plaster* contains mercury, olive oil, resin, and lead plaster, and is used as a resolvent in *indurated glands*, enlarged *chronically inflamed joints*, etc.

Pilulæ Hydrargyri.—*Pills of Mercury*.—"Blue Mass."—"Blue Pills" are made by extinguishing mercury with confection of rose and powdered liquorice-root, and so dividing that each pill weighs three grains and contains one grain of mercury. They are used for the same purposes as calomel, but are milder, and, when given to affect the system, less apt to disturb the bowels.

Hydrargyrum cum Creta.—*Mercury with Chalk*.—*Gray Powder* is made by extinguishing three ounces of mercury with five ounces of chalk. It is a

smooth, grayish powder, and is similar in its medical properties to blue mass, but much weaker. It is chiefly used as an alterative cholagogue in *bowel-complaints* of children. The dose for a child is from two to four grains, according to age.

HYDRARGYRI CHLORIDUM MITE—MILD CHLORIDE OF MERCURY
($\text{HgCl} - \text{Hg}_2\text{Cl}_2$).

Calomel is made by boiling mercury and sulphuric acid together in such proportions as to form a bisulphate of the deutoxide of mercury, reducing this to the simple sulphate of the protoxide of mercury by triturating it with more of the metal and subliming with the chloride of sodium. The sublimate is to be well washed with water, to remove any of the bichloride that shall have been formed owing to the imperfect reduction of the bisulphate to the sulphate. When the washings are no longer affected by the addition of ammonia, it may be known that the drug is free from the soluble corrosive sublimate. Calomel is sometimes manufactured by precipitating corrosive sublimate by sulphurous acid; but this method is not officinal, and is subject to serious disadvantages.

PHYSIOLOGICAL ACTION.—Owing to the great insolubility of calomel, a good deal of discussion has occurred as to the way by which it finds entrance into the system. The theory of Mialhe (*Chimie appliquée*), a modification of one originally advanced by Snow (*Lancet*, 1840), has been pretty widely accepted, though with some hesitation, but is, I think, untrue. According to the chemist named, the calomel is converted by the chlorides of the stomach into corrosive sublimate, and as such is absorbed. The action of calomel upon man is so different from that of corrosive sublimate as to render this theory exceedingly improbable, and, at temperatures even higher than that of the stomach, Mialhe was never able to obtain the formation of more than a sixteenth of a grain of the sublimate by the gastric juices. Further, Bucheim, Oettingen, and Winckler (quoted by Prof. Stillé, *Therapeutics*, 2d ed., p. 655) affirm that this conversion does not occur at all at the temperature of the body. Jeannel (*Schmidt's Jahrbücher*, Bd. cxliii. p. 9, from *Journ. de Bordeaux*, 4e sér., ii. p. 67, 1869) has confirmed this, and has suggested what seems to be the way in which calomel is absorbed. He finds that when the protochloride of mercury is placed in a solution of an alkaline carbonate it is decomposed and the gray oxide precipitated. A small portion, however, of the latter is held in solution, as much as 0.02 part in 50 parts of water (by weight); and if a fatty oil be mixed with the alkaline solution a very large part of the mercury is dissolved. From these facts it would seem to follow that the calomel entering the stomach escapes unchanged into the alimentary canal, and is there decomposed by the alkaline juices and dissolved by the fatty matters always present. The physiological evidence would seem to confirm this, for calomel, being absorbed in this way, ought to resemble blue mass rather than corrosive sublimate in its action,—which it does.

The varying constitution of the alimentary juices and the complex chemical relations of calomel would indicate that its solution in the alimentary canal is accomplished in more ways than one,—an indication which is confirmed by the varying results following the ingestion of the drug. It is probable that at times, when the stomach contains more than usual of chlorides and of hydrochloric acid, a very slight portion of the calomel is converted into corrosive sublimate, and that when there is an excess of sulphuretted hydrogen in the alimentary canal a soluble sulphide may be formed. Calomel has been used subcutaneously (Dr. F. Zambon, *Gaz. Med. Ital. Prov. Venete; The Doctor*, 1871), with asserted good results, in syphilis;* and, according to Bellini, its solution is due to the presence of alkaline chlorides in the system. He states that very severe constitutional and local symptoms may follow the hypodermic injection of calomel in persons taking the alkaline iodides, bromides, or sulphates.

The influence of calomel upon the system has been sufficiently discussed. It remains only to state that its freedom from all irritant properties is shown when taken internally or when used externally. Probably no single dose of it is capable, in the average man, of acting as a violent poison, owing to but a limited portion of it being absorbed, and the remainder being swept out of the alimentary canal by the diarrhœa produced. At least it is stated that in the Western United States it is very frequently taken in teaspoonful doses, that sixteen grains of it will act as vigorously as an ounce, and that a pound of it has been given in a case of cholera without visible effect.†

THERAPEUTICS.—Calomel is employed as a mercurial purgative, and also to produce constitutional mercurialization.

ADMINISTRATION.—The dose of calomel as an alterative is usually given as a half to one grain; as a purgative, six to ten grains, followed, six hours after its administration, by Seidlitz powder, or other saline, if required. Minute doses (one-sixth of a grain) of calomel given every hour afford a very good method of impressing the system rapidly. When it is desired to get its constitutional influence, it is generally necessary to conjoin opium with it, to prevent purging.

HYDRARGYRI CHLORIDUM CORROSIVUM—CORROSIVE CHLORIDE OF MERCURY (HgCl_2 — HgCl_2).

Bichloride of Mercury, or *Corrosive Sublimate*, is made by subliming the bisulphate of mercury with common salt. It occurs in the form of colorless crystals, or of white, semi-transparent, crystalline masses, of an acrid, metallic, styptic, and very persistent taste, soluble in sixteen parts of cold and in three of boiling water. It is at once distinguished from the other mercurial prepara-

* See paper by Prof. R. Bellini (*Lo Sperimentale*, June, 1873; *London Medical Record*, 1873).

† Prof. Geo. B. Wood's *Therapeutics*, vol. ii. p. 565.

tions by its color, taste, and solubility, and by its forming a yellow precipitate with lime-water.

PHYSIOLOGICAL ACTION.—Corrosive sublimate is a violent irritant, and in concentrated form caustic. When given in small, repeated doses, although capable of inducing salivation, it is less apt to do so than is calomel or blue pill. In overdoses it produces symptoms of irritant poisoning of a severity proportionate to the dose. If the latter be small, the manifestations may be only some nausea, slight burning in the stomach, colicky pains in the abdomen, and diarrhœa. After large doses these symptoms are intensified. The subject first experiences a peculiar metallic, coppery taste at or shortly after swallowing the poison. If the solution be concentrated, deglutition is interfered with by a spasm of the muscles of the throat and larynx, causing a feeling of suffocation, and sometimes even the rejection of the draught. Then burning pains are experienced in the œsophagus and stomach, followed by violent vomiting, at first mucous, then bilious, and finally bloody, and by severe abdominal pain and tenderness, with profuse purging, at first serous in character, but afterwards affording only small, mucous, bloody stools, which are often voided with much straining. The breath generally becomes fetid and offensive in a very short time. In the course of two or three hours, very rarely in less than one hour, collapse occurs, with small, frequent, irregular pulse, pinched, anxious face, cold extremities, and finally death, preceded, it may be, by fainting, convulsions, and coma. The urine is very much lessened in quantity, is sometimes albuminous, or even bloody, and not rarely is suppressed. If the patient survive several days, a petechial eruption may appear, and salivation sometimes, but not always, occurs. In some cases, after the collapse there is an attempt at a febrile reaction, which soon, however, gives place to a second and fatal prostration.

When recovery occurs after severe poisoning, the convalescence is slow and protracted.

In regard to chronic poisoning with corrosive sublimate, sufficient has been said under the general heading, except that colicky pains and abdominal disturbance are more apt to occur with this than with the less irritating preparations of the metal.

Severe purging, and even fatal poisoning, may result from a single external application of this preparation of mercury;* and in animals killed by hypodermic injections of it (see experiments of Dr. J. Rosenbach, *Schmidt's Jahrbücher*, Bd. cxliii. p. 9) diarrhœa and other indications of gastro-intestinal irritation are prominent symptoms.—facts which indicate that the bichloride is eliminated unchanged from the alimentary canal.

* See case reported by Dr. Meeres, *London Lancet*, Sept. 16, 1871, in which a solution (grs. ii to f5i) was applied with a camel's-hair brush to the head of a child nine years old for the cure of tinea tonsurans. The symptoms were diarrhœa, profuse salivation, and great prostration, ending in death.

HYDRARGYRI IODIDUM VIRIDE—GREEN IODIDE OF MERCURY ($\text{HgI}—\text{Hg}_2\text{I}_2$).—This *Protiodide of Mercury* is made by the direct action of iodine upon the metal. It is a greenish-yellow, odorless, and tasteless powder, insoluble in water, ether, and alcohol. The protiodide, when compared with the biniodide or the bichloride, is a mild preparation, and has been used to produce constitutional impression in *syphilis*, especially when of long standing. The iodide of potassium converts it into the biniodide and metallic mercury (U. S. Dispensatory), and should, therefore, never be given in combination with it.

The alterative dose is one-fourth of a grain three times a day, increased to a grain if necessary.

HYDRARGYRI IODIDUM RUBRUM—RED IODIDE OF MERCURY (HgI_2).—The *Biniodide of Mercury* is made by precipitating the bichloride of the metal with iodide of potassium. It is a scarlet-red powder, insoluble in water, but sparingly soluble in alcohol. It is a powerful local irritant, producing, when taken in overdoses, symptoms and results very similar to those caused by corrosive sublimate. It is used a good deal in *tertiary syphilis* and in *syphilitic rheumatism*; also to some extent as a local application in *lupus* (see *Annuaire de Thérap.*, 1852). It is much more active than the protiodide, and should be used as cautiously and in the same doses as corrosive sublimate.

Besides the above preparations of mercury, the U. S. Pharmacopœia recognizes the following:

The deutoxide in two forms, the *Yellow* and the *Red Oxide* (HYDRARGYRI OXIDUM FLAVUM, HYDRARGYRI OXIDUM RUBRUM). Both are used upon *ulcers*, *chancres*, etc., solely for their local effects, and are stimulant and alterative when diluted, mildly escharotic when in powder. The red *Precipitate Ointment* (*Unguentum Hydrargyri Oxidi Rubri*, $\S i$ to $\S vii$), as well as its colleague (*Unguentum Hydrargyri Oxidi Flavi*, $\S i$ to $\S vii$), very generally requires dilution with lard, and is much used in chronic *skin-affections*, in obstinate *conjunctivitis*, in *psorophthalmia*, etc.

The *Red* or *Bi-sulphuret* (HYDRARGYRI SULPHURETUM RUBRUM) was formerly used a good deal in fumigations, but is now rarely employed. *Turpeth Mineral*, or *Yellow Sulphate of Mercury* (HYDRARGYRI SULPHAS FLAVUS), a lemon-yellow powder, sparingly soluble in water, is a basic sesquisulphate of the deutoxide of mercury, prepared by throwing the bisulphate into water, which causes it to break up into a supersulphate, which remains in solution, and the salt in question, which precipitates. Turpeth mineral has been used as a harsh emetic, and also as an alterative, but is now never employed in the former capacity, and very rarely in the latter. The alterative dose is from a quarter to half a grain. Forty grains of it (*Guy's Hospital Reports*, vol. x., 3d series) have caused death; profuse salivation came on in six hours.

White Precipitate, or *Ammoniated Mercury* (HYDRARGYRI AMMONIATUM), is a white complex powder, made by precipitating the bichloride with

water of ammonia. It is used in the form of ointment (*Unguentum Hydrargyri Ammoniaci*, grs. xl to ʒi) as a local application in various skin-affections.

Black Wash and *Yellow Wash*, two non-official but favorite preparations, are respectively made by the addition of a drachm of calomel to a pint of lime-water, and of half a drachm of corrosive sublimate to a pint of lime-water. They depend for their virtues upon the black and yellow oxides of mercury, and are used exclusively as local applications to *chancre*s and other *syphilitic ulcers*. The yellow wash is much the more stimulating of the two.

IODINIUM—IODINE. (I.)

Iodine is a soft, friable, opaque substance, occurring in crystalline scales with a semi-metallic lustre and of a bluish-black color. Its odor resembles that of chlorine; its taste is hot and acrid. It is somewhat volatile at ordinary temperatures, but when heated to 225° F. melts and emits the beautiful purple or violet vapor to which it owes its name. It is freely soluble in glycerine, alcohol, and ether, but requires seven thousand times its weight of water to dissolve it. With starch it strikes a deep-blue color, and this test is so delicate that it will indicate the presence of iodine in four hundred and fifty thousand times its weight of water. In testing animal liquids, such as urine, for iodine, a small quantity of nitric acid should be added to insure its being free in the liquid.

PHYSIOLOGICAL ACTION.—Iodine, when applied to any part of the body, acts as a very powerful irritant, or, if in highly concentrated form, as a mild caustic. The tincture stains the skin yellow, and causes, if applied with sufficient freedom, smarting, some erythematous inflammation, and finally desquamation. Its repeated application blisters and destroys the cuticle. Upon mucous membranes its action is more intense than upon the skin.

When taken internally, a single moderate dose of iodine causes merely some gastric uneasiness and a disagreeable metallic taste in the mouth; when larger amounts are ingested, the gastric uneasiness may be intensified into violent vomiting, with increased salivary flow, abdominal pains, and even purging. In sufficient quantity it is a poison; although very few deaths have been recorded as caused by it. The usual symptoms, when toxic doses of it have been taken into the stomach, are vomiting, purging, severe abdominal pain, headache, giddiness, sometimes violent excitement and convulsions, ending in faintness, pallor, and collapse.

Dr. E. Rose (*Nothnagel's Arzneimittellehre*, Berlin, 1870, 252) has reported a case in which death resulted from a large injection into an ovarian cyst. Very soon after it was given, there ensued severe thirst, with great dryness of the throat and mouth, and then painless vomiting of watery matters containing iodine. The whole surface became very pale, the extremities cyanosed; the radial pulse very frequent, but so small that it could not be counted; the urine very scanty, dark brown, and rich in iodine. After a time, reaction occurred. For three days the vomiting persisted, the pulse

was very frequent, full and hard, and the cheek put on the glow of high fever, but the temperature did not rise above 37.18° C. On the fourth day, exanthematous blotches, not disappearing on pressure, appeared on the skin and in the mouth; the sputa became bloody; and menstruation occurred, two and a half weeks too soon. The urine remained scanty, and on the eighth day, when all other symptoms save swelling of the parotids had disappeared, still contained iodine, and was albuminous. On the tenth day, in the midst of apparent convalescence, the patient died suddenly.

Very large quantities of iodine are asserted to have been taken without serious results. Julia de Fontenelle (quoted by Stillé, *Therapeutics*, ii. p. 731) tells of a man who took two and a half drachms of iodine without experiencing any remarkable effects; and Magendie relates the case of a child four years old who swallowed ten grains without serious consequences.

If full doses of iodine be exhibited continuously for a length of time, a train of phenomena result, known as *Iodism*. In regard to these there has been a good deal of difference of opinion and statement, a difference which seems explainable only upon the supposition that different individuals are differently affected by the drug. Rilliet (Trousseau's report on his memoir, *Bull. de l'Acad. Roy.*, xxv.), who has had wide opportunities and has apparently studied the subject very closely, makes three forms of iodic intoxication: first, that in which the symptoms are those of gastric irritation; second, that characterized by nervous troubles, neuralgia, ringing in the ears, convulsive movements, disturbed intellection, with coryza, ophthalmia, salivation, vomiting, diarrhoea, polyuria, and cutaneous eruptions, and in some cases atrophy of the mammae in the female and of the testicles in the male; third, iodic cachexia, caused either by iodine or iodide of potassium continuously used for many months. It is said to be most easily induced in goitrous persons, and is characterized by rapid emaciation, commencing mostly in the face, and severe nervous palpitations of the heart, with excessive appetite, which sometimes precedes and sometimes follows the loss of flesh. So long as the drug continues to be taken, these symptoms continue to progress, and after a time hysteria or hypochondriasis, with insomnia, manifests itself. The goitre, the mammae, and the testicles waste away together; but if the medicine be suspended and health gradually return, whilst the abnormal growth reappears, the sexual glands remain wasted.

In the experiments of Jörg and his pupils, doses of iodine of a grain to a grain and a half gave rise to colicky pains, increased appetite, watery stools, an increased secretion of urine, malaise, and some headache. When the dose was augmented to two grains, a diffused sense of heat and sexual excitement were superadded. Other observers have noted this abnormal sexual excitement, and some have stated that at times it precedes atrophy of the mammae or of the testicles. Prof. Stillé affirms that the menstrual flow may become excessive, or that during pregnancy abortion may be caused.

The second form of iodism of Rilliet, in which the nervous symptoms are

prominent, has been spoken of by other authorities; and Brodie has especially noted disturbances of vision, and paralysis.

Most authorities affirm that iodine and iodide of potassium produce similar symptoms. I cannot help believing that their therapeutic value is different, and consequently their physiological action. I have given the salt in enormous doses, and have seen nervous symptoms in only a single case,—a man who received for a long time two hundred and seventy grains a day, and who was intensely sleepy and stupid, presenting symptoms exactly similar to those of bromism, including an eruption of acne.

Of the physiological action of iodine, all that we know is that it modifies nutrition,—how or in what way we cannot say. It certainly is absorbed and as certainly is eliminated chiefly by the kidneys, but probably to a greater or less extent by all the mucous membranes; and Dr. R. W. Taylor (*American Journal of Syphilography and Dermatology*, April, 1873) believes that he obtained in a case evidences of the free escape of the iodine through the skin. Dr. C. Handfield Jones (*Beale's Archives*, i.) analyzed the urine of six patients taking large doses of iodide of potassium, with the following results: first, water increased in three cases very much, in one slightly so, in two diminished; second, acidity increased in three and diminished in two; third, urea increased in three and diminished in three; fourth, phosphoric acid and sulphuric acid increased in four and diminished in two; fifth, chlorine increased very greatly in two cases, moderately in one, and decreased in two; sixth, uric acid increased very greatly in two cases and diminished in four.

According to the experiments of Dr. Hermann von Boeck (*Zeitschrift für Biologie*, iii. 393, 1869; *Schmidt's Jahrbücher*, Bd. cxlv. p. 142), the ingestion of iodine does not increase notably the elimination by the kidneys or bowels; but it is said that Rabuteau's* studies yielded a different result.

THERAPEUTICS.—As an alterative, iodine is of especial value in *chronic scrofula*. In those cases in which there is indolent enlargement of the lymphatics, which exhibit no tendency, or but little tendency, to suppurate, it is of especial value. Except in very acute cases, however, it should always be tried, even when the glands do tend towards suppuration, especially as it exerts a very beneficial influence upon the ulcers left after suppuration. In other forms of scrofulous disease, in *chronic enlargements* of the joints, and *bone-affections* of such nature, iodine is often of great service. As scrofulosis is generally, if not always, associated with lowered nutrition and with anæmia, in the great majority of cases, cod-liver oil and iron in some form should be administered as adjuvants to the iodine. At the same time that the drug is exhibited internally in these cases, its ointment should be freely applied to the enlarged and indurated glands.

Experience has demonstrated the value of iodine in goitre, whether of the ordinary variety or of that known as *exophthalmic goitre* or *Graves's disease*.

* I have never seen the original paper, which is, I believe, in the *Gazette Hebdomadaire*, ix., 1869.

All tumors of the thyroid body are not goitre, however; cystic degeneration of it is very common, and is in no wise benefited by iodine. It is in simple hypertrophy of the gland that iodine used internally and applied externally over the tumor is so beneficial. During the acute stage of enlargement the use of leeches is often of great benefit, and whenever much tenderness exists should precede the exhibition of the drug.

In *phthisis*, iodine sometimes does good, but only in the most chronic cases; and inhalations of its vapors, as have been recommended by Piorry, can only be of service by stimulating the bronchial mucous membrane and the surfaces of cavities. When softening is progressing and the lung breaking down, iodine sometimes appears to hasten the process.

Local Application.—As a simple counter-irritant, iodine is very frequently employed when it is desired to maintain a mild, persistent influence, as in *chronic rheumatic affections* and sometimes in *phthisis*. For this purpose the tincture is generally preferred, and it should be applied freely once or twice a day, or every other day, according to the susceptibility of the patient's skin.

In various affections of the skin, iodine has been employed with asserted advantage. In *erysipelas* of the skin, very beneficial results have been ascribed to its local use, and, I think, with justice; but great care is necessary lest it be applied too strong. I have seen very serious results from the destruction by it of the skin in this affection. If the full strength of the tincture be used, it should be applied at first very lightly, and not more than once in the twenty-four hours.

In *psoriasis*, in *acne*, in *parasitic skin-diseases*, it has been used, but holds only a second rank among remedies.

In a similar manner it is employed in various chronic diseases of the mucous membranes, such as *ozana*, *leucorrhœa*, *chronic cystitis*, *chronic dysentery*, and *serofulous ophthalmia*,—whenever, in a word, an alterative, stimulant action is desired.

In cases of retraction of the gums, with consequent loosening of the teeth, Prof. Stillé recommends the application, with a camel's-hair brush, after each meal, of a watery solution (gr. i to fʒi) of iodine, the mouth being immediately afterwards washed.

The most important external use of iodine is as a resolvent in cases of indolent *glandular* hypertrophic enlargement, and where there are large watery exudations, as in some forms of *chronic pleurisy* and of *diseased joints*.

Iodine has been very largely employed by injection into serous cysts, as in *hydrocele*, for the purpose of exciting inflammation and causing obliteration of their cavity; but this use of it is purely surgical, and the reader is referred to treatises upon such subjects.

In chronic *empyema*, the injection of iodine after free exit has been formed for the pus is often of the greatest service. The solution in the beginning should be very weak, containing not more than six grains each of iodine and

of iodide of potassium in a pint of water; with this the pleura should be daily washed out, the strength of the solution being gradually increased.

ADMINISTRATION.—Iodine is never administered in solid form; nor should the tincture be given internally, because the iodine is precipitated by the watery juices of the stomach.

Taking advantage of the solubility of iodine in a watery solution of iodide of potassium, the framers of the Pharmacopœia have directed the following preparations, which may be used internally; they should always be given well diluted, after meals:

Liquor Iodini Compositus—Compound Solution of Iodine—*Lugol's Solution* (Iodine, $\mathfrak{v}$ i; Iodide of Potassium, $\mathfrak{z}$ iss; Water, Oj).—Dose, gtt. v to xv.

Tinctura Iodini Composita—Compound Tincture of Iodine (Iodine, $\mathfrak{z}$ ss; Iodide of Potassium, $\mathfrak{z}$ i; Alcohol, Oj).—Dose, gtt. x to xx.

For external use, the following preparations are official:

Tinctura Iodini—Tincture of Iodine ($\mathfrak{z}$ i to Oj).

Unguentum Iodini—Iodine Ointment (Iodine, gr. xx; Iodide of Potassium, gr. iv; to $\mathfrak{z}$ i).

Unguentum Iodini Compositum—Compound Iodine Ointment (Iodine, gr. xv; Iodide of Potassium, gr. xxx; to $\mathfrak{z}$ i).

POTASSII IODIDUM—IODIDE OF POTASSIUM (KI).

This salt occurs in white or colorless, generally cubic, crystals, soluble in two-thirds of their weight of water and in from six to eight parts of rectified spirits. If to its solution starch be added, no blue color should arise, but on the passage of chlorine the characteristic iodine reaction should take place, owing to the liberation of the metalloid by the gas; or if sulphuric acid be added, a purple tint gradually appears, and deepens into blue; a spontaneous blue color betrays the presence of the iodate of potassium, a harmful adulteration. At a dull red heat iodide of potassium fuses into a crystalline mass; by a bright heat it is decomposed.

PHYSIOLOGICAL ACTION.—Iodide of potassium influences nutrition in a manner similar to iodine; indeed, most authorities teach that their action is identical; yet in therapeutics they find a different range of employment, and, I believe, act differently.

THERAPEUTICS.—In certain forms of *rheumatism*, iodide of potassium is of the utmost value. In the early, active stages of *inflammatory rheumatism*, it is much less efficient than the alkalies; but when all the good that can be has been obtained from these, and when the joint symptoms persist in a sub-acute form, the iodide comes very well into play.

In *subacute* or *muscular rheumatism*, the iodide is the most efficient remedy. Often when the symptoms are very acute it may be advantageously combined with the alkalies, and in lingering cases, especially where there is reason to suspect a gouty taint, with colchicum. In *sciatica*, in *lumbago*, in *rheumatic*

neuralgia following exposure to cold or wet, as in all other forms of subacute rheumatism, very much is to be hoped for from its use.

In *gout*, it is of less service than in rheumatism, but in the chronic form of the disease, and in the irregular, inherited gout which so frequently appears as neuralgia or other anomalous affection, it adds to the efficiency of small continuous doses of colchicum.

In *rheumatic gout*, or *rheumatoid arthritis*, it should be tried; although little is to be hoped for from its use.

In *tertiary syphilis*, including in the term all cases of syphilitic bone, visceral, or nervous disease, the remedy is really of inestimable value. It must be given freely, and, when there is no cachexia, may be advantageously combined with the bichloride of mercury. It is scarcely in place here to enumerate all the forms which tertiary syphilis may assume; but the iodide is useful wherever the dyscrasia has existed for a length of time.

The iodide of potassium appears to have the power of promoting absorption of serous fluids, and certainly is of value in *chronic pleuritis* with effusion, in *chronic pericarditis*, and even in *chronic hydrocephalus*.

In various chronic *metallic poisonings*, the iodide of potassium is of great service. With both lead and mercury it forms double salts, which are soluble, and there is very good reason for believing that the formation of these salts takes place in the economy, and that the metal which has been lying in an insoluble condition in the various tissues is taken up and excreted. Severe salivation and ulcerative stomatitis have sometimes resulted from the use of the potassium salt in those who had previously taken large quantities of mercury;* and in Melsen's experiments, dogs to which insoluble preparations of mercury had previously been given without the induction of severe symptoms afterwards died under the action of the iodide, the mercury also having appeared in their urine. The experiments of Mayençon and Bergeret, quoted in the article on Mercury, afford striking confirmation of these facts, and seem to render the evidence irresistible that the iodide does cause the elimination of mercury.

In regard to lead, the researches of Drs. Parkes, Goolden, Swift, Malherbe, and Sieveking† have shown that very frequently, in cases of chronic lead-poisoning, the exhibition of iodide of potassium causes the appearance of lead in the urine.

This chemical evidence is abundantly corroborated by clinical experience, so that in all cases of chronic metallic poisoning, especially by lead and mercury, the persistent use of the iodide of potassium should be tried.

ADMINISTRATION.—The ordinary dose is ten grains three times a day; but much larger quantities may often be given with impunity, and, in internal syphilitic affections, may be necessary. In the latter class of diseases, the

* See Dr. Budd, *British and Foreign Medico-Chirurgical Review*, xi. 202, for a striking case.

† See Stillé's *Therapeutics*, vol. ii. p. 735, Blanchard & Lea, 1864.

best plan is to begin with twenty grains three times a day, and rapidly to increase the amount until drachm doses are reached, or frontal pain or other symptom of iodism produced. The best substance for disguising the very disagreeable taste of the drug is the compound syrup of sarsaparilla.

LIQUOR ARSENICI ET HYDRARGYRI IODIDI. — *Solution of the Iodides of Arsenic and Mercury* is made by dissolving seventy grains each of the iodide of arsenic and the red iodide of mercury in a pint of water. It was originally suggested by a surgeon of Dublin, by whose name it is very generally known. *Donovan's Solution* is a powerful alterative, used chiefly in very obstinate chronic scaly *skin-diseases* when the local action is of a very low grade, and in *chronic rheumatism*. According to the U. S. Dispensatory, twenty drops of it contain the twenty-fourth of a grain of arsenious acid, a little over the twelfth of a grain of the deutoxide of mercury, and about a quarter of a grain of iodine. It is therefore an exceedingly active preparation, very capable of acting as a corrosive poison, and when administered a little too freely is said sometimes to cause salivation. When applied locally, it acts as a violent irritant. The dose of it is from three to eighteen drops, well diluted.

IODOFORMUM—IODOFORM.

This substance was discovered by Sérullas in 1822, and was introduced as a remedy by Dr. Glover in 1837, but has not become officinal, and has only very recently attracted much attention. It occurs as small, pearly-yellow crystals, having a strong, persistent, saffron-like odor, insoluble in water, but readily soluble in alcohol and ether.

PHYSIOLOGICAL ACTION.—The action of iodoform upon the system has not been sufficiently investigated to render any positive conclusions possible. According to M. Maître, when taken by man in doses of five or six grains it causes no notable symptoms, but two hours after the drug has been ingested, iodine can be found in the urine. In animals, large non-toxic doses of it are said to cause symptoms of intoxication, tottering, weakness, loss of appetite, but no vomiting; and lethal doses to produce violent opisthotonos, convulsion, hurried breathing, and finally death.

When iodoform is applied in strong solution, or in substance, it acts as a very powerful local anæsthetic. Thus, a suppository containing it, if introduced into the rectum, will so benumb the parts that defecation may take place without the person or animal being aware of it.

THERAPEUTICS.—Iodoform has been used internally as an alterative and analgesic in *syphilitic rheumatism* and *night-pains*, and in other forms of *neuralgia*. Although very highly commended by some (Dr. Stiles Kennedy, *Medical and Surgical Reporter*, Jan. 1870, p. 50), it has not come into general use, and, in the extensive trials made with it at the Philadelphia Hospital, has failed to sustain its first reputation in these disorders.

Whatever position iodoform may finally acquire as an internal remedy, there can be no question as to its value when used locally. It is useful in

cases of painful *ulcers*, even when they are *cancerous*,* serving to alleviate pain and to promote cicatrization. The testimony is especially strong as to the value of the remedy in true and soft *chancre*, and, indeed, in all *syphilitic ulcers*; and Dr. A. A. Izard (*New Treatment of Venereal Diseases*, translated by Dr. Howard F. Damon, Boston, 1872), who has had a large experience, even recommends it very highly as a resolvent application to simple, non-virulent *boobes*, but affirms that in phagedenic chancre it is of no value. He further states that its local application should never interfere with or be allowed to replace constitutional treatment. In the Philadelphia Hospital it has become a standard application to *indolent leg-ulcers*, and is thought to act not only as a local anæsthetic, but also as a decided stimulant to nutrition.

ADMINISTRATION.—Iodoform may be applied to ulcers in powder, in solution, or in ointment. When there is a great deal of pain, especially if there be much discharge, the powder may be preferred. In *uterine cancer*, in painful *hemorrhoids*, cacao butter suppositories, containing from five to ten grains of the drug, should be employed. Owing to the bad odor of the drug, its application about the mouth and throat is often objected to. According to Dr. Louis Elsberg (*Philada. Medical Times*, Oct. 4, 1873, vol. iv. p. 4), if to four parts of absolute ether (Squibb's) one part of crystallized iodoform be added, and the whole shaken in a *red glass flask*, a solution is obtained of sufficient strength for effectual use in diseases of the mouth, and free from odor other than that of ether.

OLEUM MORRHUÆ—COD-LIVER OIL.

Cod-liver oil is obtained from the liver of *Gadus morrhua* and other species of *Gadus*. In the manufacture of the so-called *shore oil*, the only variety usually employed in medicine, the fish caught near land are brought at once to the shore, and the oil is obtained from the fresh livers by one or other of several processes. The original custom was to put the livers into large kettles, add water, boil to a pulaceous mass, drain off the liquid, allow it to stand, and finally to skim the oil as it rose to the top. A more modern method is to heat the livers by steam applied to the outside of the vessel containing them, to allow drainage, and to proceed as in the process just described. I am informed that at present some of the finest brands of oil are prepared by forcing currents of steam at high pressure through the mass of livers, tearing them in this way to pieces and melting out their oil.

Shore oil should be a perfectly limpid, yellow, thick oil, free from rancidity, and having the peculiar taste and smell of the oil well developed.

Straits oil or *Banks oil* is prepared from those fish caught at the "Banks," far from land.

* Consult papers by Dr. G. Völker (*Bulletin de Thérapeutique*, t. lxxiii., Dec. 1867) and Dr. Féréol (*Ibid.*, t. lxxiv., May, 1868).

The livers are thrown into casks and allowed to stand for a greater or less length of time and to undergo more or less complete putrefaction, until, on the return of the fishing-smack to port, they are thrown into water-boilers and treated in a manner similar to that previously described.

Of this oil there are two varieties: the *brown* oil, which is much darker than shore oil, and much more nauseous to the taste and smell; and the *black* oil, which is very dark, and still more disgusting in its evidences of rancidity. Both of these varieties are largely used in the preparation of leather.

When a mineral acid (especially the sulphuric) is added to cod-liver oil, the well-known biliary play of colors occurs; but this does not prove the genuineness of the drug, or demonstrate that it is derived from the codfish: it only shows that it is a *liver* oil. It is scarcely to be doubted that not rarely the livers of other fish are largely mixed with those of the *Gadus morrhua*, but it is not probable that this often happens to such an extent as to interfere with the therapeutic value of the product: indeed, it is far from certain that the liver of the cod yields an oil having superior medicinal qualities to that produced by the same organ of other fishes. Be this as it may, the physical properties afford the only known test as to the genuineness of the drug.

Cod-liver oil is a very complex substance, containing, according to the analysis of De Jongh, glycerine, oleic, margaric, butyric, and acetic acids, gaduin, various biliary principles, such as fellinic, cholic, and billifellinic acids, iodine, chlorine, traces of bromine, phosphorus, phosphoric acids, and various other substances. According to the U. S. Dispensatory, the proportion of iodine never exceeds one part in two thousand. In De Jongh's analysis the greatest amount was found in the light-colored oils, and was only four-hundredths of a grain in one hundred grains of the oil. *Gaduin* is a peculiar, dark-brown substance, which is probably medicinally inert. When to cod-liver oil ammonia is added, there can be obtained, by distillation, a peculiar volatile alkaloid, *propylamia*, which exists in no other official oil, but occurs in the ergot.

PHYSIOLOGICAL ACTION.—As is well known, all fatty substances when taken into the system have a tendency to cause deposition or formation of fat in the body. Cod-liver oil certainly shares this property in an eminent degree. Dr. Pollock, as quoted by Prof. Stillé, has found that if there be given of it to pigs from one to two ounces *per diem*, to sheep one ounce, and to bullocks from three to nine ounces, it is digested, and aids in fattening the animal; larger amounts than those noted, in Dr. Pollock's experiments, always deranged very seriously the digestive function.

No close studies of the effect of cod-liver oil upon healthy men have, that I am aware of, been made. Undoubtedly it tends to produce obesity; but, as no other oil is able to supply its place in various chronic diseases, it must have some influence upon nutrition not shared by ordinary fatty matters, and therefore is an *alterative*.

The history of the clinical use of oleum morrhuæ certainly indicates that it influences the constitution of the blood. It is an every-day occurrence to see pale, anæmic patients become, whilst taking it, rosy and plethoric. According to the analysis of the blood of a patient made by Simon, there is, during its use in phthisis, a great increase in the amount of solids in the blood, a diminution of the fibrin, and an increase in the albumen. The examinations of Dugald Campbell (*British and Foreign Med.-Chir. Review*, 1856, vol. xvii. p. 21) have confirmed the results of Simon. It is very probable that cod-liver oil has some peculiar influence upon the blood-making organs.

Upon the various single functions of the body, except the digestive, cod-liver oil has no apparent immediate effect, disturbing directly neither the nervous, motor, respiratory, circulatory, nor secretory movements. When by its use the general nutrition is improved, all the functions seem to share equally in the improvement.

Cod-liver oil has undoubtedly, when given with sufficient freedom, a tendency to cause indigestion and looseness of the bowels. All oils are of difficult digestion, and when too much of the oleum morrhuæ is exhibited in man, as in animals, it exerts a deleterious local effect upon the alimentary apparatus.

Much speculation has been indulged in as to which of its ingredients cod-liver oil owes its peculiar medicinal properties. Certainly, however, no real light has been shed upon this matter, and the present probabilities are that it acts as a whole,—i.e., that its virtues depend upon the peculiar combination.

Almost the only effort which has been made with any degree of success to explain the superiority of cod-liver oil over other animal fats is that of Dr. Oswald Naumann (*Archiv der Heilkunde*, sechster Jahrgang, 1865, p. 536). If the experiments of this investigator be as accurate as they appear, he has certainly proven that cod-liver oil has physical properties which must aid in its usefulness, although it is not probable that its value depends solely, or even in great part, upon them. He first tested the rate at which various oils pass through fresh moist animal membranes when pressed upon by a column of mercury or by the weight of the atmosphere over an exhausted receiver, and found that cod-liver oil passed much more rapidly than any of a number of oils tried. Apparently this power depended in some measure upon the presence of the biliary principles, since if it was deprived of them the rate of its passage was greatly lessened, but was again increased by the addition of a little bile. The investigator then, opening the abdomen of cats, separated in each animal by ligatures two knuckles, of equal length and entirely similar, from the remainder of the intestines. Into each of them he injected a certain amount of bile, and then into one ordinary oil, into the other cod-liver oil; and when the animals died, some hours afterwards, it was always found that much more of the oleum morrhuæ was absorbed than of the other oil.

These experiments were, unfortunately, too few and incomplete to be decisive, but certainly they indicate that the oleum morrhuæ is more easily and rapidly

absorbed than other animal oils. Dr. Naumann's last series of experiments were directed to discovering the comparative ease with which animal oils and the cod-liver oil were oxidized. For this purpose he used a test solution of the permanganate of potassium, and on adding to given bulks of this, in test-tubes, equal amounts of the various oils, noted the changes of color induced by the reduction of the permanganate. He found that cod-liver oil was the first to be affected. It is evident that to be easily absorbed and easily oxidized fit a fat for use in the animal economy; but surely the peculiar value of *oleum morrhuæ* does not depend solely upon these properties.

THERAPEUTICS.—Cod-liver oil is especially useful in that condition of system in which, with general lowered tone, there is a tendency to cellular hyperplasia, to the formation of "exudations" composed of imperfectly-developed cells, which, in the great majority of cases, from the very beginning are incapable of development into perfect entities, having only one potential quality, that of dying. There are various types of this diathesis, or condition of system. In one of them there is a tendency to increase in the lymphatic glands; to multiplication, at the expense of development, of their cellular elements,—*i.e.*, to the formation of numerous imperfectly-developed cells, and, finally, to the destruction of them. The death of these cells is partly due to their inherent qualities, and is partly the result of the pressure which they exert upon one another and upon their sources of food-supply. If they undergo a slow, fatty degeneration, with desiccation, cheesy deposits are formed; if a rapid, fatty change, with abundance of moisture, pus and abscesses are produced: in either case, ulceration is the final result. This is the so-called *scrofulous* diathesis,—*scrofulosis*. In another of this group of diatheses, the tendency to cellular hyperplasia affects the mucous membranes of the air-passages, and the patient, on the slightest provocation, suffers from catarrh, until finally a multiplication of cells occurs so rapidly as to fill up a greater or less number of the air-vesicles of the lungs, generally those of the apex, and "*consumption*" results; or else, an attack of pneumonia being produced by some exposure, the exudation is cellular rather than fibrinous, and catarrhal pneumonia, ending in the majority of cases in a more or less rapid phthisis, occurs.

As already stated, it is especially in these conditions of system that cod-liver oil is so extremely useful. Preceding the development of active disease in these cases there is very generally a recognizable stage, marked by weakness, a tendency to emaciation, more or less anaemia, and other symptoms, which the present is scarcely the place to consider in detail. In this stage, cod-liver oil is exceedingly efficacious, and its use, combined with proper hygienic measures and the exhibition of other suitable drugs, may often succeed in warding off fatal disease.

In *scrofulosis*, it is useful in all stages, but should never be relied on to the exclusion of other drugs. Its effects are more marked during the *ulcerative* and *suppurative* stages, but in most cases it aids iodine very materially to reduce the enlarged glands in the earlier periods of the disease.

There are various diseases of the bones, dependent upon or resulting from a scrofulous taint, which are most favorably affected by cod-liver oil. Sometimes the disease attacks the articulating surfaces, giving origin to chronic inflammations of the joints,—*white swellings*; sometimes it is the body of the bones, especially of such as are composed chiefly of spongy tissue, that is affected, and *caries*, with subsequent abscesses, results. In these, as in all other forms of scrofulosis, cod-liver oil is almost a specific.

An affection probably not identical with, but closely allied to, scrofula, is *rickets*; cod-liver oil is of great value in this complaint.

In certain pale, cachectic children there may be found a swollen, tumid belly, perhaps with evident enlargement of the liver, and very generally, if not always, with enlargement of the mesenteric glands. This is the so-called *tubes mesenterica*, which is sometimes relieved, or even cured, by the exhibition of cod-liver oil.

The value of cod-liver oil in what is often very incorrectly called the “*pre-tubercular*” stage of phthisis has already been alluded to, but is so important that it will bear reiteration. There can be no doubt that consumption often commences with catarrh, and is often developed slowly as the result of frequently “catching cold.” Whenever a patient is feeble, pale, somewhat anæmic, complains of his liability to catch cold on the slightest exposure, even though no local disease exists anywhere, or rather because no local disease exists anywhere, there is cause for alarm; and it is of the most vital importance that the patient be put upon a tonic treatment whose basis is cod-liver oil, be fed upon nutritious diet, and have the hygiene of his daily life regulated, especial care being taken to avoid any exposure to cold. In the advanced stage of chronic *phthisis* the remedy is less efficacious, in that it much more rarely effects a cure than in the pre-tubercular stage; yet it does more good than all the other remedies of the Pharmacopœia combined,—alleviating the cough, increasing the strength, weight, and general health of the sufferer, often retarding or even arresting the pulmonic disorganization, almost always greatly prolonging life, and sometimes, in conjunction with other measures, effecting a cure.

It is a question of some importance to decide how the remedy does good in these cases. Its value, before the occurrence of any lesion, as a preventive of consumption, indicates that its influence during phthisis is not directly upon the local lesion, but upon the general condition of the system. This inference is borne out by clinical experience. The general symptoms commence to improve before the local lesions, and sometimes, although the patient fattens and gets stronger for a time, the pulmonic affection steadily increases; again, in some instances the oil fails to increase the weight of the patient or sensibly to affect the general nutrition, and in these cases it never does any good. In true *tuberculosis*, cod-liver oil, like all other remedies, is of very little, if any, value.

In cases of *defective nutrition*, when pallor, anæmia, loss of strength, and,

perhaps, emaciation, occur without any obvious cause, cod-liver oil is often of great service, especially when the subject is a child. Indeed, in children these symptoms are simply the result of a very mild action of the same depressing causes whose more intense malign influence produces scrofula.

In persons broken down with any of those chronic diseases which take the form of a dyscrasia, the remedy is often of great service. Thus, in the cachexia of *tertiary syphilis* it is invaluable.

The disease by whose relief and cure cod-liver oil first achieved its reputation is *chronic rheumatism*. I think, however, that it is much more efficacious in those cases in which the rheumatic disease has been grafted upon a scrofulously-tainted constitution, or in those cases in which the patient has been broken down by the disease, or by other agencies, so that there is what may be styled a general nutritive dyscrasia, than in simple chronic rheumatism; yet in obstinate *sciatica* and *lumbago* trial of it should always be made. In *gout*, oleum morrhue is of little service, and should be used only in the very chronic form of the disease, and when there is a generally disordered nutrition.

In *nervous affections*, especially in *neuralgia*, in *skin-diseases*, in fact, in any chronic disorder in which the patient is feeble and presents a condition of general depraved nutrition somewhat similar to that seen in consumption, cod-liver oil may be exhibited with advantage.

ADMINISTRATION.—The dark oil has been esteemed most highly by some authorities, especially by De Jongh, who asserts that it contains more of the biliary principles than does the pale oil, and even believes that the products of decomposition in it increase its beneficial action. It is, however, never employed at present, because of its exceedingly repulsive taste, and because it is very prone to disagree with the stomach. One of the difficulties in the use of even the pale oil is the very common real or imagined inability of the patient to take it. Without doubt, this very often arises from its nauseous taste, to lessen or disguise which various expedients are resorted to, with more or less doubtful success. Sometimes a piece of salt taken into the mouth just before the oil, which is also immediately followed by another lump of salt, suffices. It is said that some prefer the oil in emulsion made with some strong aromatic water. The addition of an equal part of glycerine and a half to one drop of the oil of bitter almonds to the dose certainly lessens the taste of the medicine. Some patients take it best in the froth of ale or porter, the glass being first half filled with the malt liquor, then the oil being carefully floated on the top of it without touching the sides of the glass, and the remainder of the vehicle put upon the top of it. Most of the patients requiring oil are also benefited by the use of alcohol; and my experience with the remedy is that the most generally successful plan of exhibition is to place, according to the exigencies of the case, from one to three tablespoonfuls of whisky or brandy in a tumbler, add not so much water, put the oil in the centre, and *toss* the whole down the throat, the head being held

well back, the mouth wide open, and the lips not touched by the medicine. The stimulus of the alcohol often enables the stomach to digest the oil when otherwise it could not do it. Sometimes it is necessary to commence with a single small daily dose, even a single teaspoonful, which is best taken at bed-time, and gradually to increase the amount as the patient becomes habituated to it. Children almost always learn to tolerate the taste of the oil, or even become in a short time fond of it.

The usual dose is for an adult a tablespoonful three or four times a day; for a child one year old, a teaspoonful.

ACIDUM PHOSPHORICUM—PHOSPHORIC ACID.

Phosphoric acid, which results from the burning of phosphorus in the air, is prepared by the action of sulphuric acid upon bone-ash, which consists chiefly of the phosphate of calcium. There are three varieties or forms of it: the *monobasic*, which exists in combination with one part of water, and which unites with a single part of a base; the *bibasic*, which requires two equivalents of water or of a base; and the *tribasic*, which unites with three parts of water or of a base. Phosphoric acid is officinal in two forms.

ACIDUM PHOSPHORICUM GLACIALE, or *Glacial Phosphoric Acid* of the U. S. Pharmacopœia, is the monobasic or *metaphosphoric* acid of the chemists. It is a colorless, transparent, ice-like solid, of a very sour taste, slowly deliquescent, and slowly soluble in water. It is the only variety of phosphoric acid capable of coagulating albumen, and is used in medicine only for the preparation of the *Acidum Phosphoricum Dilutum*, or *Dilute Phosphoric Acid*, which is commonly made by the action of nitric acid on phosphorus, but may be formed by the influence of a very small quantity of nitric acid upon the monobasic phosphoric acid. Dilute phosphoric acid is tribasic. It is a colorless, inodorous, sour liquid, of a syrupy consistence, which has a very acid reaction, but is not corrosive to animal tissues.

THERAPEUTICS.—Phosphoric acid has been used to a considerable extent abroad as a tonic and alterative in *scrofulous affections*. Upon the digestive organs, in my experience, it has little or no effect, and I have never been able to perceive that it is at all astringent or exerts any alterative influence upon the glands of the alimentary canal. In scrofulosis and rickets it is, I think, inferior to the phosphates.

PHOSPHATE OF CALCIUM.—The phosphate of calcium is, as is well known, an essential ingredient of bones, of which, indeed, according to the analysis of Berzelius, it forms more than fifty per cent. (*Traité de Chimie*, Paris, 1833). It should not be forgotten, however, that it exists in notable quantity in all the tissues, and is probably as essential an ingredient of their structure as of that of bone. Whenever it is taken out of the food of animals, although they be otherwise well fed, sooner or later they waste, sicken, and die. Chossat fed pigeons exclusively on corn containing very little of the

phosphate of calcium, and found that after some months they wasted, were affected with diarrhoea, and died (*Comptes-Rendus*, t. xiv.).

According to Roloff (*Virchow's Archiv*, Bd. xlv., p. 302), a herd of cows which had been fed upon hay from a certain meadow were very much out of health, and suffered from *fragilitas ossium*. On examination, the hay was found to be nearly free from earthy salts, and upon bone-meal being given to the cows they recovered their health in four weeks. The same authority further states that, in some meadows with which he is acquainted, the disease is endemic among the cows because the grass is so poor in phosphates. Haubner also affirms (*Schmidt's Jahrbücher*, Bd. cli. p. 138) that cattle fed exclusively upon potatoes, or upon roots very poor in phosphates, fail to fatten, become weak, and are apt to suffer from caries, but that if the phosphate of lime is given they rapidly improve.

Hegar (*Schmidt's Jahrbücher*, Bd. cli. p. 138) has considered the absorption of the phosphate of calcium, when given as a medicine, very doubtful, because when he exhibited it freely there was no increase in the amount of the phosphoric acid or of the earthy bases in the urine. Böcker (*Ibid.*), on the other hand, has found that if the drug be given to those wet-nurses whose milk contains an abnormally small amount of phosphates, the milk soon becomes rich in the earthy salts. Further, Albert Riesell (*Hoppe-Seyler's Medicinisch-chemische Untersuchungen*, p. 318) has shown that the phosphates are eliminated by the intestines, and that therefore the fact that the renal excretion of them is not augmented does not prove that they are not absorbed. The researches of Prof. Beneke* (*Schmidt's Jahrbücher*, Bd. cli. p. 138) are said to have shown that in many diseases the increased elimination of the earthy phosphates through the kidneys, which plainly occurs, is not accompanied by any increase in the amount ingested in the food, or decrease of the amount eliminated by the intestines, and that, consequently, there is a very decided wasting of the normal phosphates of the body. This being so, the use of phosphates in these diseases is as rational as that of iron in anaemia.

THERAPEUTICS.—According to Dusart (*Archives Générales*, 6e sér. t. xv.) and to Beneke (*loc. cit.*), the diseases in which the phosphate of calcium is especially indicated are *rachitis*, *osteomalacia*, and *scorfulosis*. It is evident that the indications for the earthy salts are very strong in the first two of these affections; and clinical experience has certainly borne out the results of *a priori* reasoning.

In *scorfulosis*, the call for the drug is not so plain; but Prof. Beneke states that in many cases, if the urine be examined, it will be found to be abnormally rich in earthy phosphates, and that under these circumstances the remedy is of the greatest value. Cases are not rare of children of slow development, often seemingly well nourished and robust, and yet really pale and with flabby

* I have not had access to the original memoir of Beneke, *Zur Würdigung des Phosphors. Kalkes in physiolog. und therapeut. Beziehung*, Marburg, 1870.

flesh, but without any distinct symptoms or marks of scrofulosis or of rachitis. Under these circumstances, the child is in a condition allied to that of the diathesis spoken of, and of the value of the phosphate of lime I have no doubt.

In cases of *delayed union* after *fracture*, the present remedy is seemingly indicated, especially since Dusart (*loc. cit.*) has experimentally proven that when given to animals whose bones have been broken it hastens union and makes the callus abnormally heavy and firm.

The phosphate of lime has been recommended in various diseases other than those mentioned, but its value is much more doubtful. Bennett commends it in *chronic phthisis*; Piorry (*Journ. de Chim. Méd.*, t. ix., 1863), in *syphilitic periostitis*; and Beneke, in *syphilitic gummata*; Schönian, and also Kugelmann, in the *menorrhagia* of anæmic women. Prof. Beneke calls attention to the use of it during *pregnancy*, and believes that it exerts an influence on the fœtus, so that women who have borne, it may be, only rachitic or scrofulous children will bring forth healthy offspring.

ADMINISTRATION.—The U. S. Pharmacopœia recognizes the *Precipitated Phosphate of Calcium* (*Calcii Phosphas Precipitata*), a white, inodorous, tasteless powder, which is prepared by dissolving bone-ash in muriatic acid and precipitating with ammonia. This may be employed in doses of ten grains three or four times a day, but, owing to its insolubility, is not so useful as the so-called *lacto-phosphate of lime*. This preparation, originally suggested by Dusart and Blache (*Archives Générales*, t. xv. p. 67), is made by the action of lactic acid upon the phosphate of lime, and was found by those experimenters to be soluble in all proportions, not only in water but also in the gastric juice. There is prepared by the druggists in this city an emulsion containing fifty per cent. of cod-liver oil and two grains of the lacto-phosphate of lime to the drachm, which has appeared to me to be the best of all the alterative preparations in cases of the character spoken of in the section on therapeutics. It certainly is very often more easily digested than the pure oil. The dose is a teaspoonful to a tablespoonful, or even more, according to the age.

Under the name of *chemical food*, or *compound syrup of the phosphates*, a very complex preparation has been much used in disorders attended with impaired nutrition, such as the lacto-phosphate of lime has been recommended in. I have had no experience with it, but very much doubt its being superior, or even equal, to the latter drug.

COLCHICI SEMEN—COLCHICUM SEED.

COLCHICI RADIX—COLCHICUM ROOT.

Colchicum autumnale, whose products the above drugs are, is a little plant growing in Europe and England, where it is commonly known as the *meadow saffron*.

It is not really the root of the plant which is officinal under the name of colchicum root, but the thickened swollen end of the stem, with the little bulblet whose office it is to develop a new plant. This *corm* is solid and fleshy, an inch and a half to two and a half inches in length, with a longitudinal groove, having a nail-like process (the bulblet) at its base. In the shops it is very commonly kept in transverse slices, which are notched and cordate; the taste is bitter, hot, and acrid. Colchicum seeds are nearly round, about an eighth of an inch in diameter, and of a bitter, acrid taste. The active principle of both seed and root is an alkaloid, *Colchicia*, whose individuality was first made out by Geiger and Hesse. According to Hübler, it is slowly soluble in water, readily so in alcohol, and not at all so in ether. By the action of mineral acids and by other agencies it is resolved into a brownish-green insoluble resin and a crystallizable neutral substance, soluble in water, *Colchicina*.

Colchicia makes with concentrated nitric acid a violet solution, which when diluted with water becomes yellow; with concentrated sulphuric acid it strikes an intense yellow color.

PHYSIOLOGICAL ACTION.—When taken in dose of sufficient size, colchicum acts upon man as a poison, producing repeated, uncontrollable vomiting, with nausea and retching, and also violent purging of at first serous character; afterwards the passages become smaller, more mucous, with flakes in them, and finally in some cases bloody. Abdominal pain may be absent or present, but if present is generally griping; sometimes there is gastric burning.

Nervous symptoms have been prominent in some of the severe cases. In one instance, it is said, a feeling of numbness or prickling was complained of by the patient; but this seems not to be common. Spasms are very frequent, and sometimes convulsions, which may be fatal, are present. Muscular pains are not rarely experienced, in some cases replacing the spasms, and probably in all other cases coincident with them is great muscular weakness, amounting, as death approaches, to paralysis.

Finally, a condition of collapse develops itself, the circulation fails more and more, the pulse, which has been frequent and feeble, becomes rapid and thready, the skin cold, pale, or livid, bedewed with sweat, and death from exhaustion results. Consciousness is preserved until the last. The effect of lethal doses of colchicum on the urinary secretion varies; sometimes the kidneys seem to be nearly unaffected almost to the last; sometimes their functional activity is decidedly increased, but in other cases it is diminished, and even suppression of urine has been noted.

The symptoms produced by the largest therapeutic doses of colchicum are slowing of the pulse, nausea, vomiting, abdominal uneasiness, borborygmi, and free purging, together with a sense of prostration and of weakness.

The occurrence of the inflammatory changes after the hypodermic injection of *colchicia* proves that the irritant principle of colchicum acts by absorption, and that the intestinal symptoms are not simply due to a local action,—deductions confirmed by the chemical experiments of Aschoff, who found *colchicia* in

the liver, kidneys, heart, lungs, and blood of a rabbit poisoned with one and a half grains of it.

A very curious fact noticed by Schroff, which, if it be confirmed, shows that colchicia really has little or no direct influence upon the nervous system, is that the rapidity of death is not at all in proportion to the size of the dose. Thus, in rabbits the fatal result occurred after one and a half grains of the alkaloid, in fourteen hours; after fifteen grains, in eleven hours. This seems to be explicable only by the supposition that colchicia kills solely by virtue of its irritant action on the alimentary canal, and, not being in any dose corrosive, requires time to work out the fatal result, through the instrumentality of a gastro-enteritis.

This deduction is confirmed by the long-protracted course of the poisoning after small doses. Thus, Aschoff noted death on the ninth day in a pigeon which had received one-fourth of a grain of the alkaloid.

After death from colchicum, the blood is generally found very dark and imperfectly coagulable; but whether this is due to a direct action of the poison, or is the result of the slow death by asphyxia and exhaustion, is uncertain. According to Heinrich,* 0.15 grain of colchicia will produce poisonous symptoms in man, and in Krahmer's experiments (*Journal für Pharmakodynamik*, ii. p. 561) 0.3 grain caused in an adult violent serous purging, lasting for four days, and accompanied with severe tenesmus.

Upon most animals colchicum acts very much as it does upon man, producing, in poisonous doses, as prominent symptoms, severe and often bloody purging, vomiting, great prostration, embarrassed respiration, finally more or less pronounced paralysis, and death, not rarely preceded by convulsions. Reflex actions, according to Albers, are lessened, and finally abolished, in the frog; and Schroff makes similar assertions in regard to the rabbit. In no case do reflex spinal convulsions occur at any time.

Geiger (*Annal. Chem. Pharm.*, vii. 274), Hoppe, Aschoff (*Vierteljahresschrift für Prakt. Pharm.*, vi.), Schroff (*Oester. Zeitschrift f. Prakt. Heilk.*, 1856), and Albers (*Deutsches Klinik*, 1856, xxxvi.), have experimented with colchicia, and have shown that it causes symptoms similar to those produced by colchicum,† of which it is without doubt the active principle. Unfortunately, their experiments have not been carried far enough to enable us to make out what is the exact influence of the drug upon the various portions of the body.‡ They all agree in that the chief force of the poison is expended upon the alimentary canal, at least in mammals, and that after death, even when the alkaloid has been given hypodermically, the intestinal mucous

* Quoted by Husemann.

† Dr. R. Lewins's experiments (*Edinburgh Medical and Surgical Journal*, vol. lvi., 1811) may be used for comparison. He used colchicum.

‡ Unfortunately, the works of nearly all these investigators are known to me only at second hand. Not having had access to the original papers, I have been forced to depend upon abstracts in various journals.

membrane is found much inflamed, as is also very frequently the inner coat of the stomach.

The action of poisonous doses (one to two grains) of *colchicina* has been studied upon dogs by Dr. Saml. R. Percy. The symptoms are very similar to those produced by *colchicia*: they are increase in the frequency of the pulse, severe purging, with tenesmus, vomiting, finally great slowing of the pulse and failure of the heart's action, and death without convulsions. The urine, at first increased, was afterwards suppressed. On post-mortem examination, the mucous membrane of the intestines was found highly inflamed, that of the stomach slightly so, and the heart and arteries were filled with black tarry blood, similar to that of *colchicia*-poisoning.

The main interest to the therapist in the physiological study of colchicum, of course, is in regard to its action in small therapeutic doses. The most prominent result of the ingestion of such amounts is gastro-intestinal disturbance, as shown by abdominal uneasiness, colicky pains, borborygmi, loss of appetite, moderate purging, and sometimes nausea,—symptoms differing in degree only from those of poisoning by the drug. Before they come on, however, there is a lowering of the pulse-rate, sometimes as much as twelve beats per minute. Upon the skin the medicine occasionally acts, producing in some cases *diaphoresis*, and it is said that the amount of this action is in inverse ratio to the effect on the bowels. According to Schrott, the one-hundredth of a grain of *colchicia* is rather more than the therapeutic dose, and produces purging, lasting for several days, with griping pains, cerebral distress, a pulse at first lowered but afterwards accelerated, and a secretion of thick lateritious urine. Any nervous symptoms, such as vertigo, headache, muscular weakness, which may be present as the result of the administration of colchicum, are probably sympathetic upon the gastro-intestinal irritation.

It is evident that colchicum influences the bowels powerfully, and probably in this way acts as an eliminative. In the minute doses often used with great advantage in disease, purging, however, does not occur, and consequently increased elimination, if it takes place, must be through the kidneys: great interest therefore attaches to the influence of the remedy upon the urinary secretion.

In considering this, the effects of poisonous and of therapeutic doses must not be confounded, for it is very evident that an irritation which causes suppression of urine may, when present in a much milder degree, produce an increased flow.

It seems very certain that in moderate doses, repeated at regular intervals, colchicum very often increases the flow of urine. In 1828, Chelius announced that during its administration in gout the amount of uric acid eliminated is nearly doubled. Dr. R. Lewins (*Edinburgh Medical and Surgical Journal*, 1841, vol. lvi. p. 200) submitted the urine of several persons suffering from gout, taken before and after the administration of colchicum, to Prof. Chris-

tison, who found in the colchicum-urine the *proportion* of urea nearly double, and that of uric acid greater,* than that of the other specimens.

In 1852 Dr. Mackagan (*Edinburgh Monthly Journal of Med. Sci.*, 3d series, vol. xiv. p. 24) analyzed the urine of three cases of rheumatism before and after the exhibition of colchicum: in two instances the *proportion* of urea was very greatly increased, that of uric acid slightly so. In the third case the effect just noted happened at first, but not afterwards.

On the other hand, Prof. Stillé states that Graves and Gardner affirm that the urates diminish under the use of the medicine. It is evident that these different results are not so contradictory as they seem, for it is possible that in one case the colchicum may so act as to increase the elimination of urea, in another that of uric acid; and that when one of these is increased the other may be unaffected, or even diminished.

Further, when the medicine purges freely it is very probable that elimination by the kidneys is lessened; and no account of this is taken by any of the observers whose original papers I have seen. Moreover, these observers also all contented themselves with noting the proportion of urea and uric acid in the urine, when it is evident that the mere proportion, unchecked by the absolute amount of urine secreted during the twenty-four hours, is no criterion as to the absolute amount eliminated. Although it seems to me probable that in many cases colchicum does increase in gout and rheumatism the solids in the urine, yet so many evident sources of fallacy have hitherto been overlooked, and even when kept in view are so difficult to avoid, that far more precise investigations are needed to settle the question.

The action of the drug upon the urine during health is evidently very closely connected with the question just discussed. Here again we find conflicting testimony. Dr. Bird (*Urinary Deposits*, Phila., 1859, p. 354) quotes Prof. Kramer's† experiments as showing that colchicum does not increase the amount of solids eliminated, and intimates that his own investigations had given similar results. Dr. Hammond (*Proc. Biol. Dept. Acad. Nat. Sci.*, Dec. 1858), on the other hand, in a series of experiments in which every care to avoid fallacies, by maintaining equality as to diet and exercise, was observed, found that while squill and digitalis only increased the watery part of the urine, both the organic and the inorganic solids were remarkably increased by colchicum.

In regard to *colchicine*, the experiments of Prof. S. R. Percy (*Amer. Med. Times*, April, 1862, p. 167) indicate, but are much too few to prove, that in gout it increases the elimination of urea and uric acid.

* Dr. Harley (*The Urine and its Derangements*, Phila., 1872, p. 81) makes the assertion that colchicum diminishes the excretion of uric acid and even the urea: as, however, he does not deem it necessary to give any authority for the assertion, and as he does not appear to have made any elaborate chemical examination himself, not much weight is to be attached to his testimony.

† Kramer's paper was published in *Keller's Archives*, Dec. 1847, and is entirely inaccessible to me.

THERAPEUTICS.—Our knowledge of the use of colchicum in disease is purely empirical, based upon clinical experience; and our acquaintance with its physiological action is not sufficient to enable us even to explain fully what experience has taught, much less to guide us in our use of the drug. *Gout* is the one disease in which colchicum is almost universally recognized as a specific. It may be advantageously employed both as a preventive of the paroxysm and to lessen its severity when developed.

During an attack of gout, from ten to twenty drops of the wine of the colchicum root may be exhibited every four hours until some decided evidence of its action, such as nausea or slight purging, is induced.

It should always be borne in mind that although looseness of the bowels may be useful, yet when colchicum purges the gouty patient actively it mostly fails in achieving the desired therapeutic result. Its action is most favorable when its influence is felt chiefly upon the skin and the kidneys. To effect this desired result, it is often well to restrain the tendency of the drug to act upon the bowels, by combining it with opium. This is especially the case in debilitated subjects, in whom anything like over-purgation must be avoided with the most scrupulous care.

By large purgative doses of colchicum the paroxysm of gout may often be suppressed; but experience has shown that this use of colchicum is dangerous, the suppression being sometimes followed by serious internal disease, apparently due to a transfer of the gouty irritation. Between the paroxysms, colchicum may be steadily exhibited to the gouty subject in small doses (ten drops of the wine of the root three times a day); and often great advantage is derived from its combination with the iodide of potassium. This combination is especially useful in irregular atonic gout, such as is most frequently seen in women of feeble nervous organization who have inherited the diathesis, but is sometimes present even in robust men. Ten grains of the iodide and ten drops of the colchicum wine may be given three times a day. Speculations as to how colchicum cures gout seem to me useless in the present state of our knowledge: until we know more of the physiological action of the drug and of the nature of the disease, one theory seems as good as another.

In *rheumatism*, colchicum has been highly recommended by some, but has never come into such general use as in gout. In the inflammatory variety of the disease it is of but little value, except in purgative doses, and is mostly administered in the form of *Scudamore's Mixture*, which is composed of magnesia and its sulphate with wine of colchicum root. In *subacute rheumatism*, the combination of colchicum and iodide of potassium, already spoken of, is very useful.

Colchicum has been administered in various diseases, but when there is no rheumatic or gouty taint is at present very rarely used.

TOXICOLOGY.—The symptoms of poisoning by colchicum have been already enumerated. The fatal dose varies, but is small. Prof. Geo. B. Wood (*United States Dispensatory*, 13th ed., p. 1504) states that death has

been produced by two drachms and a half of the wine of colchicum root, and Taylor (*Medical Jurisprudence*, 2d ed., vol. i. p. 330) records a case in which three drachms and a half proved fatal. On the other hand, recovery has taken place after the ingestion of an ounce.*

According to the experiments of Schroff, *colchicia* is eighty to one hundred times stronger than the fresh corm. Casper has seen death result from a quantity of the wine containing 0.025 to 0.03 gramme (0.37 to 0.45 grain) of colchicia; but, according to Husemann, recovery has taken place after the ingestion of 0.045 gramme of the alkaloid. Very recently (Dr. Geo. W. Major, *Canada Medical and Surgical Journal*, Dec. 1873) seventeen cases of poisoning from one bottle of wine of colchicum seeds occurred in Montreal, seven proving fatal. The patients had been vomiting and purging almost continuously for many hours when first seen, and the symptoms were exactly those of the stage of collapse of severe cholera morbus. In no case was the purging bloody. Consciousness was preserved to the last, and in only one case was there anything like convulsions. There was decided numbness of the extremities, and a peculiar hoarseness of the voice was especially noted.

The treatment of colchicum-poisoning is as follows. If the stomach and bowels have not been freely evacuated, administer at once an emetic and a cathartic, so as to empty the alimentary canal; allow the patient to drink freely of warm water, to aid in these operations and to act on the kidneys. Give freely of tannic acid, as the only known chemical antidote; although experiments upon animals have shown that it is not to be relied upon. To check the vomiting and purging, administer opium freely; and to allay the irritation, cause the patient to drink freely of albuminous matter, such as white of egg dissolved in water: the tannic acid having been given as soon as possible after the taking of the poison, the demulcents are useful in the more advanced stages. Symptoms of gastro-enteritis or of collapse are to be met as they arise.

ADMINISTRATION.—Colchicum is never used in substance, but in one of the following preparations, of which the wine of the root is deservedly the most popular.

Preparations from the seeds:

Tinctura Colchici—*Tincture of Colchicum* (℥ii to Oj). Dose, half a teaspoonful to one and a half teaspoonfuls.

Vinum Colchici Seminis—*Wine of Colchicum Seeds* (℥ii to Oj). Dose, half a teaspoonful to one and a half teaspoonfuls.

Extractum Colchici Seminis Fluidum—*Fluid Extract of Colchicum Seeds*. Dose, two to six minims.

Preparations from the root:

Vinum Colchici Radicis—*Wine of Colchicum Root* (℥vi to Oj). Dose, ten to fifteen drops; as a purgative, half a fluidrachm.

* See case in *L'Union Médicale*, Aug. 1848.

Extractum Colchici Aceticum—*Acetous Extract of Colchicum*. Dose, one to two grains.

Extractum Colchici Radicis Fluidum—*Fluid Extract of Colchicum Root*. Dose, two to four minims.

SARSAPARILLA—SARSAPARILLA.

The root of *Smilax officinalis* and other species of *Smilax*, woody vines inhabiting Mexico and the northern portions of South America. There are in commerce a number of varieties of sarsaparilla, the two most important of which are the *Honduras* and the *Brazilian*. The former of these is almost the only sarsaparilla used in this country. It occurs in bundles two or three feet long, composed of several very long, thin roots, folded upon themselves, the whole being bound round by a number of turns of the root. The Brazilian sarsaparilla also comes in cylindrical bundles, each of which is closely wrapped about by a very flexible stem: it mostly has fewer rootlets than the Honduras variety. The crude sarsaparilla has little or no smell, but its taste, which is at first simply mucilaginous, soon becomes, if the root be chewed, persistently acrid. According to Prof. Geo. B. Wood, the degree of this acidity is the best measure there is of the activity of any specimen of the drug.

Sarsaparilla contains a crystallizable principle, first discovered by Palotta in 1824. From its discoverer this substance received the name of *Paraglin*. By subsequent authorities it has been variously called *Smilacin*, *Salseparin*, *Sarsaparillin*, and *Parallinic Acid*. Paraglin, according to Poggiale, crystallizes in fine needles; according to Thubeuf, in star-like clusters of plates. It has a neutral reaction, and in solution a bitter, acrid taste, but when solid is nearly tasteless. It is very slightly soluble in cold water, more so in boiling water; moderately soluble in dilute alcohol when cold, freely so when hot. Palotta asserts that thirteen grains of it cause vomiting, constriction in the throat, weakness, diaphoresis, and depression of the circulation. On the other hand, Böcker, of Bonn, has exhibited it in doses of a like amount without producing any symptoms whatever (*Journal für Pharmakodynamik*, Bd. ii. Heft i. p. 23).

PHYSIOLOGICAL ACTION.—It has already been shown that there is very little reason for believing that paraglin has much immediate action on the system; and the question naturally arises, Has sarsaparilla any more influence? The answer to this question must, I believe, be a negative one, since the only sensible effects that follow even the largest draughts of the decoction are simply the results of slight gastric disturbance. It has been claimed that sarsaparilla acts as a diuretic and diaphoretic; but the only record I have met with of any careful experimentation is that of Böcker (*loc. cit.*). That investigation seems to show conclusively that the drug has no marked influence upon these secretions. If, therefore, sarsaparilla have any value whatever in disease, it must be simply as an alterative,—as a remedy which in some unknown way modifies nutrition.

THERAPEUTICS.—Sarsaparilla has been used, and still is used, to such an enormous extent in medicine, that it seems impossible to believe that it is destitute of therapeutic virtue. It is not in accordance with the plan of the present work to enter into an elaborate discussion of the recorded clinical experience with it: suffice it to say that, although the evidence is contradictory, on the whole there is a decided preponderance in favor of the value of the drug in *chronic syphilis* and in *chronic scrofulous diseases*. I have used it largely, but always in combination with more powerful alteratives, so that it is impossible to decide how much of the good achieved has been due to its influence.

There are two distinct methods or objects of use of sarsaparilla in syphilis: one as an adjuvant to mercury in the secondary stage; one as an adjuvant to the iodide of potassium, or as a sole reliance, in the advanced tertiary cases, especially where the constitution is very much broken down by the disease. It is stated that, in the latter condition, very often during its use the appetite will gradually increase, the spirits rise, the secretions become more and more normal, and the strength grow day by day.

ADMINISTRATION.—Sarsaparilla is never given in substance, but in one of the following preparations:

Decoction Sarsaparillæ Compositum—*Compound Decoction of Sarsaparilla*.—This contains sarsaparilla, bark of sassafras root, guaiac wood, liquorice root, and mezereon, and is an imitation of the famous "*Lisbon Diet-Drink*." The dose is three or four fluidounces three or four times a day.

Syrupus Sarsaparillæ Compositus—*Compound Syrup of Sarsaparilla*.—This contains sarsaparilla, guaiac wood, pale rose, senna, liquorice root, oil of sassafras, oil of anise, and oil of gaultheria.

It is a very popular preparation, on account of its pleasant taste: it affords the only vehicle I know of capable of disguising the taste of the iodide of potassium.

The dose of it is one to two tablespoonfuls three or four times a day.

Extractum Sarsaparillæ Fluidum—*Fluid Extract of Sarsaparilla*.—Dose, half a teaspoonful three times a day.

Extractum Sarsaparillæ Fluidum Compositum—*Compound Fluid Extract of Sarsaparilla*.—This contains sarsaparilla, liquorice root, sassafras, and mezereon. The dose is a teaspoonful.

GUAIACI LIGNUM—GUAIAO WOOD.

GUAIACI RESINA—GUAIAO RESIN.

The wood and the resin of the *Guaiacum officinale*, a large tree growing in the West Indies, are both included in the *Materia Medica* list of the U. S. Pharmacopœia. Guaiac wood, or *lignum-vitæ*, is imported in billets, but very generally is kept in the shops in the form of raspings or shavings. It is a very dense wood, the central heart-wood having a dark-olive or brown-

ish-green, and the outer sap-wood a light-yellowish color. It is inodorous, but becomes somewhat fragrant when rubbed or heated. Besides the resin, it contains an extract which is believed to have medicinal properties.

Guaiaec resin is obtained to a slight extent by spontaneous exudation from the living trees; much more largely by boring a hole into the centre of one end of a billet, placing the other end in the fire, and catching the melted resin as it runs out; and still more frequently by boiling the chipped wood in salt and water and skimming off the resin as it rises to the surface. Guaiaec occurs in irregular lumps or masses, of a dark reddish-brown greenish color externally, offering a conchoidal fracture with somewhat translucent edges. The odor is feeble, peculiar, agreeable, increased by heat. The taste is at first very slight, but, as the resin melts in the mouth, becomes very acid and persistent. It is a very complex body, containing three acids, the *guaiaconic acid* of Hadelich, the *guaiaec acid* of Righini, and the *guaiaecresinic acid* of Hlasiwetz, besides *guaiaec yellow*, a peculiar resin, and other substances. Landerer asserts that he has found in it a peculiar crystallizable substance, which he calls guaiaecin (Husemann, *Die Pflanzenstoffe*, p. 1106.)

When a lump of guaiaec is freshly broken, it offers a dark, blackish surface, which on scraping or bruising becomes yellowish; on exposure, the well-known greenish tint is acquired, owing to a spontaneous oxidation. The ease with which the resin undergoes oxidation is its most distinctive characteristic. As already stated, the change occurs on simple exposure to the light. According to Wollaston, it is most rapid and perfect in the focus of the violet rays of the spectrum, whilst in the focus of the red rays the original color is regained. Oxidizing or ozonizing agents, such as nitric acid, chromic acid, iodine, bromine, and chlorine, produce this oxidation very rapidly and very thoroughly, the resin acquiring an intense blue color.

PHYSIOLOGICAL ACTION.—Guaiaec is believed by some to act as a diaphoretic, to do good by increasing the elimination of the skin; but, as I have not been able to obtain either from medical literature or the exhibition of the medicine any distinctive proof of its having such action to any marked extent, I have preferred to consider the drug as an alterative. When taken internally, very little sensible effect results, unless the dose be so large as to irritate the stomach.

THERAPEUTICS.—Guaiaec has been very largely employed in *chronic syphilis*, in connection with sarsaparilla, and is an ingredient of its most favorite preparations.

Guaiaec has also had a good deal of repute in *rheumatism*, and in the sub-acute and chronic forms of the affection is often of service.

For an account of its employment as an emmenagogue, see the chapter upon that class of remedies. Guaiaec is best administered in tincture, either the simple (*Tinctura Guaiaci*) or the ammoniated (*Tinctura Guaiaci Ammoniata*). The dose of either of these is from one to two teaspoonfuls, administered, preferably in milk, three or four times a day.

MEZEREUM.—*Mezereon* is the bark of two species of *Daphne*, native shrubs of Europe. It is a thin, grayish, tough, flexible bark, occurring in long strips folded upon themselves, nearly odorless, but having a very acrid taste. It contains a neutral, crystallizable, bitter glucoside, *Daphnin*, besides a volatile acrid principle. *Mezereon* is intensely irritant, and its ointment (*Unguentum Mezerei*) is used as a stimulant dressing to very indolent ulcers, and to keep blisters sore or to maintain issues. Internally, in small doses it is believed to be an alterative, but is never used except in combination with sassaaparilla. In overdoses it is an active poison, producing severe vomiting, purging, and gastro-intestinal inflammation.

SASSAFRAS.—The bark of the root of the *Sassafras officinale* is believed by some to be an alterative, on, as I think, very slight foundation. It contains some tannic acid and very largely of a volatile oil (*Oleum Sassafras*), which, on account of its cheapness and agreeable odor and taste, is much used both in medicine and in the arts as a flavor and perfume.

TARAXACUM.—The root of the common dandelion, *Taraxacum officinale*, is included in the *Materia Medica* list of the U. S. Pharmacopœia, and is believed to have the property of altering the action of the liver; although no effect is to be witnessed from a single dose of the drug, let it be never so large, other, at least, than some nausea. Diuretic properties have also been ascribed to taraxacum; but the only evidence brought forward to establish this is the vulgar name which the plant bears both in English and in French. If useful at all, it is in those cases of dyspepsia in which there is habitual torpor of the liver, with costiveness. It must be given very freely and continuously for weeks before any good effect is to be looked for. The U. S. Pharmacopœia recognizes a fresh juice *extract* (*Extractum Taraxaci*) and a *fluid extract* (*Extractum Taraxaci Fluidum*), either of which may be used in doses of one or two drachms, administered after meals.

DIVISION II.—LOCAL REMEDIES.

CLASS I.—EMETICS.

EMETICS are those drugs which are employed in the practice of medicine for the purpose of producing emesis, or vomiting. The mechanism of vomiting has been so frequently written upon, and has so little connection with the application of emetics, that it is not necessary here to enter upon an elaborate discussion* of it. Suffice it to state that emesis is the result of a very complicated series of actions, in which the chief expulsive force is supplied by the abdominal muscles and the diaphragm,—the stomach, however, participating in the general contraction, and not being, as some have thought, entirely passive. The exact relations and functions of the various nerves concerned are not, I think, fully made out. It has been generally believed that the pneumogastrics were the afferent nerves, and that, although emetics introduced into the circulation after their section acted, yet irritation of the gastric mucous membrane was not capable of so doing. But Schiff found in his experiments that, even when the nerves were cut in the neck, the introduction of semi-solid food into the stomach gave rise to efforts at vomiting, which were in some cases successful; and MacLagan† has obtained similar results with the sulphates of zinc and copper. Further, I have invariably failed to induce vomiting with veratria, even when given immediately after section of the par vagum. Evidently, further investigations are needed.

Vomiting occurs under two provocations, or in two manners. Thus, a mental impression, or a disordered state of the blood, may influence the nerve-centres directly, and emesis, spoken of as *centric*, results; or a peripheral irritation in the stomach itself, or in some other organ, as in the kidneys, may induce vomiting precisely similar in the method of its production to the more ordinary reflex movements; such vomiting is called *reflex* or *eccentric*.

Emetics produce their results in both of these methods. Thus, tartar emetic affects the centres directly, and causes centric vomiting. Sulphate of copper irritates the mucous membranes of the stomach, and reflex vomiting is

* For a very elaborate discussion of this subject and other matters relating to the action of emetics, see Prof. Joseph Carson's Lectures in the *Phibada. Medical Times*, June, 1872.

† *The Action of Medicines in the System*, by F. W. Headland, M.D., Amer. ed., 1839, p. 110.

induced. It is evident that medicines of the former class must be absorbed to act; whilst those of the latter, often spoken of as "*mechanical emetics*," act without absorption. It is also obvious that the latter are more prompt than the former, and it is a clinical fact that they act more certainly when the nerve-centres are obtunded, as in narcotic poisoning, and that they always cause less nausea and general systemic disturbance than do the centric emetics.

Another evident practical fact is, that whilst centric emetics will act in whatever way they are introduced into the system, the mechanical must be exhibited by the stomach. Thus, apomorphia may be given by hypodermic injection, but mustard must be taken by the mouth.

In regard to the phenomena of vomiting, there are a few points to which it is necessary here to call attention. First of these is the fact that *nausea* always produces, or is accompanied by, muscular relaxation. Vomiting may exist, as from mustard, without much relaxation; but when it is accompanied by much nausea the whole system is as it were unbent, the skin soft and bedewed with perspiration, the pulse soft and feeble, the muscular system limp and incapable of exertion, the mental acts almost suspended. During violent vomiting the blood is driven to the head, so that the whole exterior, and probably the interior also, of the cranium becomes very much congested. The abdominal circulation also is very much affected, and the blood is as it were squeezed out of the portal vein and its tributaries. The matters rejected consist of the contents of the stomach, and, in repeated vomiting, also those of the duodenum. The secretion from the gastric mucous membrane is very much enhanced, and without doubt is more or less modified. Bile in ejecta is to be recognized by the green color and the bitter taste, or more infallibly by testing with the proper reagents.

The indications for the use of emetics are as follows:

1. *To unload the stomach.*—For this purpose they are employed in poisoning; in the existence of crude articles of food or indigestible substances in the stomach; or in the presence of acrid, perverted secretion. The symptoms induced by irritating materials in the stomach are various, and sometimes it requires a good deal of tact or experience to recognize their cause. Among them may be mentioned a feeling of weight or load in the stomach, gastric distress, or severe cramp or spasmodic pains, with or without some nausea and retching. In other cases no local manifestations of trouble may be present. Thus, *convulsions* in children are very frequently the result of gastric irritation, and are at once relieved by emptying the stomach. In adults, *apoplectic form coma* may offer a similar history. Occasionally *urticaria*, or hives, and not rarely severe *headache*, have a similar origin, and require a similar treatment, to the conditions just mentioned.

2. *To affect the abdominal viscera and circulation.*—Emetics have been recommended by some in *congestions* of the *spleen*; but evidence is wanting as to their power materially to affect other viscera than the liver.

In *congestion* of the *hepatic* and *portal* circulation, not dependent upon

organic cause, and in the condition of digestive derangement known as "*biliousness*," they are often of service. In *catarrhal jaundice* they may effect much good by causing dislodgment of the mucus plugging the ducts. They have been employed in cases of *biliary calculi*; but the chances of forcing out the calculus by external violence are probably no greater than those of lethal rupture of the gall-bladder.

3. *To dislodge substances from the respiratory passages.*—For this purpose emetics are sometimes used when foreign bodies have found entrance into the larynx; but it is chiefly in *membranous croup* that the present indication is met with. The emetics chosen for this purpose should be such as act with violence without producing much nausea or systemic disturbance; the mechanical emetics are therefore the best.

4. *To produce muscular relaxation.*—The introduction of anaesthesia has rendered the use of emetics to meet this application almost obsolete. Occasionally, however, in *asthmatic* or other *spasmodic affections* of the respiratory organs, emetics are still employed. For this purpose the drugs causing much nausea are preferred. In adults, lobelia is the best; in children, ipecacuanha. Nauseating rather than emetic doses should be employed.

5. *To lessen arterial action and reduce inflammation.*—Almost the sole disease in which advantage can be derived from this use of emetics is acute bronchitis in its early stages. In very many cases a "cold on the chest" in its outset may at once be subdued by ipecacuanha, or, better still, tartar emetic: small doses should be given at short intervals, to produce continuous nausea, terminating after a time in vomiting. This method of cure is so disagreeable, although very efficacious, that patients will rarely submit to it, unless, as sometimes in the case of public speakers, relief within a short period of time be a matter of great importance. It should be added that the cure is wrought in these cases not merely by the lessening of arterial action, but also by the induction of free bronchial secretion.

6. *To create a shock to the system.*—Under this head may be included several empirical uses of emetics, in which advantage is gained, but in a method which is not very clear. Thus, in epileptic attacks, when the fits have a tendency to recur every few minutes, the unconsciousness persisting, it may be, for hours, emetics will sometimes break up the succession of disordered nervous action. Again, not unfrequently an ague-fit can be set aside by an emetic given just before its expected recurrence.

Contra-indications.—The chief contra-indications to the use of emetics are the existence of congestion of the brain, or of gastric inflammation. Advanced pregnancy or hernia, whilst they do not positively contra-indicate the use of emetics, should cause great caution to be practiced in their employment.

ADMINISTRATION.—Emetics should, as the general rule, be given in a full dose, so as to avoid unnecessary repetition, and should be administered dissolved in water or syrup. Their action should be assisted by frequent and copious draughts of tepid water, which also have the advantage of ren-

dering the vomiting less painful. When for any reason protracted nausea is desired, the doses should be small and repeated at short intervals.

Hyperemesis may advantageously be divided into two varieties: first, such as is due to overdoses of depressing centric emetics; second, such as arises from irritation of the stomach, as by mechanical emetics. The treatment of the first of these consists in the enforcement of absolute quiet in the horizontal position, the free use of opium enemata, the application of counter-irritants to the epigastrium, and the use of alcoholic stimulants. The latter should be given in hot water, and should not be too much diluted. I have seen raw brandy arrest at once the most alarming centric emesis, after the failure of other methods. Creasote, chloroform, or chloroform and volatile oils, are sometimes of value in this form of hyperemesis.

When excessive vomiting is due to some irritant emetic, the stomach should be thoroughly washed out by large draughts of warm mucilage. Opium should be exhibited by the rectum, a mustard plaster or blister, or, often better still, leeches, be applied to the epigastrium, and no medicine at all be taken into the inflamed viscus. The swallowing of small pieces of ice is sometimes of service. If these remedies fail, the treatment of this form of hyperemesis soon resolves itself into that of gastritis.

VEGETABLE EMETICS.

IPECACUANHA.

The root of the *Cephaelis Ipecacuanha*, a small, shrubby plant, growing in Brazil, where the drug is gathered by the Indians to be exported in large bales or bags. *Ipecacuanha* occurs in pieces of two or three lines in thickness, variously bent and contorted, marked on their surface with numerous prominent rings, and composed of an outer, thick, active, hard, and horny cortex, and an inner, light, inert, woody centre. Varieties of *ipecacuanha*—the *red*, the *gray*, and the *brown*—have been formed from the color of the bark, but the distinction is trivial.

The root has very little odor, but the brown powder has a decided and peculiar smell, and in some persons excites sneezing, or even violent asthmatic dyspnoea. The taste is bitter, acrid, and nauseous. The active principle is *Emetia*, an alkaloid first discovered by Pelletier in 1817. The cortex also contains small quantities of ipecacuanhic acid, which is related to tannic acid.

Pure *emetia* is a white, uncrystallizable, odorless powder, of a bitter, burning taste, soluble in one thousand parts of water at 50° C. (Lefort), freely soluble in dilute and absolute alcohol, in chloroform and benzole, scarcely so in ether. Its solution in acidulated water, according to Draggendorff, has a decided blue fluorescence. Its salts are, according to Pelletier, uncrystallizable.

Concentrated sulphuric acid turns it a dirty brown, nitric acid a yellowish brown. Pure *emetia* is very difficult to prepare, and, according to Mr. Wil-

liams (*St. Bartholomew's Hospital Reports*, vol. v.), only two grains of it can be obtained from an ounce of the root. The ordinary impure alkaloid of the shops occurs in brownish-red, transparent, very deliquescent scales, which are very soluble in water.

PHYSIOLOGICAL ACTION.—Locally applied, ipecacuanha is a decided irritant, manifesting its action not only upon mucous membranes and upon denuded surfaces, but even, when used by inunction, producing an eruption upon the sound skin. According to Dr. Dyce Duckworth, this eruption consists at first of small, discrete pustules, with a rather large areola; afterwards, if the application be persisted in, of large pustules, followed by severe ulceration. When exhibited in small repeated doses to man, it produces malaise, with nausea, and perhaps an increase of the secretions of the salivary glands and of the mucous membrane of the bronchial tubes and of the stomach. In large amounts it causes vomiting, accompanied by only a moderate amount of nausea but by a decided increase of the secretions mentioned above. The vomiting, even when very large amounts are taken into the stomach, is not apt to be severe, nor the prostration marked,—no doubt because the excess of the drug is rejected before absorption. In large doses of emetia, this mildness of action, in all probability, would not occur: certainly animals are readily killed by the alkaloid.

Although ipecacuanha was made known in 1649 by Piso, and although it has been enormously used since its introduction into Europe in 1672, its physiological action is not as yet well made out. That the active principle is absorbed, and that the vomiting is so produced, is shown by the experiments of Orfila (*Toxicologie*, i. 651) and of Dr. Dyce Duckworth (*St. Bartholomew's Hospital Reports*, vol. v.), both of whom found that vomiting followed the hypodermic use of emetia in the dog.

There is an enormous mass of clinical testimony to show that it has a decided tendency to act upon the bronchial mucous membrane and is therefore an "expectorant." Whether given in large or small doses, it is very apt to act upon the bowels, to increase and to modify their secretions. It probably influences the liver, since Pecholier (*Gazette Médicale*, 1862) affirms that in animals killed by it no hepatic glucose can be found. Moreover, great advantage from its use may often be obtained in the condition known as "biliousness." In "bilious dysentery" it will often produce large tarry discharges; and I have seen a change in the color of the stools follow its use in catarrhal jaundice. The mechanical effect of the vomiting induced by it in these cases, however, must not be lost sight of; yet it does not seem to me at all sufficient to account for the results, especially as some observers state that the effects noted are produced even when little or no vomiting occurs.

Upon dogs ipecacuanha acts as upon man, the impure emetia producing vomiting and sometimes purging. If the dose be sufficiently large, intense prostration sometimes comes on and ends in death.

Pecholier* found that in the rabbit, which for anatomical reasons cannot vomit, emetia given in lethal doses produces diminution in the number and energy of the heart's beats, the pulse sinking from 160 to 120, lessening of the respiration, and fall of the temperature in the mouth and ear from 1° to 3° C., that of the rectum remaining stationary, or even rising 0.5° C.

Dr. Dyce Duckworth failed, however, to find that emetia influences materially the circulation, at least until very late in the poisoning. The pulse-rate was not constantly affected; sometimes it was apparently lowered, sometimes it remained about the same, and sometimes it was seemingly increased. In two experiments made with the kymographion upon dogs, the arterial pressure remained as it was before poisoning for a long time. In one experiment, a fresh injection of emetia into a jugular vein was followed by an immediate suspension of cardiac action; in the other dog, similar phenomena were witnessed, —a half-grain was injected, and suddenly the pulse fell in rate and became irregular; in a minute and a half the arterial pressure descended from 135 to 20, and in a moment the animal was dead of cardiac paralysis. In neither of these experiments was there any vomiting.

It is evident that with the insufficient and contradictory evidence at our disposal we can only wait for more light, and perhaps draw the conclusion that in therapeutic doses ipecacuanha has no marked direct action upon the circulation.

Dr. Dyce Duckworth is also in disagreement with Pecholier in regard to the action of the drug upon the respiration, since he found that the respiratory acts were increased in frequency.

The post-mortem results obtained in animals poisoned with ipecacuanha by these authorities were even more contradictory. The former observer found great paleness of the lungs, with intense hyperæmia of the stomach and the upper half of the intestines; the latter, intense hyperæmia of the lungs, which were in some places emphysematous, but in other portions collapsed and even affected with true consolidation. The lesions were much less marked in the intestines than in the lungs, which resembled very closely those taken from the bodies of animals killed by section of the vagi. The pulmonic lesions were found to be most intense in the rabbit; the intestinal, in the dog, cat, and guinea-pig. After section of the cervical vagi, Dr. Dyce Duckworth found that emetia failed to cause vomiting.

The anatomical lesions found by the English investigator are the same as those noted by Magendie forty years ago, and are in all probability correct. It is evident that until further studies are made it is impossible to frame any accurate theory as to the action of ipecacuanha.

* Dr. Dyce Duckworth (*St. Bartholomew's Hospital Reports*, vol. v. and vol. vii.) has experimented extensively with emetia, and, so far as the action of the drug on animal heat is concerned, corroborates these results of Pecholier, since he found that, whilst the surface of the body was often remarkably cooled, the temperature in the internal parts was about normal.

THERAPEUTICS.—The most ordinary use of ipecacuanha is as an emetic. Whenever it is desired to unload the stomach or to act by emesis upon disease, without inducing much prostration, this drug commends itself by its safety and efficiency. In *narcotic poisoning* it is less certain than the “mineral emetics,” but, as it produces no irritation of the stomach, can be given more freely than they can, and is constantly used as an adjuvant to them. It is especially useful in the diseases of children, never causing the serious depression which tartar emetic is so apt to produce. When, however, very violent emesis is desired, as in *membranous croup*, other emetics, such as zinc or alum, are to be preferred, on account of the greater force of their action.

In *sick stomach* of nervous origin, as in *pregnancy*, minute doses of ipecacuanha have so often met with success that there can be no doubt of their value. One drop of the wine in a teaspoonful of water should be given every hour.

The use of ipecacuanha as an expectorant will be spoken of under that heading.

One of the most important uses of ipecacuanha is in *acute dysentery*,—all forms of which have been treated with it with asserted advantage. I think, however, its beneficial action is best seen in “*bilious dysentery*” and in “*malignant dysentery*,” as is indicated by the fact that its use is most common in tropical climates. In “*sthenic inflammatory dysentery*” it seems to be less available; although even in this it has its advocates.

In *catarrhal jaundice*, in *intermittents* or *remittents* accompanied by congestion of the portal circulation, ipecacuanha is often very serviceable.

As a *hæmostatic*, ipecacuanha has been recommended by Trousseau and other authorities, but as such it is at present very seldom used; at least I have never known of its having been so employed. It has been even given, with asserted advantage, to arrest *flooding* after child-birth.

ADMINISTRATION.—As an emetic, ipecacuanha is generally administered in powder, thirty grains being given every fifteen or twenty minutes until the desired effect is produced. For a child a year old the emetic dose is five grains. Its action should be aided and hastened by large draughts of lukewarm water. As a nauseant, the dose is from two to five grains.

In dysentery it is generally best to begin with a full emetic dose, or with ten grains repeated every half-hour until emesis is produced. Two or three hours after vomiting, fifteen drops of laudanum should be exhibited, followed in twenty minutes by five grains of ipecacuanha in *pill-form*: this should be repeated every two or three hours, the amount of the opium being lessened and that of the ipecacuanha increased, according to circumstances. The object is to have as much of the ipecacuanha retained as possible. Another plan is to give larger doses (twenty grains), repeated every two, four, or six hours, mustard being applied to the epigastrium; and it is said that after two or three doses tolerance is established and the drug retained. In India, enemata of ipecacuanha are often employed, either as a substitute for or a

succedaneum to its use by the mouth. A scruple of the powder may be given with starch and laudanum, and repeated every four hours.

As a counter-irritant, ipecacuanha is rarely used in this country; but in England a liniment is employed composed of four parts of the powder to fourteen parts of olive oil.

The following preparations are for internal use:

Syrupus Ipecacuanhæ—*Syrup of Ipecacuanha* (℞i to Oj). Dose, as an emetic for an infant, one or two teaspoonfuls.

Vinum Ipecacuanhæ—*Wine of Ipecacuanha* (Fld. ext. ℞ii; Sherry, ℞xxx). Dose, the same as the syrup.

Extractum Ipecacuanhæ Fluidum—*Fluid Extract of Ipecacuanha*. Dose, as an emetic, for an adult thirty drops.

Emetia is not officinal, but has been used by Dr. Dyce Duckworth (*London Pharmaceutical Journal*, March, 1872) and his colleagues in doses of from one-twelfth to one-sixth of a grain.

SANGUINARIA—BLOODROOT.

This is the rhizome of an indigenous perennial herb, *Sanguinaria Canadensis*. It occurs in pieces two or three inches long, reddish brown externally, a bright somewhat orange red internally, and, when fresh, full of a similarly-colored juice. It contains an alkaloid, first discovered in it by Dr. Dana, and named by him *Sanguinarina*, but which is identical with the alkaloid of *Chelidonium majus*—*Chelerythrin*. As it was discovered by Dana in 1829, and the latter by Probst in 1839, it is obvious that the name *sanguinarina* has the priority and should be adopted.

Sanguinarina occurs in colorless stars or groups of fine needle-like crystals, tasteless when dry, of a sharp burning taste when moistened with alcohol, very irritating to the nasal mucous membrane, and uniting with acids to form brilliant red, mostly soluble, salts. Two other alkaloids, *Porphyræxin* and *Puccin*, are said to have been found in bloodroot.

PHYSIOLOGICAL ACTION.—Although *sanguinaria* has been more or less used for so many years, we are still without any really definite account of its action. Little or nothing has been added to our knowledge since the paper of Dr. Tully, in 1830. That observer claims that when given in small repeated doses it acts as a very decided cholagogue; and more recently it has been affirmed that it is also a stimulating expectorant. In full doses it is certainly a harsh emetic, and in overdoses, according to Tully, it produces, with the vomiting, burning at the stomach, faintness, vertigo, diminished vision, general insensibility, coldness, extreme reduction of the force and the frequency of the pulse, together with great irregularity of action and often palpitation of the heart, great prostration of muscular strength, and sometimes a convulsive rigidity of the limbs.

Fatal poisoning of several persons by it occurred at Bellevue Hospital; but the only symptoms recorded are "racking, burning pains, and tormenting thirst."

THERAPEUTICS.—As an emetic, sanguinaria has fallen into well-deserved disuse. Indeed, I have never known of its employment except as a stimulant expectorant in obstinate *bronchitis*, and even then with doubtful advantage.

ADMINISTRATION.—The crude drug is very rarely used; the emetic dose of the powder is from ten to sixty grains.

Prof. R. P. Thomas, in experiments upon himself and others with the alkaloid, found that in a dose of from one-eighth to one-twelfth of a grain it acted as an expectorant, without disturbing the stomach; one-sixth or one-fourth of a grain, given every two or three hours, generally nauseated, the emetic dose being half a grain repeated every ten minutes. One-sixth of a grain every three hours, in the course of two or three days, reduced the pulse from five to twenty-five beats per minute.

Tinctura Sanguinariæ.—*Tincture of Sanguinaria* (3ii to Oj). Dose, as an expectorant, gtt. xx to xl.

APOMORPHIA.—This non-officinal alkaloid was discovered by Dr. Matthieson and C. R. A. Wright (*Proceedings of the Royal Society*, xvii. p. 455), who made it by the action of a strong solution of hydrochloric acid upon morphia. A probably better method of preparation is that of E. L. Mayer (*Berichte der Deutschen Chem. Gesellschaft zu Berlin*, 1871, iv. p. 121), in which the morphia is treated with a solution of the chloride of zinc at 120° C.

Apomorphia occurs as a snow-white powder, which is permanent when dry, but when moist soon becomes green. Its solution suffers this change, which is probably an oxidation, in a few hours, and in the course of some weeks the green tint deepens into a black. Bichromate of potash turns it a dense yellow orange; bichromate of potash and concentrated sulphuric acid make a dark red with it, and with neutral chloride of iron it strikes an amethyst color; heating the solution causes the green color to appear in a short time (Matthieson and Wright). It differs from morphia in being soluble in cold water.

PHYSIOLOGICAL ACTION.—Dr. Gee (*St. Bartholomew's Hospital Reports*, vol. v. p. 215) announced that apomorphia is a certain and prompt emetic, having the great advantage of acting in very small dose, a tenth of a grain being sufficient, when injected under the skin, to cause vomiting in ten minutes. His assertions have been confirmed by various observers, among whom may be mentioned Dr. F. M. Pierce (*Brit. Med. Jour.*, 1870, p. 274) and Drs. Franz Riegel and Rudolf Boehm (*Glasgow Med. Jour.*, May, 1872).

According to Dr. Pierce, the original dose employed by Dr. Gee is too large, one-fifteenth of a grain sufficing in almost all cases to cause vomiting in from five to twelve minutes.

Apomorphia certainly has other influences upon the system than those of an emetic; but we have no definite knowledge as to their exact nature. Dr. Gee states that sometimes a contra-stimulant effect is very marked, and the patient suffers even from alarming muscular weakness; and, according to Riegel and Boehm, headache, giddiness, and a disposition to yawn are often present, and free perspiration and great drowsiness, with apathy, are quite

common. According to the latter observers, in none of their experiments was the pupil affected; the pulse was very generally diminished in force and volume.

THERAPEUTICS.—Apomorphia bids fair to be a very useful emetic, but neither our clinical nor our physiological knowledge of it is as yet sufficient to establish its exact value. Its great promptness of action, its ease of administration, and the freedom of its emesis from nausea, seemingly fit it for all cases in which evacuation of the stomach is desired. The effect of narcotics upon its action is not determined; but Dr. Loch has reported (*Schmidt's Jahrbücher*, Bd. clv. p. 272) a case of poisoning by oil of bitter almonds in which the effects of a hypodermic injection, 0.013 grain, were most happy. The same observer has, on the other hand, noted a case of acute gastric catarrh in which the alkaloid did not cause vomiting, although exhibited until the patient was unable to stand. He has also used it in a child thirteen months old suffering from capillary bronchitis: 0.002 grain was given, and caused free vomiting, which left the child exceedingly exhausted.

I am not aware that abscesses have been reported as produced by the hypodermic injection of the alkaloid, but the pain caused is said to be in some cases very severe. If apomorphia be given by the stomach, about double the hypodermic dose should be exhibited. Owing to the great rapidity with which apomorphia undergoes decomposition when in solution, it should always be dissolved at the time of using. Dr. Loch reports a case in which most alarming symptoms were caused by the use of an old solution.

In the secondary list of the U. S. Pharmacopœia are included the roots of *Gillenia stipulacea* and *G. trifoliata*; also *Euphorbia corollata* and *E. ipecacuanha*, all indigenous plants. The former of these are conjointly recognized as *Gillenia*, which is said to be a mild emetic, resembling ipecacuanha in its action, but far less certain: the latter bear the names of the plants that yield them, and are very harsh emetics, capable in overdoses of acting as acrid poisons. I have never known of either of these drugs being exhibited.

A very useful stimulating emetic is *mustard flour*, very prompt and even violent in its action. It acts as a mechanical emetic, and is to be used when it is desired simply to evacuate the stomach rapidly, and especially when there is torpor of the viscus. As it is very generally to be had at once, it is especially useful in such emergencies as *narcotic poisoning*. It has also been commended in *nervous collapse*, such as is seen in *malarial pernicious chill*. The dose is a heaped dessertspoonful in half a pint of water, repeated, if necessary, in ten minutes. As mustard is irritant to the stomach, if it fail to act it should not be repeated more than three or four times, even in narcotic poisoning.

Squill (Scilla) is sometimes used as a harsh, stimulating mechanical emetic.

MINERAL EMETICS.

The mineral emetics in ordinary use are tartar emetic, sulphate of zinc, and sulphate of copper. As they are all considered elsewhere, it is only necessary here to say a few words as to their use as emetics.

Tartar emetic is the most depressing of all the substances here spoken of as emetics. It is rather slow in its action, but the vomiting which it causes is preceded and accompanied by intense nausea, and is exceedingly violent and persistent. For these reasons, tartar emetic is rarely used simply to unload the stomach, except in the absence of more eligible substances. Its use in inflammatory diseases is discussed elsewhere. It is a centric emetic acting by absorption.

Sulphate of zinc is an excellent and prompt mechanical emetic, producing little or no irritation, and is to be preferred above all others when an emetic of such nature is needed. In *narcotic poisoning* it should be given in combination with ipecacuanha, and perhaps be preceded by mustard whilst it is being obtained from the apothecary. Thirty grains of it with fifty of ipecacuanha may be given as the first dose, and a mixture of fifteen grains of the former to thirty grains of the latter be administered every fifteen minutes until the desired effect is produced or one hundred grains of the zinc are taken. Beyond the latter amount it would be hardly safe to go, for fear of causing gastro-enteritis.

The *sulphate of copper* resembles the corresponding zinc salt as an emetic, but is more severe and irritating, and more capable of causing gastro-enteritis. The full dose in narcotic poisoning is from five to ten grains, which should not be repeated more than once.

Powdered *alum* is a mechanical emetic, which has been especially recommended in *membranous croup*, on account of its being believed to act beneficially upon the diseased surfaces in its passage up and down the throat. A heaped teaspoonful of it may be given in molasses or syrup. In my experience alum has proved an unreliable emetic.

CLASS II.—CATHARTICS.

PURGATIVES, or cathartics, are those drugs which are employed in medicine to act upon the intestinal tract so as to produce purgation, or catharsis. The question has recently been a good deal discussed as to whether purgation is produced by an increase of the watery secretions, or of the peristaltic movements of the intestines.

M. Thiry (*Sitzungsberichte der k. k. Akad. d. Wissensch.*, Bd. I.) experimented upon this subject by drawing out a knuckle of intestine through a wound in the linea alba, cutting it free from the remainder of the gut without injuring its nerves or blood-vessels, sewing together the distal and proximal ends of the main portions of the intestines so as to re-form a continuous tube, and then, after closing up one end of the knuckle, forcing the other into the wound so as to make an intestinal *cul de sac*, which could be studied through a fistulous opening. In dogs which had recovered after this operation, Thiry found that large doses of the sulphate of magnesia, of senna, or of croton oil failed alike to increase the secretion of the separated piece of intestine, although they induced violent purging; further, neither concentrated solutions of Epsom salts, nor infusion of senna, even though kept in the *cul de sac* for some time, were able to increase its secretion by exosmose. More recently, Dr. S. Radziejewski has made a most elaborate investigation of the subject (*Reichert's Archiv für Anatomie*, 1870, p. 37). As a result of a number of very careful analyses, he asserts that there is nothing to be found in the stools produced by the sulphate of magnesia, by calomel, castor oil, croton oil, senna, or gamboge, to indicate that they are anything besides the ordinary contents of the upper and lower bowels. Dr. Radziejewski confirms the fact observed by C. Schmidt, that the stools of purgatives contain a great deal of soda, but denies that this proves at all that they are transudations, asserting that the alkaline salts are derived simply from the pancreatic fluid. Dr. Radziejewski also corroborates the confirmation by Asp (*Ludwig's Arbeiten*, 1868) of the discovery of Moreau,* that division of the intestinal nerves is followed by free serous exudation into the gut, but denies that purgatives act by paralyzing the vasomotor nerves, because croton oil injected into a loop of intestine which had been separated by two ligatures from the remainder of the gut, caused both

* *Comptes-Rendus*, t. lxxi., 1868; also Asp, *Ludwig's Arbeiten*, 1868.

vomiting and purging. As no emulsifying substance was contained in the intestine, he declares that no absorption could have occurred, and that consequently the general intestinal disturbance was simply due to increased peristaltic action, caused by the internal local irritation of the oil propagated along the intestines.

The experiments of Thiry have also been repeated by Radziejewski with croton oil and with sulphate of magnesia, as well as by Schiff (*Nuove Ricerche sul Potere digerente*, II. Morgagni, 1867) with aloes, jalap, and sulphate of soda. In all cases the results were the same as those already noted as obtained by Thiry.

Carrying his investigations still further, Radziejewski, by forming intestinal fistule at such positions as would enable him to study the rate of passage of the intestinal contents, found that after a dog is fed upon flesh the small intestine empties the partially-digested food into the colon so rapidly, and in such quantity, as to constitute, so to speak, a normal diarrhœa, and that the long delay in the exit and the hardening of the fœces occur in the large intestine. The liquid which passed into the ascending colon agreed in all its characteristics with the stools of purgation.

Dr. Radziejewski also claims to have established by direct experimentation that the peristaltic movements of the small intestine were affected very decidedly by drastics, and to some degree by Epsom salts, and that in all cases the large intestine was still more intensely acted upon.

Although these experiments are very interesting, it cannot be allowed that they prove what is claimed for them, namely, that purgatives cause no increase of intestinal secretion, but only of peristaltic action. So much violence to natural conditions is done in the experiments after the method of Thiry that they seem worthy of very little weight. The assumption of Radziejewski, that croton oil confined in a loop of intestine is not absorbed, is a pure assumption, and his experiment does not warrant the conclusions drawn from it.

The other facts brought forward seem to prove only that increased peristalsis, especially of the large bowel, plays a more important rôle in the induction of diarrhœa than has been assigned to it.

Leaving out of sight for the moment all clinical evidence, the fact that previous section of the par vagum prevents the action of purgatives* is opposed to the German theory, since it is almost certain that the division of the nerves of the neck does not arrest peristaltic movements. Further, Armand Moreau (*Archives Générales*, 6e sér., t. xvi. p. 234) has found that a solution of Epsom salts, placed in a knuckle of intestine isolated by means of two ligatures, does cause a serous exudation into it, and in repeating M. Thiry's experiments (*Gazette Médicale*, 1871) he has obtained opposite results. His experiments indicate three possible sources of fallacy in the work of the previous investigators: first, if the Epsom salts be not kept in the intestine for a

* See paper by the author, *American Journal of the Medical Sciences*, vol. lx., 1870.

sufficient length of time (some hours), no transudation occurs; second, in some cases the inner end of the isolated piece of intestine fails to adhere, so that the opening is not obliterated, and the matters injected into the arrested *cul de sac* really pass into the peritoneal cavity; third, atrophy of the mucous membrane and glandular apparatus of the *cul de sac* often follows almost at once upon the operation, and of course necessitates a negative result in the subsequent experiments. Recently, Dr. Lauder Brunton, in a communication to the Medical Society of London, stated that he had repeated Moreau's experiments, and found that sulphate of magnesia injected into the intestine of a cat caused about two-thirds of a drachm of fluid to be secreted in four hours by each inch of the bowel operated on, although the proportion of sulphate was only one grain to an inch. (*Med. Press and Circular*, Dec. 31, 1873.)

The experimental evidence bearing upon the question under discussion is, so far as I know, all included in the above summary, and is far from demonstrating that increased peristalsis is the only cause of the watery stools produced by purgatives. The clinical evidence is overwhelmingly in favor of increased secretion. The facts proven by clinical observations and by experiment—that various purgatives act when taken into the blood, and that in these cases elimination by the bowels occurs; that at least some purgatives (Headland, *Action of Medicines*, London, 1867, p. 443), when given by the mouth, are absorbed, disappearing from the alimentary canal and reappearing when purgation occurs; that the stools induced by overdoses of various drastics, as elaterium, are so enormous as to cause the profoundest depression, and even choleraic collapse, in a very few hours; that the discharge of hydragogues contains a very large percentage of soda in the alkali of the serum; that the relief obtained in portal congestion by the depletion of salines is very marked—are, when viewed together, to my mind, incompatible with any other belief than that purgatives cause increased secretion, as well as increased peristalsis, in the alimentary canal.

The question of the action of drugs upon the flow of bile is a very important one, the evidence concerning which is best considered under two headings: first, the experimental; second, the clinical.

The experiments upon this subject which have attracted most attention are those of Dr. Scott, and of the Edinburgh committee, of which Prof. Bennett was chairman and Drs. Rutherford and Gangee the workers. The method employed both by Dr. Scott and by the Edinburgh committee was to make biliary fistulæ in dogs in the usual physiological method, and, after recovery from the operation had taken place and the bile regularly escaped by the external orifice, to administer the drugs, especially calomel and podophyllin, and study the effects upon the excretion of bile. Of the accuracy of their experiments I do not think there can be any reasonable doubt. I believe they prove that in dogs with biliary fistulæ mercury has no effect upon the flow of bile unless given in such quantities as to deteriorate the general health, when it diminishes the biliary secretion. This result does not, how-

ever, warrant the further conclusion that mercury does not increase the flow of bile in healthy dogs. The animals were in such an unnatural condition that, in spite of the daily ingestion of much more than the normal amount of food, they progressively emaciated, and finally died apparently of inanition: moreover, the innervation, and very possibly the blood-supply also, of the liver was very much interfered with. Under these circumstances it is clearly conceivable that the mercurial or other purgative might in the uninjured dog affect the biliary secretion, and yet fail to do so in the experiment, hindered by some obscure yet efficient cause.

Very recently, Dr. A. Röhrig has experimented (*Stricker's Medicinische Jahrbücher*, 1873) in a method which simulates more closely the natural conditions; although even the results thus obtained do not seem to me entirely conclusive. In curarized dogs in which life was maintained by artificial respiration, he placed a glass tube in the gall-duct so that the bile could escape only through it. Under these circumstances, of course, after a time secretion ceased, and Dr. Röhrig experimented not only on the effect of remedies upon the secretion whilst naturally going on, but also upon their power of re-establishing it. He found that large doses of croton oil (eighteen drops) thrown into the duodenum caused an immediate very great increase, or a re-establishment, of the secretion. After the oil, the vegetable cathartics were most active, decreasing in power in the following order: colocynth, jalap and aloes, rhubarb and senna. Castor oil had very little influence, as had also the bitter salts. Calomel, even in large doses (twenty grains), very rarely re-established the secretion, but its power of increasing and maintaining it beyond the natural time for cessation was very marked.

What is to be drawn from these various facts? Evidently, I think, but one conclusion,—that the experimental evidence at present does not warrant any deductions as to the effect of purgatives upon the biliary secretion of healthy dogs. The canine diet and digestion are so different from the human, that it is to be expected that medicines acting upon the digestive apparatus will influence dogs differently from man; and I have given doses of elaterium that would have killed a man to some of the carnivora without causing the slightest purging. In view of these facts, the only fairly deducible conclusion in regard to the experimental evidence that has been brought forward is, that it must be all laid aside when we desire to study the question as to the cholagogue action of remedies upon man, and that our conclusions must be based solely upon clinical evidence.

In regard to the drastics, there can be little doubt that almost any irritant purgative will to a greater or less extent increase the escape of bile, probably both by increasing its flow into the duodenum and by sweeping it out of the small intestine before absorption can take place. There are, however, two actively purgative substances for which it is especially claimed that they are cholagogues: namely, calomel and podophyllin. The discussion of the action of these will be found under their respective headings.

Various divisions of purgative medicines have been proposed by different authors ; but probably the most convenient arrangement is as follows :

1. *Laxatives*.—Medicines which simply unload the bowels, and are not able to cause active purgation, even when given in very large doses.

2. *Purges*.—Medicines which purge actively, but are not capable of acting as poisons, even in very large amount.

3. *Hydragogues*, which produce very large watery stools without much irritation. In overdoses, medicines of this class assume some of the characters of those of the next.

4. *Drastics*, which cause great irritation of the alimentary mucous membrane, and in overdoses are violent poisons.

It must be borne in mind that this classification is somewhat artificial ; that the effects of the remedies depend much upon the doses in which they are administered, so that in sufficiently minute quantity a drastic may act as a laxative ; and that the dividing-lines between the groups are not very distinct.

The indications to fulfil which cathartics are used are as follows :

1. *To unload the bowels*.—It is not necessary, in a work like the present, to say anything about the evil results of retained fecal matter, but only to point out the methods of relief. Before this can be done to advantage, however, a summary of the causes of *constipation* is required. Constipation may be well divided into acute and chronic. *Acute or temporary constipation* is that which occurs under special, transient circumstances, as in convalescence from acute disease, and in pregnancy. It is to be relieved by the use of laxative articles of diet, and, this not sufficing, by laxatives or purgative medicines. It should never be forgotten that acute constipation is sometimes due to organic affections of the alimentary canal, such as enteritis or intussusception, or is caused by a mechanical obstacle, such as a hard foreign body or an enormous gall-stone. It is evident that such cases are not simple constipation ; that the treatment required is essentially different from that of the latter affection, and is various according to the lesion. For the diagnosis and treatment of these diseases the reader is referred to works on the practice of medicine. *Chronic constipation* may be due to sedentary habits of life ; to habitual overwork, especially of the nervous system ; to a deficiency of intestinal secretion and of peristalsis, apparently natural to the individual and without obvious cause ; to long-continued voluntary habit of restraining the desire to go to stool ; to lead- or other forms of poisoning ; and to diseases of the nervous system producing a paralytic state of the intestinal muscular fibres. It is evident that in the treatment of these various forms of constipation due regard must be paid to the cause, which should always, if possible, be removed. There are also certain cardinal principles which apply to the treatment of all forms of chronic constipation. They are as follows :

1. A voluntary effort at defecation is to be daily made at a fixed hour, whether the desire exist or not.

2. Medicines are to be avoided as far as possible, a sustained effort being made to regulate the bowels by means of diet.

3. In very many cases the daily use of enemata of cold water, with attention to diet, suffices to attain the desired result.

4. If medicines become necessary, as small an amount as will suffice, and the mildest drugs, are to be used. Purgatives or laxatives are at best merely temporary devices, and if abused in costiveness increase the trouble. So far as can be, the attempt should be to produce a permanent impression, an alteration of the intestinal glandular action or peristalsis. Thus, when atony of the muscular coat exists, strychnia, or, according to recent experiments and clinical observations, Calabar bean (see page 275), may be employed; if the hepatic or other glands are habitually torpid, nitro-muriatic acid may be administered.

When constipation is attended with low spirits and a coated tongue, it is almost always due to a deficiency of secretion, and may be looked upon as a form of dyspepsia: in such cases nitro-muriatic acid is especially valuable, but sometimes a mild mercurial course seems almost imperative.

A second use of cathartics under the present indication is to remove offending materials, as indigestible or irritant food, foreign bodies, acrid discharges, etc. For these purposes a brisk, quickly-acting purgative is generally best.

2. *To deplete.*—On account of the large serous flow which they produce, the hydragogue cathartics when freely exhibited cause a very decided general depletion.

Local depletion by means of cathartics is called for in *congestion* of the *portal circulation*, as well as in *dysentery* and other acute intestinal inflammations. Under the first of these conditions may, we think, be included without violence cases of the so-called "torpidity of the liver," which will be discussed in the article upon calomel. In acute *intestinal inflammations* the salines are to be preferred when depletion is desired, as they produce very large serous discharges and are not at all irritant.

3. *To promote absorption.*—By emptying the blood-vessels the cathartics favor the absorption of the exuded fluid in general dropsy. For this purpose the hydragogues, and especially elaterium, are the best purgatives. The production of catharsis is the surest method of relief in general *dropsy*, also in *ascites*; in other forms of local effusion its effects are less marked. As, however, purgation is the most exhausting of all the plans employed for the cure of dropsy, due regard must always be had to the strength of the patient. It is frequently necessary actively to support or even to stimulate while it is being carried out.

4. *To revulse.*—The long tract of the alimentary canal affords a great extent of surface upon which to practice revulsion in certain brain-diseases, as in *mania* and rheumatic or gouty *irritation* of the *cerebrum*. In *hyperæmia* of the brain, purgatives do good by depleting as well as by acting as revulsives. The drastics should be preferred.

5. *To eliminate*.—It cannot be doubted that the use of purgatives in such diseases as fevers and cholera, with the idea of eliminating some *materies morbi*, rests simply upon a crude, unproven, and probably false pathology. In rheumatic disease and in gout it is more probable that they do good in this way, although it is by no means certain that the advantage derived from their use is not simply due to depletion. In cases of retained renal secretion, the evidence is very decided that they do aid in separating the products of retro-grade metamorphosis from the blood.

6. *To influence the pelvic circulation*.—The only purgative used for this purpose is aloes, in the article upon which all that is necessary will be said upon the subject.

LAXATIVES.

As has been already stated, constipation should always, when possible, be overcome by laxative food. There are two qualities by virtue of which food is laxative. Chief of these is *bulk*. All aliment which contains a large amount of innutritious material affords a large residuum, which, by distending the intestine, stimulates peristalsis. Contrariwise, articles of diet which are highly nutritious and afford but little residuum are constipating. This holds good, more or less strictly, among the lower animals. Thus, the flesh-eating carnivora are habitually constipated; the grass-eating herbivora very generally lax. Owing to its containing so little of the innutritious portion of the grain, the finest white flour favors a costive habit, whilst the “cracked wheat,” in which the whole grain is eaten, is laxative; as to a still greater degree is bran, which is composed almost wholly of the husk of the wheat, the least nutritious portion of it, and therefore leaves a large residuum after digestion. Cracked wheat is boiled into a sort of jelly-like mass, and eaten with cream and sugar, whilst bran is taken in the form of bran bread, bran crackers, or bran mush. Unbolted flour, containing the whole of the grain, is about equal to cracked wheat, and is often made into bread. Indian meal, in the form of cakes or of mush, is highly nutritious, and somewhat laxative; oatmeal is decidedly laxative, scarcely so much so as bran, but much more nutritious. When it agrees with the stomach, and is easily digested, it is probably the best of all these laxative articles of food. As the oats produced in warmish climates are very inferior, care should be taken to procure oatmeal manufactured from Northern grain. It should be thoroughly cooked, and is best eaten in the form of a thick porridge. In dyspepsia all of these articles sometimes disagree with the stomach and cannot be used.

Some dietary articles seemingly possess *dynamic* laxative powers,—i.e., they exert a direct action which is not mechanical, but is similar to, although far less active than, that of the true purgatives. They intensify the intestinal action. Chief among substances of this class are molasses, and its congener, brown sugar; white sugar probably does not share these laxative powers. Of course, great care is usually necessary in taking advantage of this laxative

virtue of molasses, on account of the danger of producing fermentation and acidity in the *primæ viæ*. An obvious deduction, however, is to encourage the use of brown instead of white sugar in those of constipated habit.

There are certain foods which combine the two methods of action spoken of. Chief among these are the fresh acidulous fruits—such as apples, pears, etc.—and the dried fruits. Of the latter, the fig is one of the most palatable, and, owing probably to the great number of small seeds which it contains, is the most efficient. Prunes are nearly as agreeable as figs. To a limited extent the finest varieties of them may be eaten raw; but they are especially to be recommended stewed. When it is necessary, a pinch of senna-leaves may be cooked with them, and, if it be not made too large, increases the activity of the dessert without affecting its flavor.

Among constipating articles of diet, it is only necessary to call attention to milk as one of the most decided of the class.

The laxative remedies of the U. S. Pharmacopœia are as follows:

TAMARINDUS—TAMARINDS.

The preserved fruit of the *Tamarindus Indica*, a large tree, native of the East and West Indies. The fruit is a broad, compressed pod, usually from four to six inches long, somewhat curved, with an exterior brown hard rind. It contains seeds inclosed in cells formed of a tough membrane, between which and the rind is an acid pulp, the medicinal part of the fruit. Tamarinds are preserved for market by stripping off the outer rind, packing the inner portion in layers, and pouring on boiling syrup. In the market they are offered as adhesive masses composed of pulp, membranes, strings, and seeds, and having a sweet, acidulous taste. They contain a good deal of citric acid, much less of the tartaric, and a little of the malic acid. They are used chiefly in making refrigerant acidulous drinks for fever, and in convalescence as a laxative article of diet, half an ounce to an ounce or more being eaten like preserves. They enter into the officinal confection of senna.

MANNA—MANNA.

An exudation of the European ash, *Fraxinus Ornus*, chiefly produced in Sicily and Calabria. There are three varieties of it. The best, *flake manna*, occurs in unequal, rough, stalactite-like pieces with a crystalline or granular fracture, and is obtained in the hottest and dryest weather in July and August. The next quality, *manna in sorts*, consists of pieces of flake manna, mixed with a soft brownish matter; it is obtained in September. *Fat manna*, a soft viscous mass, which exudes during the wet weather in the latter part of October and in November, is the least valuable variety. Manna has a slight odor, a sweet, mawkish taste, and should contain from forty to eighty per cent. of the saccharine, active, crystalline principle *Mannite*, which differs from ordinary sugar in not containing equal parts of hydrogen and oxygen, and is therefore not readily convertible into grape sugar or its derivative, alcohol.

THERAPEUTICS.—Manna is a gentle laxative in large doses, sometimes causing flatulence and pain. It is rarely used by itself, but is added to infusions of more powerful purgatives, to cover their taste and aid in their effects. The laxative dose for an adult is half an ounce to two ounces; for a child, one to four drachms in an aromatic infusion.

CASSIA FISTULA.—*Purging Cassia* is the pulp of a hard, blackish, cylindrical pod from one to two feet in length and about an inch in diameter, having on one side a single and on the other a double dark band, running the whole length of the pod, and marking the positions where its valves are united. The pods are produced by a large tree, *Cassia fistula*, a native of Egypt and India. The dark, sweetish, acidulous, officinal pulp may be used as a laxative in doses of half an ounce, but is apt to cause griping. It enters into the confection of senna.

MAGNESIA.

The U. S. Pharmacopœia contains in its primary list the *carbonate of magnesium*, and among its preparations *magnesia*, which is made by heating the carbonate to a red heat in an earthen vessel. In commerce there are two varieties of both substances, which are known as the *heavy* and the *light*, and differ only in their physical characters, the particles being differently aggregated. The carbonate of magnesium is manufactured by precipitating a solution of the sulphate of magnesium by one of the carbonate of sodium. If the two solutions be concentrated, the dense or heavy carbonate falls; on the other hand, if the solutions be dilute, the precipitate is a light carbonate. Heavy *magnesia* is obtained by calcining a heavy carbonate; light *magnesia*, by using a light carbonate.

Both of these substances are of a milk-white color, and occur in powder; the carbonate sometimes in very light cubical blocks. They are both practically insoluble in water, freely so in dilute acid, and in the presence of acids they both act as alkalies.

THERAPEUTICS.—*Magnesia* and its carbonate act in the same manner upon the human economy, being both antacid and laxative. For their purgative powers they are probably dependent upon the presence of acids in the primæ viæ, and hence their effects vary. They are sometimes taken as habitual laxatives by persons suffering from *acid dyspepsia*; but, as they are said at times to accumulate in the intestines and to do harm mechanically, this use of them should be discountenanced. They are very frequently employed in conjunction with Epsom salts, senna, or other of the more powerful purgatives, on account of their antacid properties.

Their chief use is in acute *acid dyspepsia*, in *sick headache*, in some forms of *diarrhœa* with excessive acidity in children, in *gout*, *rheumatism*, and in various *cutaneous affections*; wherever, in a word, a laxative antacid is indicated.

ADMINISTRATION.—The dose of the carbonate is, for a child a year old,

from five to twenty-five grains, according to the effect desired; for an adult, half a drachm to half an ounce; that of the magnesia is about one-fifth less.

SULPHUR. (S.)

Sulphur is official in three forms: SULPHUR SUBLIMATUM, or *Flowers of Sulphur*; SULPHUR LOTUM, or *Washed Sulphur*; and SULPHUR PRÆCIPITATUM, or *Precipitated Sulphur*. The first of these is made by subliming sulphur into cool chambers, and always contains some sulphuric acid, generated during the process. When freed from the acid by washing with warm water, it constitutes the washed sulphur. The U. S. Pharmacopœia directs the precipitated sulphur to be prepared by boiling lime and sulphur together, so as to form a sulphide of calcium, and precipitating this with muriatic acid.

The sublimed and the washed sulphur occur as sulphur-yellow, crystalline powders; the precipitated as a whitish powder, whose particles are often coherent into friable lumps.

Sulphur is not only a crystalloid dimorphous, but is capable, under the action of heat, of assuming various allotropic forms. For an account of these, and for its chemical properties in general, the reader is referred to works on chemistry. It is insoluble in water, but soluble in alkaline solutions, in alcohol, the fixed and volatile oils, chloroform, ether, etc.

PHYSIOLOGICAL ACTION.—When applied locally, sulphur is almost without influence. Taken internally, owing to its solubility in the alkaline juices of the intestines, it is absorbed to a greater or less extent, and has been detected in the milk, sweat, urine, and even in the breath. At the same time, if in sufficient quantity, it acts as a mild laxative, producing soft, semi-liquid, feculent stools, accompanied generally with much offensive flatus of sulphuretted hydrogen. It is affirmed that in some instances the latter gas has been so freely generated and absorbed as to cause systemic poisoning. Cases have also been reported in which the flowers of sulphur acted as an irritant poison; but this, without doubt, has been owing to their containing a large quantity of sulphuric acid. Its continued use has probably some effect upon nutrition; the secretions generally are slightly increased, and some have affirmed that the temperature is somewhat elevated; but the truth of this is certainly very doubtful. The results of clinical experience indicate that it has an especial tendency to act upon the skin and mucous membranes.

THERAPEUTICS.—As an habitual laxative, sulphur has been used with asserted advantage in cases of *hemorrhoids* and of chronic *rheumatism*. In subjects of the latter disease, it is claimed that it exerts a beneficial alterative influence, especially in *sciatica* and in *lumbago* and other varieties of *muscular rheumatism*. It has also been employed as an alterative in various cutaneous affections; and in the form of natural sulphur-waters, used externally and internally, there is much testimony as to its value in both rheumatic and skin diseases.

Sulphur is very largely used as a parasiticide in cases of *itch*. Dr. Tilbury

Fox recommends its application in the following manner. He says, "I have applied to *all papules and vesicles* the following ointment: sulphur, half a drachm; ammonio-chloride of mercury, four grains; creasote, four drops; oil of chamomile, ten drops; and an ounce of lard. This is rubbed in night and morning for three days, especially to the interdigits and wrists; the same shirt is kept on till the third day, when it is changed and a warm bath given. The use of the parasiticide for two or three days should be followed by a good washing and the discontinuance of the remedy for a night. If the patient be not troubled with itching during the night, we may conclude that the acari are killed, and all we need to do is to guard against the hatching-out of fresh acari by the light application of our parasiticide once a day to any 'pimple' or itchy place for a few days longer, taking care that the foul clothes are well heated or scalded. 'Not too strong and not too long,' is my rule in the use of remedies for scabies. The occurrence of red, rough, erythematous patches is a sign that the remedy itself is creating a disease."

ADMINISTRATION.—Sulphur is generally given in powder, mixed with syrup or molasses. Dose, as an alterative, ten to twenty grains three times a day; as a laxative, one to three drachms at bedtime.

POTASSII SULPHURETUM—*Sulphuret of Potassium* (KS). This is prepared by heating together sulphur and carbonate of potassium. It occurs in liver-brown fragments, which form an orange-yellow solution in water. Its taste is acrid, alkaline, and very disagreeable. When moistened, it feebly emits the odor of sulphuretted hydrogen.

Locally applied, the sulphuret of potassium is a very decided irritant. Taken in large quantities, it is a violent corrosive poison, and is said to have produced fatal gastro-intestinal inflammation. In medicine it is employed only externally. It has been employed as a stimulating ointment (℞ss to ℥j) in various skin-affections, and is also used for the formation of sulphur-baths, the strength of which should vary, according to the requirements of special cases, from two to six ounces of the drug in thirty gallons of water. They should be taken warm, the patient remaining in from twenty minutes to two hours, and are said to cause a general excitement, amounting in some susceptible persons to high fever. When employed strong, they sometimes occasion a papular eruption. They have been used in *chronic rheumatism* and in various *scaly skin-diseases*.

PURGES.

OLEUM RICINI—CASTOR OIL.

A fixed, nearly odorless oil, of a nauseous taste, obtained from the seeds of the *Ricinus communis* by expression. The seeds are slightly warmed before being put under pressure, so as to liquefy their contained oil; and the crude oil obtained from them is boiled with a small amount of water,

so as to coagulate its albuminous impurities. The oil was formerly manufactured by means of alcohol, also by heating the seeds or by boiling them in water, and several varieties of it existed; but these are no longer in the market. Castor oil is remarkable for being soluble not only in ether but also in alcohol. The *castor-oil seeds*, or *beans*, as they are commonly called, contain an acrid fixed principle, which makes them exceedingly poisonous, so that death from gastro-intestinal inflammation has often been produced by them.

PHYSIOLOGICAL EFFECTS.—Castor oil acts upon the human organism as a mild but decided purgative, producing copious fluid fecal discharges, and in overdoses sometimes vomiting, and always purging freely. It is a matter of some doubt whether the properties of the officinal oil are or are not due to the presence of a minute quantity of the acrid principle of the seeds; although the probabilities are in favor of the affirmative proposition. That the oil or its active principle is absorbed is proven by analogy, and by the facts that in children it sometimes purges when rubbed upon the skin of the abdomen (Canvane, *Stillé's Therapeutics*, 2d ed., vol. ii. p. 416), and that when taken into the stomach it has been known to exude from the skin (Ward's case, *London Medical Gazette*, vol. x. p. 377). In regard to its existence in the stools, the testimony is conflicting. Thus, Bucheim (*Virchow's Archiv*, Bd. xii.), although he submitted the passages produced by it to careful chemical manipulation, failed to detect it or any of its derivatives in them; but Bird (*Stillé's Therapeutics*, 2d ed., vol. ii. p. 416) and other observers affirm that it can be seen by the eye in the dejecta, either as oil or in the form of caseous flakes. According to the experiment (quoted by Stillé) of Hale upon himself, half an ounce of castor oil injected into a vein produces malaise, nausea, faintness, anxiety, and general dullness and depression, without purging.

THERAPEUTICS.—On account of the mildness of its action and an especial property which it appears to have of soothing an irritated bowel, castor oil is constantly employed whenever it is desired simply to evacuate the intestinal canal; not so much, however, in *chronic constipation* as when a temporary action is alone required.

In various inflammatory or irritative affections of the alimentary canal, castor oil is of the greatest service, partly, no doubt, by removing acrid irritating secretions or foreign materials, such as undigested food, and partly by causing a depletion of the congested vessels, but also apparently by virtue of an almost specific power, which renders it the most satisfactory cathartic in these cases. This is especially seen in the acute *diarrhæas* and even in the *chronic enteritis* of children, but also holds good in the *diarrhæas* and *dysenteries* of adults.

Within the last few years a good deal has been written in regard to the use of the leaves of the castor-oil plant as a *galactagogue*, and sufficient evidence has been brought forward to render them worthy of some confidence.

A poultice made of the fresh leaves should be applied to the breasts, and a teaspoonful of a fluid extract, prepared from the same, should be exhibited three or four times a day.

ADMINISTRATION.—Castor oil is very repulsive to the palate, so much so as to nauseate, or even vomit, by its taste, some susceptible individuals. It has been the habit to administer it in emulsion with a strong mint-water, or to give it in the froth of porter or in a cup of hot coffee; but by far the best plan is to mix it with an equal part of glycerine and to add two or three drops of the oil of cinnamon or of gaultheria to each dose. The substances do not stay mixed, but separate on standing: when used, they may be made temporarily to recombine by shaking the bottle. As glycerine has feeble laxative power, an ounce of this mixture represents a little more than half an ounce of the oil. It should be taken directly out of the spoon. The full purgative dose of the oil is half an ounce to an ounce for an adult; for an infant a year old, one to two teaspoonfuls. In dysentery it is sometimes advantageous to give the drug in small dose every three hours until a decided purgative operation is induced.

TOXICOLOGY.—Whilst castor oil itself is harmless, the castor oil beans, as already stated, contain an acrid principle, which renders them exceedingly poisonous, three of the beans having sufficed to destroy the life of a man (*Medical Times and Gazette*, May, 1861). The symptoms do not usually come on until from two to five hours after the ingestion of the poison, when severe abdominal pain is felt, and is accompanied by violent vomiting and by purging, which after a time may become bloody. This purging soon ushers in a stage of collapse, with or without severe muscular cramps, with cold sweating skin, contracted features, thirst, restlessness, small rapid pulse, and sometimes the general appearance of Asiatic cholera. After death, intense redness and even abrasion of the stomach and of the small intestines are found. The treatment should consist in the evacuation of the stomach and bowels by mild emetics, such as ipecacuanha and warm water, and by mild cathartics, such as castor oil, provided nature has not already sufficiently fulfilled the indications, and in the free use of opium and demulcent drinks, the early external application of leeches and of emollient poultices, and the swallowing of small pieces of ice: in other words, the treatment, after emptying the parts affected, should be that of acute gastro-enteritis.

HYDRARGYRUM.

The only preparations of mercury which are used as purgatives are *calomel* and *blue mass*. Of these the first is by far the more active, and indeed is the only one which can be relied on to purge, since the *pilulæ hydrargyri* very frequently will, if given by themselves, fail to induce liquid stools.

The chief interest in the purgative action of mercurials centres in the question as to their influence upon the liver. The evidence at present derivable

from experiments upon the lower animals has already been discussed, and the decision arrived at that it must be rejected.*

When calomel is given to a healthy man in moderate purgative doses, green liquid stools are produced, which, after larger doses, are replaced by brown passages. The color of these passages has been always supposed by clinicians to be due to the presence of bile; but recently it has been affirmed that the green tint is owing to a compound of the mercury itself. Although no chemical proof of the presence of the metal or its salt has, that I am aware of, been furnished, yet it can scarcely be doubted that, in the first passages produced by calomel, mercury is present. The question evidently is not, Is mercury ever present in the "spinach-stools"? but, Is it always present? or, in other words, Is it an integrant portion of them? The evidence is not so abundant upon this point as is desirable, but yet seems sufficient to furnish a negative answer to the last question. Simon (*Animal Chemistry*, Sydenham Soc. Translations, vol. ii. p. 386) and Golding Bird (*London Medical Gazette*, 1845, p. 801), in careful analyses, both failed to detect the metal; and, as the recognition of mercury is an exceedingly simple chemical problem, it seems impossible that these chemists could have overlooked the metal if it had been present. Simon's analysis was performed upon the fifth stool after the administration of a large dose of calomel. The passage was fluid, perfectly green, had no fecal odor, exhibited a mild acid reaction, and showed, under the microscope, a great number of mucous corpuscles and epithelium-cells. Ether extracted from the solid residue (obtained by evaporation) a considerable amount of fat, which had an acid reaction, contained cholesterin, and was colored by biliverdin. All the other substances, which were separated from the stool by water and alcohol, were more or less colored by bile-pigment. Bilin, with bilifellinic acid and biliverdin, were found in large quantity.

The most satisfactory evidence is, however, that furnished by Michéa (*London Lancet*, 1849, vol. i. p. 15), who examined chemically the feces under four different conditions. First, the spontaneous dejections of six healthy individuals: no bile was detected. Second, green stools of three persons suffering from gastro-intestinal derangement: bile-pigment was found in one case only, and in that could not be detected after persistent vomiting had ceased. Third, calomel having been given to eight healthy persons, five men and three women, bile was readily demonstrated in the green passages produced in all of the subjects. Fourth, saline and resinous purgatives were given to five persons, but no bile could be detected in the liquid stools.

To the evidence brought forward in favor of the proposition that calomel given to healthy men causes an increased escape of bile from the alimentary canal, may be added the conclusive fact that in some persons, whose idiosyn-

* For a very elaborate and careful review of the clinical evidence as to the action of calomel on the liver, see Dr. Thos. R. Fraser's paper in the *Edinburgh Medical Journal*, April, 1871.

crasies render them very susceptible to the action of calomel, it produces not merely purging, but also vomiting of bile, which is scarcely at all altered.

From the facts which have just been passed in review, the conclusion seems inevitable that mercurial purgatives given to healthy persons cause the escape of large quantities of bile from the alimentary canal.

As is well known, when from any cause bile does not pass into the duodenum, the stools become very pale, of a peculiar potter's-clay, or even white, color. Very frequently under these circumstances, which may co-exist either with diarrhœa or with constipation, mercurials will modify the color of the passages and alleviate or cure any symptoms present. In many cases the mercurials are, of course, powerless to effect the desired result; but this depends upon the cause being organic, or of some other nature not to be overcome by a mere stimulant to secretion.

As mercurials in health increase the flow of bile from the intestine, and as they will sometimes re-establish it in disease when the secretion has altogether ceased or has been very materially diminished, the conclusion seems to me inevitable that mercurials have the power of directly or indirectly increasing the secretion of bile. The only objection of any force to be urged against this deduction is founded upon the idea that the drug simply increases peristalsis in such a way as to cause the bile naturally in the duodenum to be swept out instead of being absorbed. The answer to this is embraced in the following facts: mercurials restore the color of the passages when pale from arrested secretion, often without producing diarrhœa; other even more active purgatives fail to induce the same bilious passages; when diarrhœa exists with clayey stools, the change in the color of the passages caused by a mercurial may coincide with a not increased, or even a lessened, amount of liquidity; diarrhœa ordinarily does not cause bile to appear in the passages.

THERAPEUTICS.—A mercurial purge is especially indicated by the congeries of symptoms known as "*biliousness*:" a heavily-coated tongue, bitter, disagreeable taste, heavy headache, depression of spirits, loss of appetite, slight nausea, and light-colored passages. It should be borne in mind that one or several of these symptoms may be absent in any individual case. Of all single indications for the use of calomel, the occurrence of *potter's-clay-colored* passages is the most important; and if such stools exist, and do not depend upon an organic cause, a mercurial should be given, whether there be constipation or diarrhœa.

In *bilious fever*,—i.e., *malarial fever with congestion of the liver*,—a mercurial purge, or several mild mercurial purges, will often, by exciting the action of the hepatic gland, be of great service in preparing the way for or aiding in the action of quinine. In *catarrhal jaundice*, mercurials, on the whole, offer, I think, the most frequently successful mode of treatment. It is evident that in such cases calomel does good not merely by its cholagogue influence, but even to a greater extent by its antiphlogistic power, no doubt lessening the viscosity of the secretions and abating the inflammatory action in the hepatic ducts. In many instances it is well to exhibit a mercurial purge to start with;

but the main reliance is to be placed in the continuous exhibition of small doses of the drug until the gums are rendered slightly sore. Anything like profuse salivation is, of course, to be avoided.

In *dysentery* of an acute sthenic type, calomel has yielded, in my hands, better results than any other remedy. It probably acts as an antiphlogistic and as an alterative, not only to the liver, but to all of the intestinal glands.

In one or two cases of *obstructive enteritis*, with severe constipation, which I have seen treated with this drug after the failure of other remedies, improvement in the local and constitutional symptoms commenced simultaneously with slight ptyalism, and continued on to recovery.

RHEUM—RHUBARB.

The root of *Rheum palmatum*, and other species of *Rheum* growing in China, Chinese Tartary, and Tartary.

Rhubarb occurs in hard, irregularly cylindrical or roundish pieces, of a brownish-yellow color, peculiar bitter taste, and imparting to the teeth a sense of grittiness, due to the presence of great numbers of minute crystals of the oxalate of lime. There were formerly two chief varieties in market, the *Russian* and the *Chinese*. The first of these was the best, and was distinguished by the exterior of the pieces being cut or pared with a knife, and by a conical hole, evidently made for inspection with the point of a sharp instrument and never reaching beyond the centre of the mass. The cause of the superiority of this brand of rhubarb was the close governmental inspection which it received on the Russian frontier. Such pieces as failed to pass the officials found their way into commerce through Turkey, and constituted the so-called *Turkey Rhubarb*, which resembled the Russian in external characters, but was of somewhat inferior quality.

The *Chinese Rhubarb* was distinguished by the outside of the pieces having been scraped, and by the existence of a large hole running clear through and often retaining a portion of the cord upon which the roots had been strung to dry.

Owing to the expiration and non-renewal of the treaty between the governments of Russia and of Tartary, the only official variety of rhubarb now in the market is the Chinese.

Besides the true officinal varieties from Asia, there is a drug in commerce which from its source is known as *European Rhubarb*. It occurs in long, cylindrical pieces, or very often is cut to imitate one of the varieties of the genuine drug, from which it is to be distinguished* by its more spongy

* Dr. Carvet (*Vierteljahrsschrift für Prakt. Pharm.*, Heft i., 1873) gives the following method for distinguishing the varieties of rhubarb. If the piece be cut with a sharp knife and moistened, on attentive examination one or other of the following characters will be discovered:

European Rhubarb.—The surface marked by alternate red and white lines radiating from the centre. These lines are, in cylindrical sticks, a little way from the periphery, but, in

texture and by the complete or almost complete absence of grittiness when chewed.*

The active principles of rhubarb have not all been made out: it certainly contains chrysophanic acid, and a peculiar tannic acid, to which it owes its astringency. *Chrysophanic Acid* crystallizes out of alcohol in orange-yellow, golden, shining needles; out of benzole in orange-yellow or pale six-sided rhombic plates; with nitric acid it produces a fine yellow color; with the alkalis, a beautiful purple-red. According to Schlossberger, Bucheim, Meykow, and Auer, it is not purgative; but Schroff has found it to be so.† It certainly is not the chief purgative principle of the drug. The substances known as *Rhein* and *Rhabarbarin* are complex bodies.

PHYSIOLOGICAL ACTION.—Rhubarb is somewhat stomachic, tonic, actively purgative, and, owing to its tannic acid, secondarily astringent, leaving a decided tendency to constipation after the primary purgation. Owing probably to its chrysophanic acid, it tinges the milk of nursing women yellow, as it does also the urine, which on the addition of an alkali assumes a purplish-red color. It is generally asserted to affect chiefly the muscular coat of the bowels, and to purge by increasing peristalsis; but I have never met with any proof of the accuracy of this common belief.

THERAPEUTICS.—Notwithstanding its astringent property, rhubarb is largely used as a habitual laxative, because it does not impair, but, on the contrary, seems to strengthen, the appetite and the digestion. It should not be used in a high sthenic state of the system, or when depletion is necessary, but is, on the other hand, the best purgative when it is desired simply to unload the bowels in a debilitated subject.

In *diarrhœa*, with intestinal weakness or relaxation, it is the best purgative with which to unload the bowels of acrid secretions retained in them; and in the form of the *aromatic syrup* combined with an alkali it is especially valuable in the *summer bowel-complaints* of children when the stools are greenish and mucous.

flat pieces, very near the periphery, intersected by a brown zone, which is sometimes interrupted.

Russian Rhubarb.—The surface offers on a whitish ground, yellow, sometimes scarcely perceptible, long and short anastomosing lines, often interrupted by radiating systems. These star-like masses are roundish, or long, and of various sizes: their rays are, near to the middle point, clear yellow; farther out, brownish. The major lines have a general course to the periphery, marked by the anastomoses, irregularities, and star-clusters.

Chinese Rhubarb.—The section presents clear yellow rays which, in much bent lines, pass from the centre to the circumference. These lines often appear to anastomose, and to form irregular stars, whose outer part is composed of more rays than their inner. This is especially seen in the cambium zone. As in the Russian rhubarb, between the radiating lines there are sometimes irregular stars.

* In the past, the European rhubarb has been considered of little value; but recently it has been claimed that it is as good as the Asiatic. See *U. S. Dispensatory*, 13th ed., p. 735; also Prof. Radius (*Apotheker Zeitung*, Bd. vi., 1871).

† See *Die Pflanzenstoffe*, p. 985.

ADMINISTRATION.—Rhubarb is seldom employed in powder, but, when used, may be given in the dose of twenty grains. In chronic constipation, small pieces of the root are very often carried in the pocket and chewed by the person affected *pro re nata*. The preparations of rhubarb are as follows:

Infusum Rhei—*Infusion of Rhubarb* (℥ii to Oss). Dose, half to one wineglassful.

Syrupus Rhei—*Syrup of Rhubarb* (℥iii to Oij). Dose, one to two tablespoonfuls.

Syrupus Rhei Aromaticus—*Aromatic Syrup of Rhubarb* (℥iiss to Ovii). This preparation is too feeble for use as a purgative for adults. The dose for infants is from one to two teaspoonfuls.

Tinctura Rhei—*Tincture of Rhubarb* (℥iiss to Oj). Dose, one to two tablespoonfuls.

Tinctura Rhei et Sennæ—*Tincture of Rhubarb and Senna*—*Warner's Gout Cordial* (Rhubarb, ℥i; Senna, ℥ii to Oij). Dose, one to three tablespoonfuls.

Pulvis Rhei Compositus—*Compound Powder of Rhubarb* (Rhubarb, ℥iv; Magnesia, ℥xii; Ginger, ℥ii). Dose, half a drachm to a drachm.

Pilule Rhei—*Rhubarb Pills* (grs. iii). Dose, four to eight pills.

Pilule Rhei Compositæ—*Compound Rhubarb Pills* (Rhubarb, grs. ii; Aloes, grs. iss). Dose, two to four pills.

Extractum Rhei—*Extract of Rhubarb*. Dose, five to ten grains.

Extractum Rhei Fluidum—*Fluid Extract of Rhubarb*. Dose, ten to twenty-five drops.

JUGLANS.—The inner bark of the *Juglans cinerea*, or common butternut, or white walnut, as it is variously called, is recognized under the above title by the U. S. Pharmacopœia, which also contains among its preparations an extract of it (*Extractum Juglandis*; dose, twenty grains). *Juglans* is said to be a mild cathartic, resembling rhubarb in its action, as a substitute for which it was introduced to the profession during the Revolutionary war by the famous Dr. Rush.

ALOES.

The U. S. Pharmacopœia recognizes three varieties of aloes, which are all obtained in a similar manner,—*i.e.* by cutting off the thick succulent leaves of the various plants, standing them up, allowing the juice to drain into suitable vessels, and afterwards inspissating either by exposure to the sun or by slowly evaporating. The leaves contain a large amount of a very thick mucilaginous juice, which escapes on pressure: hence aloes prepared by expressing the leaves or by boiling them—both of which processes are sometimes used—is very inferior.

ALOE BARBADENSIS.—*Barbadoes Aloes* is prepared in Barbadoes and other West Indian islands from the *Aloe vulgaris*.

ALOE SOCOTRINA.—*Sociotrine Aloes* is the product of *Aloe Socotrina*,

which grows in the island of Socotra in the Red Sea, and on the neighboring coast of Arabia.

ALOE CAPENSIS.—*Cape Aloes* is obtained at the Cape of Good Hope, from the *Aloe spicata*.

Aloes are darkish extracts, of a bitter nauseous taste, yielding their virtues to alcohol, imperfectly to water, and very perfectly to alkaline solutions. The Cape aloes is said to be "characterized by its dark olive or greenish-black color, its smooth and very glossy surface when broken, its translucency at the edges, and the fine bright-yellow color of its powder, which is slightly tinged with green." The Socotrine aloes is distinguished by its yellowish-brown or reddish-brown color, its translucent edges, agreeable aromatic odor, and beautiful golden-yellow powder. The Barbadoes aloes is characterized by its dark-brown or reddish-brown color, its dull fracture, opaque edges, disagreeable nauseous odor, and dull olive-yellow powder. Of these varieties the Socotrine is most esteemed in human medicine. The Barbadoes is said to be the strongest, but is employed almost exclusively in veterinary surgery.

Messrs. T. and H. Smith (*Chemical Gazette*, 1851) in 1850 discovered in Barbadoes aloes a crystalline principle,—aloin,—which was shortly afterwards found by Pereira to exist already crystallized in the sap of various species of aloe plants, and was subsequently obtained by Groves (*Pharmaceutical Journal*, xvi.) from Socotrine aloes. *Aloin* crystallizes from its watery solution in sulphur-yellow granules, from a hot alcoholic solution in star-like groups of needles. It is neutral, odorless, of a taste at first sweetish, afterwards intensely bitter; soluble with difficulty in cold, freely in boiling, water and alcohol. Aloin was formerly believed to be the active principle of aloes; but at present the evidence is very decidedly against this view. According to T. and H. Smith, in doses of one or two grains it is an almost drastic purge. On the other hand, Robiquet (*Journal de Pharmacie*, t. xxix.) took fifteen grains of it without effect, and affirms that it is a simple bitter. Th. Husemann (*Die Pflanzenstoffe*, p. 1047) found that upon himself four and a half grains, upon dogs five grains,—by the mouth or injected into the veins,—exerted no purgative influence; and Wm. A. Tilden (*Transactions of the British Pharmaceutical Society*, 1872) has obtained similar results. As he states, it is most probable that the purgative property of aloes resides in the soluble, brown, uncrystallizable substance which constitutes a considerable portion of it.

PHYSIOLOGICAL ACTION.—Aloes is a stomachic, stimulant cathartic, remarkable for the slowness of its action. It has been supposed to influence chiefly, if not solely, the large intestine, and the clinical evidence is very strong that in overdoses it produces irritation of the rectum. The belief, formerly so universal, that it is capable of producing hæmorrhoids, has been very much weakened by the researches of a number of modern observers, among whom may be mentioned Trousseau and Pidoux (*Stillé, Therapeutics*, vol. ii. p. 444). Its habitual use in large doses is said to cause tenesmus, a

feeling of weight, heat, and uneasiness in the pelvis, and occasionally excitation of the sexual organs. Although I have used it a good deal, however, I have never seen these results. It undoubtedly has a tendency to increase the menstrual flow.

Aloes is certainly absorbed, as is shown by the fact, attested by Dr. Gerhard (*North American Medical and Surgical Journal*) and other observers, that it will purge when its powder is sprinkled upon a blistered surface. In regard to its action in combination, Prof. Stillé states that when other and quickly-operating cathartics are taken along with aloes it does not appear to modify their action; but if it is administered seven or eight hours before a saline or other active purgative, a combined and very powerful operation is the result.

THERAPEUTICS.—Aloes in small doses is one of the best remedies for *constipation* of atonic subjects, especially when a stomachic stimulant is indicated. In these cases it may often with great advantage be combined with a simple bitter, one or two grains of it taken directly after meals being generally sufficient. In the constipation of *plethora* it should not be employed; neither should it be administered when active abdominal or rectal inflammation exists. During pregnancy it may be used as a laxative, but, unless some especial indication calls for its use, it is best avoided. Large purgative doses of it should never be given to pregnant women, as it certainly irritates the pelvic organs, and is even said to have the power of causing abortion.

Formerly it was taught that aloes should not be used in *hemorrhoids*; but most, if not all, of the cases of this affection depend upon a condition of relaxation of the rectal veins, and Dr. Fordyce Barker (*American Practitioner*, 1872) insists upon the great value of aloes in piles, and states that Oppolzer was especially famous for his treatment of this affection, and that his prescriptions were, when piles are associated with constipation, aloes and quinine; without constipation, aloes and sulphate of iron. For bleeding piles he used R.—Ferri sulphat., $\mathfrak{z}\text{i}$; ext. aloës aq., $\mathfrak{z}\text{i}$; ext. taraxaci, q. s. Ft. pil. no. 60. S.—One morning and evening, and increase to three a day if necessary.

When costiveness accompanies atonic *amenorrhœa*, aloes alone of all the laxatives should be exhibited; and it is also of service in atonic *menorrhagia*.

ADMINISTRATION.—Aloes is very rarely or never used by itself to produce free purgation, but may be given in the dose of from ten to twenty grains; in the dose of from three to five grains it is a decided laxative. As aloes often contains sticks and other extraneous matters, the U. S. Pharmacopœia directs that an *Aloe Purificata*, or *Purified Aloes*, should be made by dissolving the crude drug in alcohol, straining, and evaporating. The following are the other official preparations:

Tinctura Aloës—*Tincture of Aloes* ($\mathfrak{z}\text{i}$ to Oij). Dose, as a laxative, one to three teaspoonfuls.

Tinctura Aloës et Myrrhæ—*Tincture of Aloes and Myrrh*—*Elixir Pro-*

prietatis (Aloes and Myrrh, āā ʒiii to Oij). Dose, as a laxative, one to two teaspoonfuls.

Vinum Aloës—*Wine of Aloes* (ʒi to Oj). Contains also cardamom and ginger. Dose, as a laxative, one-half to one teaspoonful.

Pilulæ Aloës—*Pills of Aloes*. These contain each two grains of aloes and two grains of soap.

Pilulæ Aloës et Assafetidæ—*Pills of Aloes and Assafetida*. These are useful in costiveness of hysterical or old subjects: each pill contains four grains of a mass composed of equal proportions of aloes, assafetida, and soap.

Pilulæ Aloës et Mastiches—*Pills of Aloes and Mastich*. These are the famous "Lady Webster Dinner-Pill," each containing two grains of aloes.

Pilulæ Aloës et Myrrhæ—*Pills of Aloes and Myrrh*. These are used in amenorrhœa, and contain two grains of each ingredient in every pill.

Pulvis Aloës et Canelle—*Powder of Aloes and Canela*.—*Hiera Picra*. Used in the constipation of amenorrhœa. Dose, ten to twenty grains.

SENNA—SENNÆ.

The leaflets of the shrubs *Cassia acutifolia* and *C. obovata*, of Nubia and Upper Egypt, and of *C. elongata*, of Southern Arabia. The senna leaves vary from three-fourths of an inch to an inch and a half in length, and are to be distinguished by the inequality of their bases, the two sides of the lamina or leaf-blade joining the midrib at unequal heights and angles. There are three commercial varieties of senna, which are named from the places of their export, as follows:

Alexandria Senna, the most common variety, distinguished by the presence of the shorter *argel-leaves*, with equal bases, by the ovate-pointed leaflets of *Cassia acutifolia*, and by the scattered mucronate-obovate leaflets of *C. obovata*;

India Senna, characterized by the oblong leaflets, being from one to two inches in length, entire and perfect;

Tripoli Senna, to be recognized by the great extent to which the leaflets are broken up.

The active principles of senna have not been completely isolated, but appear to be several. *Cathartic Acid*, discovered by Draggendorff and Kubly (*Vierteljahrsschrift für Prakt. Pharm.*, Bd. xvi.), is undoubtedly actively cathartic, as Kubly found that fifteen grains of it dissolved in an alkaline solution produced in six hours frequent watery discharges, with griping; but the researches of Bourgeois and Bouchut (*L'Union Pharmaceutique*, Nov. 1871) indicate that, besides cathartic acid, there are in senna chrysophanic acid, and a purgative principle which has not as yet been isolated. The *cathartine* of the older writers is undoubtedly a complex body.

PHYSIOLOGICAL ACTION.—Senna is a very powerful cathartic, producing watery fecal discharges, and acting, it is said, as readily upon swine, dogs, cats, and horses as upon man. According to Prof. Stillé, both Courten and

Regnaudot found that its infusion injected into the veins caused vomiting and purging. It is undoubtedly absorbed; and Bergius affirms that it will impart its purgative property to the milk of nursing women.

THERAPEUTICS.—Whenever a brisk, somewhat irritating cathartic is desired, senna may be selected. When given alone, it is very apt to gripe severely, and is consequently more often used in combination,—especially its infusion with Epsom salts (*Black Draught*). In obstinate *fecal accumulation* the Black Draught constitutes a most efficient and safe remedy. It is claimed that senna does not leave a tendency to constipation after its action; and hence, in small doses, it is preferred by some as an habitual laxative.

ADMINISTRATION.—Whenever senna is exhibited, an aromatic should be united with it, to lessen its tendency to gripe. The leaves are not given in substance. The official preparations are as follows:

Infusum Sennæ—*Infusion of Senna* (℞i to Oj). Dose, two to four fluid-ounces.

Extractum Sennæ Fluidum—*Fluid Extract of Senna*. Dose, two fluidrachms to half a fluidounce.

Confectio Sennæ—*Confection of Senna*. A very complex but elegant preparation, used only as a laxative, and not suited to dyspeptic cases, on account of its tendency to derange the digestion. Dose, one to two drachms at bedtime; a very good laxative in pregnancy.

SALINE PURGATIVES.

The following salts of magnesium, potassium, and sodium are official:

MAGNESII SULPHAS—SULPHATE OF MAGNESIUM (MgOSO_3 — MgSO_4).

This drug is commonly known as *Epsom Salts*, a name derived from the spring by the evaporation of whose waters it was originally prepared. It is now manufactured on a large scale from dolomite, the double carbonate of calcium and magnesium, and from native siliceous hydrite of magnesium.

Sulphate of magnesium ordinarily occurs in small, acicular, slowly-efflorescent crystals, containing about fifty-one per cent. of water of crystallization, soluble in their own weight of water at ordinary temperatures. By proper precautions, it may be obtained in large quadrangular prisms, terminating in a four-sided pyramid or a dihedral summit. The taste is bitter, saline, nauseous.

PHYSIOLOGICAL ACTION.—Epsom salts is a most active hydragogue cathartic, producing very large watery discharges without causing any irritation of the intestines. In very large doses it is, however, capable of producing fatal hypercatharsis; and Christison reports the case of a boy ten years old, said to have been killed by two ounces of the salt, without the induction of purgation.

THERAPEUTICS.—Whenever, in inflammation, it is desired to deplete

through the bowels, the sulphate of magnesium offers the most advantageous method of doing it. Especially is this the case when the intestines are affected, as in *enteritis* or in *colitis*. There is probably no other purgative in common use which produces at the same time such free serous evacuations and so little intestinal irritation. On account of the efficiency of its action and the watery character of its discharges, it is especially applicable to cases of *fecal accumulation* and of obstinate *colica pictorum*. The dose is half an ounce to an ounce, properly diluted.

LIQUOR MAGNESII CITRATIS.—*Solution of the Citrate of Magnesium* is prepared by putting into a strong bottle a syrupy solution of the citrate of magnesium containing an excess of citric acid, adding the bicarbonate of potassium, and corking tightly. On account of its agreeable taste and effervescence, this preparation is much used as a purgative. It is similar to Epsom salts in its action, but is less efficient, more apt to gripe, and more irritating. It ought not to be used in inflammatory affections of the bowels.

SODII SULPHAS.—*Sulphate of Sodium* ($\text{NaO}, \text{SO}_3 - \text{Na}_2\text{SO}_4$), or *Glauber Salts*, is at present manufactured from common salt by means of sulphuric acid. It occurs in six-sided, very efflorescent, striated prisms, which finally crumble into a white powder. Its taste resembles that of Epsom salts, but is more nauseous. On this account, and because its action upon the economy is similar to but harsher than that of the sulphate of magnesium, in medical practice it has been completely superseded by the latter salt. The dose is a quarter of an ounce to half an ounce, properly diluted.

SODII PHOSPHAS.—*Phosphate of Sodium* ($2\text{NaO}, \text{HO}, \text{PO}_3 - \text{Na}_2\text{H}, \text{PO}_4$). This salt occurs in colorless, transparent crystals, which speedily effloresce and become opaque when exposed to the air. It is a tribasic phosphate, one part of water acting as a base. This salt is soluble in four parts of cold water, and has a saline taste, closely resembling that of common salt. In large doses it is said to be a mild saline purgative, but as such is at present very rarely used.

Within a few years the use of the phosphate of sodium in chronic infantile diarrhœa has attracted a good deal of notice. Originally recommended, so far as I know, by Dr. Routh, in his work on Infant-Feeding, as being a valuable nutrient or alterative in children, attention has been especially drawn to the phosphate of sodium by Dr. William Stephenson (*Edinburgh Medical Journal*, 1867, vol. xiii. p. 336). He believes that it acts upon the liver, and adduces much clinical evidence in favor of his views. The cases in which he especially recommends it are “infants who are being artificially reared, and who are liable to frequent derangement of the bowels; also when the phosphatic elements in the food seem deficient; where, from the character of the motions, there is a deficient or defective secretion of bile. It is thus of service in cases of chalky stools or white fluid motions, and in many cases of green stools; also in duodenal dyspepsia, and in diarrhœa after weaning.”

Dr. S. G. Webber (*Boston Medical and Surgical Journal*, 1868, vol. i. p. 5) confirms these statements of Dr. Stephenson, and in a few cases I have seen very notable effects from this use of the drug.

POTASSII ET SODII TARTRAS—*Tartrate of Potassium and Sodium.—Rochelle Salts* is made by the addition of the carbonate of sodium to a solution of the bitartrate of potassium. It occurs in large, colorless, transparent, slightly efflorescent, prismatic or half-prismatic crystals, which are soluble in two and a half parts of cold water. Its taste is slightly saline. It is a mild saline purgative, decidedly less efficient, but much less offensive to the palate, than Epsom salts. The dose is from half an ounce to two ounces, properly diluted.

PULVERES EFFERVESCENTES APERIENTES—*Aperient Effervescing Powders*. The single *Seidlitz powder* is a duplex entity, occurring in two packets, that in the white paper containing thirty-five grains of tartaric acid, that in the blue paper forty grains of the bicarbonate of sodium and two drachms of Rochelle salts. When they are taken, the powders are dissolved separately, the solutions added, and the whole drunk whilst effervescing. They are very acceptable to the stomach, refrigerant and laxative rather than purgative. Seidlitz powders are used almost exclusively to evacuate the bowels, and exhibited after blue mass to "carry off" mercurials, etc. They should be taken on an empty stomach, as before breakfast. One powder is the usual dose; but not rarely even two powders will fail to purge.

POTASSII SULPHAS.—The *Sulphate of Potash* (*Potassii Sulphas*), or *Vitriolated Tartar*, was formerly used as a purgative, but is no longer employed in medicine, except in the preparation of Dover's powders.

DRASTICS.

As already stated, the *drastics* are those vegetable cathartics which are actively irritant. With perhaps one or two exceptions, in sufficient amount they are capable of causing fatal gastro-intestinal irritation. The line between the drastics and the stronger purgatives is, of course, placed more or less arbitrarily, since the various cathartics differ in action almost by insensible degrees. Thus, jalap, although included among the drastics in this work, might with perhaps even greater propriety be classed among the purgatives, since it is very little more active or irritant than is senna.

Further, these remedies in combination seem to lose, in a measure, their power of causing irritation, and to become useful purgatives. A fact, however, which makes the classification here employed clinically useful, although it be not scientifically accurate, is that none of these remedies should be used when a purgative is desired to relieve gastro-intestinal inflammation or irritation; and, on the other hand, when a revulsive action is wished for, as in some cases of brain-disease, one of the drastics should always be selected.

JALAPA.—JALAP.

The tuber of *Exogonium purga*, a convolvulaceous vine growing in Mexico. Jalap comes into the market in two forms: one, that of the younger roots, which are sold undivided; the other, that of the old roots, which are brought into the market in transverse or longitudinal slices, and in pieces. The first variety consists of very hard, irregularly globular, brittle roots, about the size of a shut fist, or smaller, and often slashed with vertical incisions, made for the purpose of facilitating drying.

The section of jalap is always distinctly resinous, if not to the naked eye, at least to the vision aided by a lens. The active principle is a duplex resin, one portion of which is soft, and soluble in ether, the other (*Rhodeoretin*) hard, and insoluble in the latter menstruum. Rhodeoretin is asserted both by Kayser (*U. S. Dispensatory*) and by Mayer (*Ibid.*) to be the purgative principle; but Mr. John C. Long (*Amer. Jour. of Pharmacy*, Nov. 1861) has shown that the soft resin is equally active. The percentage of resin varies very much in different specimens of the root, but is much greater in those that are worm-eaten.

PHYSIOLOGICAL ACTION.—Upon dogs and horses jalap (Stillé, *Therapeutics*, vol. ii.) is said to act as a powerful hydragogue cathartic, and in overdoses as a gastro-intestinal irritant. If the experiments of Cadet de Gassicourt be correct, it probably acts by absorption, as, according to that observer, diarrhoea may be produced in dogs by its free application to the shaven skin. Prof. Stillé, however, asserts that it does not impart its purgative properties to the milk of nursing women, and that in man it is not absorbed by the skin. In man jalap produces free hydragogue catharsis, often with nausea; or, if in overdoses, violent vomiting and purging.

THERAPEUTICS.—Jalap is especially indicated when it is desirable to produce large watery stools. It is, however, very rarely used alone. A favorite combination with many practitioners is of it and calomel. In the form of the *compound powder* (*Pulvis Jalapæ Compositus*: jalap, one part; cream of tartar, two parts), jalap is very frequently used with great advantage in *ascites* and also in other forms of general *dropsy*. It is believed when given in this way to exert some influence upon the renal functions: for very many cases the proportion of cream of tartar in the officinal compound powder is too small, and should be increased.

ADMINISTRATION.—The dose of powdered jalap is ten to twenty grains. The officinal *extract* (*Extractum Jalapæ*) is about twice as strong as the crude drug.

Resina Jalapæ.—*Resin of Jalap* is, like the other purgative resins, prepared by precipitating a saturated tincture with water. According to Husemann, the ordinary adulterations of jalap resin may be detected by the solubility of the substances employed in the oil of turpentine, which does not affect the genuine article. On account of its tastelessness, this preparation, as well as the similar one of scammony, is sometimes employed as a purgative for children. The dose for an adult is from two to four grains.

COLOCYNTHIS—COLOCYNTH.

The fruit, deprived of its rind, of the *Citrullus Colocynthis*, or bitter cucumber, a vine growing in the Cape of Good Hope, Japan, Syria, Egypt, Turkey, the islands of the Grecian Archipelago, etc. The fruit is a round gourd, from two to four inches in diameter, of a whitish or pale-yellow color. It occurs in the market with or without its rind. The pulp is dry and membranous, whitish, and contains the active purgative glucoside *Colocynthin*, first discovered by Herberger.

PHYSIOLOGICAL ACTION.—The experiments of Orfila and Schroff have shown that upon dogs and rabbits colocynth acts very much as it does in man, producing copious watery evacuations, and, although not so irritant as gamboge, in overdoses causing death by gastro-intestinal irritation.

If the statement of Richter that violent purgation may be induced by rubbing the abdomen with tincture of colocynth be true, the active principle is without doubt absorbed.

THERAPEUTICS.—Colocynth is rarely, if ever, used alone, but is given in combination with the other drastics, or with milder purgatives, to increase their activity. It is frequently added in small quantity, with advantage, to laxatives, especially when the constipation is somewhat obstinate. Its minute bulk is often of great advantage in these cases. Neither colocynth nor any of its combinations should be used in *dropsy*. Its chief use is, as a laxative, to unload the bowels, and to deplete directly from the portal circulation: even in cases offering these indications it is almost never used alone.

TOXICOLOGY.—Colocynth has not rarely produced death preceded by hypercatharsis and the usual symptoms of gastro-intestinal irritation. The fatal dose probably varies very much. Christison records the death of a woman twenty-four hours after taking a teaspoonful and a half of the powder. Roques chronicles a fatal result produced by less than a drachm of the powder in decoction, but, on the other hand, narrates a case in which three drachms failed to kill (Husemann, *Handbuch der Toxicologie*, p. 625).

ADMINISTRATION.—The officinal preparations of colocynth are as follows:

Extractum Colocynthidis—*Extract of Colocynth*.—Useful as an addition to laxatives. Dose, as a purgative, three to five grains.

Extractum Colocynthidis Compositum—*Compound Extract of Colocynth*.—Extract of colocynth, three and a half parts; purified aloes, twelve parts; resin of scammony, three parts; cardamom, one and a half parts; soap, three parts. A very useful purgative preparation, either as a laxative in minute dose (one to three grains), or in large dose (five to twenty grains) as an active purgative.

SCAMMONIUM—SCAMMONY.

A resinous exudation from the root of the *Convolvulus Scammonia*, a vine growing in Syria. It is said to be obtained by cutting off the root obliquely about two inches from the origin of the stems, and catching in shells the

few drachms of milky juice which exude from each root. From these shells it is emptied into some vessel and allowed to concrete. Before exportation it is usually adulterated with the expressed juice of the leaves and stalks, and with chalk, flour, ashes, sand, etc. This adulteration was formerly carried to a very great extent, but is at present indulged in to a much less degree. The pure or *Virgin Scammony* is in irregular, rough, fissured masses, of various sizes, commonly solid, with a dull resinous fracture, and of a dark greenish color, inclining to black. The smell is peculiar, resembling that of old cheese. The taste after a time is acrid.

Fuctitious or adulterated scammony occurs in cakes of various sizes and shape, and is sometimes spoken of as *amylaceous* or *erythracous* scammony, according to the material used for its adulteration. The active principle of scammony is a resin, very similar to that of jalap. The proportion of this resin in the drug varies, according to the purity of the article, from eight to ninety per cent. The U. S. Pharmacopœia directs that official scammony shall contain seventy-five per cent. of the resin.

PHYSIOLOGICAL ACTION.—Scammony acts upon the system like jalap, but is somewhat more irritating, and therefore more apt to gripe severely, and is still more strongly contra-indicated in inflammatory diseases of the intestinal canal. It is decidedly less drastic than gamboge.

THERAPEUTICS.—Scammony is never used alone, but in combination with other less active cathartics. The only official preparation is *Resina Scammonii*, which is prepared in the same way as that of jalap, than which it is more irritating and somewhat more active.

As all of the ingredients have been noticed, the present seems the proper place for the consideration of **PILULÆ CATHARTICÆ COMPOSITÆ**—*Compound Cathartic Pills*. These famous pills contain each: compound extract of colocynth, one and a third grains; extract of jalap and calomel, each one grain; gamboge, one-fourth of a grain. As there is soap in the first ingredient, the calomel is sooner or later reduced to the black oxide of mercury.

The compound cathartic pills are a very efficient purgative, generally not producing much griping. They cause large watery stools, and are used when it is desired to empty the bowels and deplete from the portal circulation.

PODOPHYLLUM—PODOPHYLLUM.

The rhizome of the *Podophyllum peltatum*, or May-apple, a perennial herb, growing in the Northern and Middle United States. *Podophyllum* occurs in simple or branched, cylindrical, brownish pieces, about the thickness of a goose-quill, smooth or wrinkled longitudinally, often obscurely marked with the scars of leaf-scales, and furnished with numerous rootlets or their remnants attached to the lower surface. The taste is bitterish, acrid, nauseous. The rhizome contains the alkaloid *Berberina*, but the purgative power resides in two resins, one soluble, the other insoluble, in ether.

PHYSIOLOGICAL ACTION.—The experiments of Dr. Snow, of Dr. S. R. Percy (*American Medical Times*, vol. iv.), of Dr. F. E. Anstie (*Med. Times and Gazette*, March, 1863), and of others, have shown that, whether administered by the mouth or hypodermically, podophyllum produces in the lower animals purging, with colicky pains and sometimes vomiting. If the dose be sufficient, the stools are bloody, and after large amounts death occurs, preceded by hypercatharsis, prostration, and slight convulsions. On post-mortem examination, Dr. Anstie found intense inflammation, with ulceration of the mucous membrane of the small intestines. Upon man podophyllum acts as upon these animals, producing in large doses violent catharsis, accompanied by much pain, and is no doubt capable of acting as a fatal gastro-intestinal irritant, although I am not aware that any case of serious poisoning by it is upon record. The experiments already alluded to show that it acts by being absorbed, which is confirmed by the experience of Dr. Percy, who found that its application to an ulcer was followed by its specific effects. It has been claimed that it acts especially upon the liver; and much clinical testimony has been adduced to support this view. I have never practically investigated this subject, but the evidence appears to be very strong that in many cases the use of large doses of podophyllum is followed by the free escape of bile per anum. This result seems, however, to be the consequence of the intense duodenal irritation rather than of a direct action of the drug upon the liver.

THERAPEUTICS.—Podophyllum in its action as a purgative resembles very closely jalap, but acts more slowly and more continuously than that drug, and is probably more apt to produce "bilious discharges." In my experience, it more often causes griping pain. In this country it is very largely used in cases of *acute constipation*, and with very good effect; and it has also been commended very highly by numerous practitioners in the so-called "*bilious attacks*." Owing to the extreme slowness of its action, it is not well adapted for combination with brisker cathartics. The same quality fits it, however, for use with calomel, it requiring nearly the same length of time as that drug to produce purgation.

ADMINISTRATION.—Although an extract is officinal, podophyllum is scarcely ever given in other form than that of the resin.

Resina Podophylli—*Resin of Podophyllum*.—*Podophyllin* is officinally prepared by precipitating a concentrated tincture with water: as berberina is soluble in the latter menstruum, the resin as thus obtained is free from the alkaloid. The portion of this duplex resin which is soluble in ether is certainly more actively purgative than the other part, to which, indeed, any purgative property is denied by some. Dose, as a purgative, from one-sixth to one-fourth of a grain; as a laxative, one-twelfth of a grain.

ELATERIUM—ELATERIUM.

A substance deposited by the juice of the fruit of the *Momordica Elaterium*, or squirting cucumber, a native of Greece, but cultivated in England. In the interior of the ovate fruit is an elastic sac, which contains the seeds and at ripening becomes so distended with juice that when the fruit falls off the vine, and the support is removed from the stem end, a rupture occurs at the latter position, and the liquid with the seeds is forcibly projected. The elaterium is said to be contained only in this inner juice. In order to avoid loss, the fruit is picked with a piece of the stalk adherent to it before ripening, and is opened by slicing.

Elaterium occurs in light, friable, slightly incurved, greenish-gray cakes about a line thick. The taste is acrid and bitter; the fracture finely granular. The active principle is *Elaterin*, which was first separated in a pure state by Morries (*Repertorium für Pharmacie*, xxxix. p. 134). It crystallizes in colorless, shining, rhombic, six-sided, odorless tables, of a very bitter sharp taste and neutral reaction. According to Dr. H. Köhler (*Virchow's Archiv*, Bd. l. p. 287), it is insoluble in water and glycerine, readily soluble in cold alcohol, and soluble with difficulty in ether and turpentine.*

PHYSIOLOGICAL ACTION.—Locally applied, elaterium is a very decided irritant, producing, according to Pereira, ulcerations in the fingers of those who handle the fruit and prepare the drug for market. When taken internally, it acts on man as a most powerful hydragogue cathartic, producing, when the dose and administration are properly regulated, enormous watery stools, without much irritation. On the lower animals its action is much less certain. Viborg asserts that a horse was unaffected by a pound of elaterium fruit; and I have given one and even two grains of a presumably active elaterium to a dog without producing very obvious results. If the dose be sufficiently large, all animals probably are, however, fatally affected by elaterium, perishing by progressive depression. Prof. Stillé asserts (*Therapeutics*, vol. ii. p. 459) that the death is not rarely preceded by violent vomiting and purging; and even when these are absent during life, post-mortem examination reveals congestion and inflammation of the gastric and intestinal mucous membranes. In none of my own experiments, which have not been very numerous, has any purging been present; further, in Dr. Köhler's elaborate investigation (*loc. cit.*), elaterium dissolved in alcohol was injected under the skin, the powdered elaterium was put into the rectum, and was given by the mouth after the gall-duct had been tied so as to prevent the flow of bile into the intestine, and in neither case was there any purging, but prostration, apathy, disturbed respiration, salivation, and violent convulsions, ending in death. From these experiments Dr. Köhler draws the conclusion that

* For the behavior of it with various reagents, and for the methods of searching for it in medico-legal investigations, the reader is referred to Dr. Köhler's memoir.

elaterium exerts a general action upon the system, for which its introduction into the blood is all that is requisite, and also a purgative influence, for which it is necessary that there be bile in the duodenum to dissolve the elaterium and cause it to act locally on the intestine. The objection to this conclusion is that our present light seems to indicate that elaterium does not purge dogs and rabbits, even when given under ordinary circumstances. Further, there is considerable evidence to show that elaterium applied externally will cause purging in man (see Stillé, *Therapeutics*, vol. ii. p. 459). If this be true, the application of the conclusions arrived at by Köhler to man certainly would be incorrect. Köhler's experiment proved that in animals elaterium is absorbed, even when given by the mouth, since he found it in the urine of the poisoned dogs and rabbits.

THERAPEUTICS.—Elaterium is certainly the most efficient of all the hydragogue cathartics, producing in properly-regulated doses the freest evacuations with comparatively little pain and irritation. It is therefore indicated whenever a powerful cathartic of such nature is indicated. It is the most efficient of all the medicines of the class in general *dropsy* or in *ascites*. As, however, its action is very exhausting, great care should be exercised not to give it in too large doses, and also to support the strength of the patient during the period of purgation, and afterwards by alcoholic stimulants, easily-digested nutritious food, and appropriate hygienic measures.

In the latter stages of dropsy, the injudicious use of elaterium may favor, and no doubt has accelerated, the fatal result, by intensifying the exhaustion.

An idea has prevailed that elaterium is especially valuable in *uræmia*, because it produces an elimination of the urea in the stools; but I have been unable to find authority for the asserted elimination. Be this as it may, however, clinical experience has demonstrated the utility of elaterium in chronic renal disease.

In order to deplete, elaterium has been employed in various diseases; but this use is not to be encouraged, and especially when there is any gastrointestinal irritation or inflammation are the salines much preferable to elaterium. In cases of plethora, however, when there is a sudden *determination of blood to the head* and a very powerful impression is needed, the vegetable cathartic may be advantageously employed.

TOXICOLOGY.—Elaterium is without doubt capable of destroying life, and that, too, when not in large quantity. I know of but one recorded death,—that of a woman who took, by the advice of a quack, two and two-fifths grains of the extract of elaterium and sixteen grains of rhubarb. Violent and uncontrollable vomiting and purging came on, and proved fatal in thirty-six hours. After death, the gastro-intestinal mucous membrane showed marked evidences of inflammation. (See *Beck's Medical Jurisprudence*, 12th ed., vol. ii. p. 719.)

ADMINISTRATION.—Formerly, elaterium varied a good deal in power;

now, however, it is very uniform, and it is never safe to commence with more than one-sixth of a grain, to which should be united a grain of the extract of hyoscyamus and a drop of some aromatic oil.

GAMBOGIA—GAMBOGE.

A gum resin, obtained in Siam by breaking off the leaves and young shoots of the tree known by botanists as the *Garcinia Morella*, and catching in suitable vessels the juice as it drops. When the receptacles consist of hollow bamboos, the juice hardens into cylindrical casts, striated externally, and with a central cavity due to the loss of substance in drying. This is the so-called *pipe gamboge*. *Gamboge in sorts* occurs in irregular masses.

Gamboge is a hard, resinoid substance, of a brittle, often conchoidal fracture, of a deep reddish-orange color on exposed surfaces, more yellowish when freshly broken, affording a bright-yellow powder, insoluble in water, with which it forms, however, an intensely yellow emulsion. It has little or no taste, but when chewed produces, after a time, an acrid sensation in the fauces. It contains, according to Christison, as much as seventy-two per cent. of *gambogic acid*, a resinous acid of a cherry-red color, forming red salts with alkalis. This would appear, however, to be only one of the purgative principles of the drug, since it is less drastic than an equal weight of gamboge (Christison (*Ann. Chem. Pharm.*, xxiii, 185), Pabo (*Additam. quæd. ad Virtutes chem. et physiol. Resinarum quarundam comparatas*, Dorpat, 1851), Daraszkiewicz (*Meletemata de Resinarum, præsertim Resinæ Guttæ, in Tractu Intestinali Rationibus*, Dorpat, 1858)); five grains producing in some persons only watery evacuations, in others not even these, and as much as seventeen grains having been taken without more serious effects than severe purgation.

PHYSIOLOGICAL ACTION.—Administered by the mouth to dogs, cats, horses, and probably other of the lower animals, gamboge acts very generally as a violent drastic cathartic; but from the experiments of Schaur and of Orfila it would appear to cause sometimes simply vomiting, and, when in very large amounts, death, without any marked symptoms other than those of progressive depression. In such cases the gastro-intestinal mucous membrane was found highly inflamed, and the intensity of the irritation probably paralyzed the intestinal functions. According to Daraszkiewicz and to Schaur, in order for gambogic acid to act as a purgative the presence of bile in the intestine is necessary. As it has been determined by Schaur (quoted by Husemann, p. 754) that the hypodermic administration of gamboge to dogs results simply in the production of local abscesses, and as A. L. Richter asserts that when applied to raw surfaces in man it acts simply as a local irritant, it would appear at first sight that gamboge does not act by absorption. This is, however, probably not true, the alkaline juices of the alimentary canal probably dissolving it so that it can readily be taken up by the villi. Further, both

Gmelin and Tiedemann assert that they have found its principles in the urine. Schaur has, however, been unable to detect it in the urine of persons or of animals taking it. Even when he injected large quantities of it into the blood of dogs, he failed to find it in the urine, although he did obtain a resinoid substance which he believes to be a derivative of gambogic acid. Lewis, Abeille, and Ferriar assert that, when given in certain ways, gamboge acts as a decided diuretic. If this be true, absorption of it must occur.

THERAPEUTICS.—On account of the intense irritation which large doses of it produce, gamboge is very rarely, if ever, used alone as a purgative, but is employed to give sharpness to purgative combinations. In very obstinate *habitual constipation* it has been used in doses of one or two grains as a decided laxative.

Its use as a hydragogue in *dropsy* is to be absolutely condemned, it being much less effective for this purpose, and more irritating, than various other substances.

The dose of gamboge is from two to five grains, made into pill with soap, or given in alkaline solution.

OLEUM TIGLII—CROTON OIL.

The fixed oil obtained from the seeds of the *Croton Tiglium*, a euphorbiaceous shrub of Hindostan and other portions of Southern Asia. This oil is quite viscid, and varies in color from a pale yellow to a dark reddish brown, and has an acid reaction. Its taste is hot, acrid, and extremely persistent; its odor faint, but peculiar. The composition of croton oil is very complex, and the active purgative principle has not as yet been separated. The most important of the substances so far discovered in the drug is that found by T. Schlippe and named by him *Crotonol*. According to this investigator, it is not the purgative principle, although to it the drug owes its power of causing a peculiar dermal inflammation.

PHYSIOLOGICAL ACTION.—Locally applied, croton oil is an intense irritant, producing upon the skin an eruption which is at first papular, but in a very short time becomes pustular. This effect of the drug will be considered more in detail under the heading of Counter-Irritants.

When given by the mouth, croton oil acts upon the horse, dog, and probably other mammals, as upon man, producing violent purging, with severe griping, and, when in sufficient amount, fatal gastro-intestinal inflammation. The question as to whether it acts by producing a simple local impression or by absorption is unsettled. In the experiments of Hertwig (quoted by Stillé, *Therapeutics*, vol. ii. p. 449) and of Bucheim (*Virchow's Archiv*, xii. 1), purgation did not follow the injection of the oil into the veins of animals; but Conwell obtained a result contrary to this, and there is considerable testimony that its external use in man is sometimes followed by its specific effects upon the alimentary canal (see Stillé, *Therapeutics*, 2d ed., vol. ii. p. 451).

The probabilities seem to me to be in favor of the purgation being due to the absorption of the oil.

THERAPEUTICS.—Croton oil is a very rapidly acting, violent drastic and hydragogue cathartic. It is chiefly used in cases of *obstinate constipation* from disease of the nervous system or from lead-poisoning. The fact that a drop of it placed upon the tongue will purge actively, peculiarly fits it for use in *mania, delirium tremens*, and other diseases when the patient refuses to take medicine. When it is desired, as in some brain-diseases, to revulse through the bowels, croton oil is probably the most available of the cathartics. The dose is one drop, which may be administered in pill, emulsion, or by simply placing upon the tongue. In overdoses, croton oil is a violent poison.

TOXICOLOGY.—Although in small amounts croton oil produces such severe symptoms, yet in larger quantities it has failed to produce as serious results as would be naturally expected. Cowan has recorded a case (Husemann, *Toxicologie*, Bd. ii. p. 443) of a child, four years old, who recovered in two days from a teaspoonful of croton oil taken on a full stomach. Adams (*Ibid.*) saw recovery in an adult after the injection of a drachm. The only fatal cases I have met with are one reported by Orfila, in which two and a half drachms produced death in a man who had already been sick four weeks with typhus fever, and one recorded by Giacomini (Stillé, *Therapeutics*, vol. ii. p. 451), in which twenty-four grains of the drug proved fatal in as many hours: although there were but four stools, the patient presented the symptoms of general collapse, preserving consciousness to the last.

HELLEBORUS NIGER.—Black hellebore is the root-stock of the *Helleborus niger*, a native of Southern and Middle Europe. It contains two active medicinal principles, *Helleborein* and *Helleborin*. Both of these substances are glucosides. According to Marmé (quoted by Husemann), helleborein is an active cardiac poison and a drastic purgative, whilst helleborin, although having to some extent the latter property, acts chiefly as a narcotic, producing in animals at first inquietude, soon followed by paresis both of motion and of sensation, constantly deepening after large doses into death.

Black hellebore when taken by man is said to produce, if in sufficient quantity, violent catharsis, with vomiting, abdominal pains, cramps, and convulsions, which have ended fatally. Locally applied, the fresh root is a violent irritant. It has been used in times past as a hydragogue cathartic, but appears to be both uncertain and very harsh in its action, and, in this country at least, is never used at present. The *extract* (*Extractum Hellebori*) and the *tincture* (*Tinctura Hellebori*) are official; their purgative doses are respectively ten grains and two fluidrachms.

CLASS III.—DIURETICS.

DIURETICS are medicines used for the purpose of increasing the flow of urine. Some of them, without doubt, act directly upon the secreting structure of the glands, but others of them induce the increased secretion indirectly, by, in some way, removing the obstacle to secretion. It is notorious that diuretics often fail in practice when their action is most urgently needed. This results, in many cases, from the nature of the disease, and is not because diuretics are powerless or uncertain. Thus, in cardiac disease the congestion of the kidneys may be so great as to render secretion impossible; and it is equally evident that when the tubules are destroyed by Bright's disease medicine must be powerless to provoke excretion.

There are certain agencies whose influence upon the kidneys should never be lost sight of in exhibiting diuretics. Thus, cold, by checking the secretion of the skin, often acts as a most efficient remedy of the class, and, whenever a diuretic action is urgently needed, it should be employed as an adjuvant to the medicinal substances.

Again, mere vascular fullness tends to provoke excretion of water by the kidneys. Hence the taking of large draughts of simple water at regular intervals has been found to act very favorably in *acute Bright's disease*, increasing the urinary flow very greatly, and at the same time lessening the irritation of the kidneys, acting, as it were, as a demulcent to the inflamed organs. In various inflammations or irritations of the genito-urinary organs, as in gravel, whenever it is desired to make the secretion less irritating or less concentrated, the value of water as an adjuvant to medicinal diuretics should always be taken advantage of.

There certainly is a very marked antagonism between the bowels and the kidneys, so that free catharsis reduces very decidedly the secretion of urine. There is also an antagonistic relation between the skin and the kidneys, so that an increase in the excretion from one of these generally results in a diminution of that of the other excretory. This should also be taken advantage of when a diuretic action is desired. Sweating and purging at such times are therefore to be avoided. When a diuretic is exhibited, the patient should be kept cool, walking about if able, or if it is necessary for him to remain in bed he should be covered lightly. Not rarely, a remedy which when administered cold and the patient kept cool afterwards will act as a diuretic, when given hot and the patient kept warm will act as a diaphoretic.

There are various substances which are of such nature that when eliminated by the kidneys they act upon the mucous membrane of the bladder and other surfaces over which they pass. It seems hardly correct to speak of these drugs as diuretics; yet they are best considered in the present class.

The chief indications for the use of diuretics are as follows:

1. *To maintain the action of the kidneys.* It is hardly necessary here to discuss the necessity to the system of excretion. In various kidney-diseases this indication is very urgent; but, as the lessened excretion too often depends upon a profound organic alteration of the renal secreting structure, it is evident that very frequently diuretics must fail when most needed. In the great majority of cases in which diuretics are used to fulfill the present indication, only the mildest of the class should be employed. Whenever there is inflammation of the kidneys, even if it be chronic, the stimulating diuretics should be avoided.

When lessened urinary excretion is purely functional in its origin, diuretics are often most serviceable. In fevers especially is it necessary to maintain the action of the kidneys; for this purpose water should always be freely given during fever. The alkaline diuretics sometimes may be exhibited; but the most generally serviceable of all remedies of the class in these diseases is the sweet spirit of nitre.

2. *To evacuate fluid.* For this purpose diuretics are employed in all forms of dropsy, and are successful in direct proportion to the universality of the effusion and the structural perfection of the kidneys.

3. *To soothe and diminish irritation of the genito-urinary organs.* The value of water in fulfilling this and the next indication has already been pointed out. By lessening the acidity of the urine and rendering soluble the uric acid which is present, the alkalies are equally important in carrying out the present and the following indication.

4. *To alter the urinary secretion so as to prevent the deposition of calculous material.* Notwithstanding it has been claimed otherwise, I think it indubitable that as yet no practicable measure has been devised of dissolving a calculus when once formed. Even to alter the urine so as to prevent further deposition is probably impracticable, except in cases of uric-acid diathesis. A discussion of the use of diuretics for this purpose will be found in the article on Potash, which is the only diuretic used to meet the present indication.

Diuretics are very naturally divisible into three sets,—the hydragogue diuretics, the refrigerant diuretics, and the stimulating diuretics. These classes, of course, run more or less into one another, but are sufficiently distinct for practical purposes. The drugs belonging to the first set simply increase the flow of water from the kidneys, and are therefore used chiefly for the relief of dropsy; those of the second division exert a marked sedative action upon the system,—very generally do not increase to a great extent the water of the urine, but mostly modify the secretion in one way or the other. They

are used in dropsy to alter the urinary secretion, and for their sedative and eliminative action in acute disease. Diuretics belonging to the third division are of such nature that their active principles are eliminated by the kidneys, and act upon the mucous surfaces over which they pass; for which purpose they are chiefly employed.

HYDRAGOGUE DIURETICS.

SCILLA—SQUILL.

The bulb of *Scilla maritima*, a liliaceous plant growing in the south of Europe, especially on the shores of the Mediterranean. The bulb varies in size from that of a child's head to that of the fist. It is composed of numerous layers or scales, which separate when it is sliced for drying. As kept in the shops, squill is in horny flakes, of a white or red color, becoming leathery when wet, and having an acrid bitter taste. It yields to water and alcohol, and also to vinegar. *Scillitin* has been asserted to be the active principle of squill; but the *scillitin* of different authors is diverse. According to Reil, there are two active principles in squill, one which he names *scillitin*, representing the diuretic and expectorant properties of the drug, whilst the toxic and irritant properties reside in a substance which he calls *sculein*. Righini believes that the squill contains veratria. Only one conclusion is to be drawn from the evidence, namely, that we are as yet ignorant as to the nature of the active principle of squill.

PHYSIOLOGICAL ACTION.—According to the experiments of Chateau (quoted by Stillé) and of Marais, squill in poisonous dose produces in dogs and other of the lower animals vomiting, then purging, dullness, stupor, intermittent paralysis, convulsions, and finally death in the course of twelve or fifteen hours. The temperature always falls.

Recently, Prof. Schroff (*Wochenblatt der Zeitschrift der k. k. Gesellsch. zu Wien*, 1864, p. 424) has reinvestigated this subject, using alcoholic extracts of the red and the white squill, and also the scillitin of Merek. Fifteen grains of the latter caused in a vigorous rabbit great weakness, mydriasis, and, after an hour or so, tremors gradually becoming violent, partial stupor, labored breathing, and finally death; twenty-three grains caused in another rabbit sinking of the pulse and respiration-rate, mydriasis, diuresis, and death, preceded by the other symptoms noted in the previous case. The alcoholic extract (fifteen grains) caused (*loc. cit.*, 424) in a large rabbit decrease of the number of respirations per minute, with rise of the pulse-rate, narrowing of the pupil, semi-stupor, and finally death. On post-mortem examination of rabbits killed with the scillitin, erosion of the gastric mucous membrane, pericardial and sub-pleural hemorrhages, pulmonary apoplexy, bloody urine, and hyperemia of the kidney and brain were found. In rabbits destroyed by the extract, gastric erosion and the various hemorrhages were wanting.

It seems evident that the scillitin of Merck does not represent squill, and also that we are still in the dark as to how the drug in poisonous doses acts upon the system.

Clinical experience has established the fact that in small repeated doses squill is diuretic as well as expectorant. The remedy is evidently a stimulant to the kidneys, and in overdoses causes an irritation whose result is lessening of the secretion, scanty bloody urine, or absolute suppression of urine, according to the ingested dose of the poison.

Its diuretic action has been noted in animals by Schroff and by Chiarenti (quoted by Stillé), and there can be no doubt as to the power that squill has of increasing the watery portion of the urine. No studies have, that I am aware of, been made as to its action on the urinary solids.

That the active principles of squill are absorbed is proven not only by its action on the kidneys, but also by the fact that its characteristic effects on the system have been seen to follow its external application. (See Stillé, *Therapeutics*, 2d ed., vol. ii. p. 534.)

In man, overdoses of squill produce violent vomiting, colicky pains, purging, and diminution or suppression of the urine, which may contain blood.

THERAPEUTICS.—As a diuretic, squill is in great repute, and is especially employed in cases of dropsy where the condition of the system is atonic and where there is no disease of the kidney. Profs. Wood and Chapman recommend it very strongly in cases of *serous effusion* into the *pleura* or the *pericardium* dependent upon chronic inflammation of the membrane. In these cases it may often be advantageously combined with calomel. The combination of squill and digitalis is very efficient in *cardiac dropsy*. The one contra-indication to the use of squill is the existence of any form of Bright's disease or of acute irritation of the kidney.

TOXICOLOGY.—Overdoses of squill produce violent purging and vomiting, with abdominal pain, lessened or almost suppressed secretion of bloody, albuminous urine, very marked reduction of the pulse-rate, ending, it may be, in collapse, convulsions, and death.

According to Husemann (*Toxicologie*, Bd. i. p. 413), twenty-four grains of it have brought about a fatal result.

The treatment consists in the evacuation of the stomach and bowels by ipecacuanha and castor oil, if nature has not already fulfilled the indication; the free use of opium; the exhibition of large quantities of water, for its action on the kidneys; and the usual measures for the relief of gastro-enteritis, if much tenderness be present. Early in the poisoning care should be exercised in the exhibition of alcoholic stimulants, for fear of increasing the gastric irritation; during the stage of collapse they may be imperatively demanded, and with their use should be combined that of dry heat applied externally, and of the other usual measures of relief during collapse.

ADMINISTRATION.—As a diuretic, squill should be given in solid form, two grains every two hours, the dose being gradually increased until some

nausea is felt. The official preparations of squill are the *Tincture* (*Tinctura Scillæ*, $\S$ ii—Oj.),—dose, $\mathfrak{m}_x$ to $\mathfrak{m}_{xx}$; the *Vinegar* (*Acetum Scillæ*, $\S$ ii—Oj.),—dose, $\mathfrak{m}_x$ to $\mathfrak{m}_{xx}$; the *Syrup* (*Syrupus Scillæ*),—dose, fʒss; and the *Fluid Extract* (*Extractum Scillæ Fluidum*),—dose, $\mathfrak{m}_{ii}$ to $\mathfrak{m}_v$.

DIGITALIS, in its general relations, has already been sufficiently discussed, and it remains only to speak of its employment as a diuretic. In the first place, it should be distinctly understood that it has no alterative effect whatever, either upon the nature of the secretion or upon the mucous membrane over which that secretion flows. In other words, when it has any effect, it is purely a hydragogue diuretic, simply increasing the watery portion of the urine. That digitalis has direct diuretic properties cannot, I think, be doubted. Nor does it seem less certain that it varies greatly in their exercise, so that when given to persons in health it will sometimes produce free diuresis and will at other times fail to do so. Another point to be constantly borne in mind during its administration is the fact that, like all the other actions of digitalis, diuresis is very slowly induced, and is very persistent when produced by the ordinary cautious method of administration. The diuresis of digitalis is not simply a result of its action on the circulation, since it will sometimes appear before the circulation is sensibly affected. At the same time, it is very evident that in disease the good effect of digitalis upon the renal organs is often in large measure due to its action upon the heart. Thus, in dropsy from a dilated heart the renal gland-cells cannot act because they are not supplied with the proper kind and quantity of blood, because their circulation, like that of the remainder of the body, is nearly stagnant. If under these circumstances digitalis be exhibited, and the circulation becomes comparatively free and active, the result is diuresis wrought out through a double mechanism, partly indirectly and partly directly produced by the drug. As a consequence of these facts, the clinicians have long since practically determined that digitalis is especially valuable as a diuretic in *cardiac dropsy*.

Digitalis is, however, also very useful in renal dropsy, both in the subacute and the chronic form. Of course, like every thing else, it frequently fails in these varieties of Bright's disease, but certainly it should always be tried. Prof. George B. Wood asserts that he has seen cases of "what appeared to be decided and obstinate attacks of Bright's disease, with universal dropsy, and unconnected with scarlatina, which yielded completely and permanently to the use of digitalis."

In *acute suppression of urine*, digitalis is often a very valuable remedy when applied externally, especially when the stomach refuses to retain medicines. At the same time, it should be remembered that large doses of the drug used in this way sometimes induce very alarming symptoms. Dr. E. F. Fanell (*British Medical Journal*, March 11, 1871) details a case in which almost fatal collapse was produced by the external use of an ounce of the tincture in a case of renal dropsy.

ADMINISTRATION.—The dose of the powder of digitalis, as a diuretic, is three grains a day (in divided doses), increased by a grain every second or third day, until some sensible effects are manifested. The infusion or the tincture may be substituted for the powder, in corresponding dose. Digitalis, in the majority of cases, is best given in combination: in cardiac dropsy it is much more efficient if given with squill; in renal diseases the bitartrate of potassa may be exhibited simultaneously. Whilst our present knowledge of the physiological action of digitalis is in accord with the ascertained clinical fact that it is safe to give the drug cautiously in the last stages of cardiac exhaustion, yet it should never be forgotten that, as a diuretic, digitalis sometimes refuses to act, and that it is possible to produce the most profound depression with it without inducing the desired result. It is, therefore, worse than useless to persist with the medicine to the danger of the patient after its constitutional effects have been distinctly produced and no diuresis has occurred.

The diuretic external use of digitalis is made by putting a poultice of an ounce of the fresh leaves upon the abdomen of the patient, or, preferably, flannel cloths wrung out of the infusion may be applied to the same part and covered with oiled silk, or half an ounce to an ounce of the tincture may be sprinkled upon previously-moistened spongio-piline. In either case the application should not be allowed to stay on for more than eight hours, at the expiration of which period it should be removed, to be replaced at the end of six hours if no effect has been produced.

SCOPARIUS.—*Broom* is the dried tops of *Cytisus Scoparius*, or the common broom-plant of Europe, cultivated in, and in some places escaped from, the gardens in this country. The drug occurs as greenish pentangular twigs with minute downy leaves, having a bitter nauseous taste, and, when bruised, a peculiar odor. Two principles have been discovered by Dr. Stenhouse in scoparius, *Scoparin*, a neutral crystallizable body, to which, he believes, the drug owes its diuretic properties, and a liquid alkaloid, *Sparteïn*. This alkaloid appears to influence the nervous system; although the testimony concerning it is discordant. According to Husemann (*Die Pflanzenstoffe*, p. 64), Mitchell found that about four grains of it administered to a rabbit caused death in three hours, preceded by a very short stage of excitation and then deep sleep without convulsions, whilst in Schroff's experiment a single drop caused in the rabbit violent convulsions, followed by a stage of muscular weakness, depression of the heart's action, renewed convulsions, and finally death. In an elaborate examination of the action of large doses of sparteïn (*Archiv für Experiment. Pathologie*, 1873), Dr. Fick arrives at the following conclusions as to the influence of poisonous doses of sparteïn: 1. So far as can be ascertained, sparteïn affects the intellectual functions both of frogs and the mammalia, and can therefore to a certain extent be regarded as a narcotic. But this effect upon the brain is not a very marked one in degree, since, even

in cases where the most decidedly poisonous results have been produced, entire suspension of consciousness has not been observed. 2. Spartein exhibits powerful toxic effects upon the spinal cord, the reflex action of which in particular is lessened to a great degree. 3. Spartein paralyzes the motor nerves, which after a larger dose of the poison lose their electrical excitability entirely. 4. A small dose of spartein causes the suspension of the electrical excitability of the vagus, so that its excitation produces no interfering influence upon the motions of the heart. In larger doses the governing centres themselves are paralyzed, so that neither by the use of muscarin nor by other means can a diastolic cessation of the heart be produced. 5. In the poisoning of mammalia by spartein, death is dependent upon paralysis of the respiratory centres. By employing artificial respiration, life may be prolonged in the poisoned animals for some time.

In very large doses, scoparius produces in man free purging, and even vomiting; but as ordinarily used it is simply a most efficient hydragogue diuretic. It is very largely used in general *dropsy*, and is one of the most reliable remedies of the class, seldom failing unless the structural lesions are such as to prevent any diuretic from acting. It is best given in decoction: half an ounce of the tops in a pint of water boiled down to half a pint. Of this an ounce may be given every three hours until some effect is produced; or a fluid extract, which is not officinal, may be given in half-drachm doses.

SPIRITUS ETHERIS NITROSI—*Sweet Spirit of Nitre* will be considered in the class Diaphoretics, in detail. Suffice it for the present to state that when given in a single large dose (a teaspoonful to a tablespoonful) and the patient afterwards kept cool, sweet spirit of nitre acts as a pretty efficient diuretic, increasing the watery portion of the urine, but not to such an extent as to render the drug available for use by itself in dropsy. It acts upon the kidneys as a mild, soothing stimulant, and is mostly employed as an adjuvant to more powerful diuretics, or by itself when there is simply diminished renal excretion of functional origin or when the kidneys suffer from slight congestion, as shown by aching in the loins without other more serious symptoms.

REFRIGERANT DIURETICS.

POTASSIUM. (K.)

The salts of potassium, like the substance itself, are very poisonous to the lower animals. According to Dr. Paul Guttman, they are all exactly alike in the character and the intensity of their action; but further experimentation is wanting before this point can be considered as decided, and I have preferred to study the bromide entirely separate from its congeners.

In the experiments of Podocæpow (*Virchow's Archiv*, 1866, Bd. xxxv. p. 460) it was found that one cubic centimetre of a solution of the chloride

(one to five), given to a frog by the stomach, would in eight minutes cause abolition both of voluntary and reflex movements. After from fifteen to twenty minutes, cardiac arrest occurred. Upon mammals similar results were obtained, but the abolition of motility was apparently not so profound as in the frog. Thus, four or five drachms of the chloride of potassium injected into the stomach of the dog caused bloody stools, reduction of temperature, muscular weakness, and death without convulsions. Although in most, if not all, of the reported cases of poisoning by potash salts, the most prominent symptoms are those due to the local action upon the alimentary canal, yet it would seem that poisonous doses act upon man as upon other mammals, as great feebleness of pulse and lowering of temperature have been noted as constant phenomena.

The most marked action of the potash salts is upon the heart. When a frog is killed by a salt of potash, the heart* is arrested in diastole, according to both Podocæpow† and Guttman. The blood-current before death is greatly lessened in force, as was determined by Podocæpow by watching the circulation in the web, and by comparative experiments in regard to the rate at which blood flowed from wounded vessels in poisoned and in unpoisoned frogs.

In mammals, potash influences the heart even more markedly than in cold-blooded animals. Ten or fifteen grains of the chloride quickly injected into the jugular vein suffice to produce instant cardiac death. The action of a poisonous dose of potash upon the heart appears to be a local one.

Traube found that when death in the dog was produced by injection into the jugular the heart-muscle failed entirely to respond to electricity. In this case, however, the heart received at once the full dose of the poison, and the careful experiments of Podocæpow and of Guttman have shown that when the potash is introduced gradually and in the more ordinary methods into the circulation, the contractility of the cardiac muscle, although very much impaired, is not at the time of death entirely destroyed; in frogs it is less affected than in warm-blooded animals. Guttman has found that previous section of the vagi has no influence upon the action of large doses of potash. The diastolic arrest is therefore independent of the inhibitory system, and is due to a direct impression upon the cardiac muscle.

Although the effects of large doses of compounds of potassium appear to be made out, definite knowledge is still wanting in regard to small doses.

Traube (*Gesammelte Beiträge*, Bd. i. p. 386) asserts, as the result of his

* The poisonous influence of potash upon the heart was, I believe, first discovered by Black (*Comptes-Rendus*, 1839), and has been confirmed by Bouchardat (*Annuaire de Thérapeutique*, 1844), by Grandeau (Robin's *Journal de l'Anatomie*, 1864), by Rabuteau, (*L'Union Médicale*, 1871, and by others).

† *Vicchow's Archiv*, Bd. xxiii. It is proper to state that Podocæpow states, on p. 506, that the arrest is sometimes systolic, sometimes diastolic, although on p. 511 he asserts that the arrest never occurs in systole.

experiments, that, injected into the blood in doses of two or three grains, the nitrate of potassium acts like digitalis, producing a fall in the pulse and a rise in the arterial pressure; and that if now the vagi be cut, both the pulse and the arterial pressure are enormously increased. Further, after section of the pneumogastrics, small doses of the nitrate produced a fall in the pulse, with increased arterial pressure; but on a repetition of the dose in the same animal, no lessening of the pulse-frequency was perceptible, while each time the pressure rose. Opposed to these results are the experiments of Podocarpow (*loc. cit.*, p. 515), who found that the increased arterial pressure produced by two grains of the chloride of potassium lasted only a minute or two. After larger doses of the potassium salts, according to Guttman, section of the vagi causes only a very brief rise in the frequency of the pulse.

The action of the drug upon the motor system is more marked in cold- than in warm-blooded animals. Podocarpow believes, but does not definitely prove, the paralysis of both voluntary and reflex movement to be of muscular origin. The much more elaborate experiments of Guttman show that the muscles of poisoned frogs are not only excitable at the time of death, but are nearly as sensitive as normal muscles, and maintain their excitability nearly as long after death. As both nerve-trunks and muscle are capable of performing their functions in the dying frog, Guttman concludes that the paralysis is of spinal origin, a conclusion which he confirmed by tying the aorta directly above its bifurcation, and then administering the potash, when the paralysis appeared as early in the protected hind legs as in the non-protected front ones.

Outside of the body, potash favors very greatly the oxidation of organic substances. Thus, when olein is exposed to ozone no change occurs, but if potash be added rapid oxidation follows. Whether a similar influence is or is not exerted within the body is not as yet completely determined, but the evidence so far as made out tends to show that it is. The fall of temperature produced by poisonous doses of potassium salts would seem to indicate lessened oxidation, but is probably simply the result of the profound depression of the circulation.

The chief arguments in favor of the theory of increased oxidation in the system as yet brought forward have been drawn from studies of the action of the drug upon the urinary excretion in health and in disease. Potash or its salts administered in sufficient quantity certainly, under ordinary circumstances, increase the watery portion of the urine; but, as Golding Bird has demonstrated, they do more. He found (*On Urinary Deposits*, second American edition, 1859, p. 356) that in a case carefully tested, under favorable circumstances, three drachms of the acetate of potassium increased the solids of a dog's urine from four hundred and sixteen to seven hundred and eighty-two grains, or, deducting all the eliminated potash, to over six hundred grains. The increase of the uric acid was about thirty-two per cent.; of the urea, about sixty per cent.; extractives, including creatine, creatinine,

etc., about twenty per cent.; or, speaking absolutely, the uric acid was increased eighty-five grains, the urea seventy-two grains, and the extractive thirty-six grains. Dr. Austin Flint (*American Medical Monthly*, Oct. 1860) has studied the effect of the nitrate of potassium upon a number of persons suffering from various diseases, and found that it very greatly increased the amount of solids in the urine. Rabuteau (*L'Union Médicale*, 1871, p. 389) has found that the daily ingestion of seventy-five grains of the chloride of potassium caused an increase of twenty per cent. in the amount of urea discharged.

Rabuteau, in his experiments with the chloride, found that the urine maintained its acidity. It is notorious, however, that large doses of the acetate, carbonate, or citrate of potassium produce alkalinity of the urine. The explanation of the apparent contradiction is that the vegetable salts are destroyed in the system and eliminated as alkaline carbonates, whilst the nitrate, and probably chlorides, sulphate, and similar compounds, pass entirely, or in great part, unchanged through the body. A proof of the latter fact is furnished by Prof. Alfred S. Taylor (*Guy's Hospital Reports*, 1863, p. 177), who, from the urine of a patient taking two hundred and seventy grains of the nitrate daily, obtained 158.7 grains of the ingested salt per diem. A portion of the potash salts escapes through the intestines, as Dr. Kramer (*Annales d'Hygiène Publique et de Médecine Légale*, vol. i., 1843) has found the nitrate in the feces of animals taking it; and it is much more probable that the nitrate not accounted for in Dr. Taylor's investigation was eliminated by the intestines than that it was decomposed in the system.

If, as there is much reason to believe, a vegetable acid when given alone passes through the system in great measure unchanged, whilst, as asserted by Dr. Münch (*Archiv des Vereins für gemein. Arbeiten*, 1863, p. 370), and as seems to follow from the facts already brought forward, the same acid is found when combined with an alkali to be destroyed by oxidation, there is in this strong reason for believing that the alkalies aid oxidation in the system. The oxidation-theory has been recently strenuously upheld by, if indeed its application did not originate with, certain clinical observers, especially Dr. Basham (*London Practitioner*, vol. v., 1870).*

It has long been customary to use potash salts in excess of uric acid in the urine, and the relief obtained has been believed to be due to the conversion of the acid into a urate. Dr. Basham affirms, however, that as the result of

* Dr. D. Campbell Black, in a letter to the *London Lancet* of September 20, 1873, attempts to correct an assertion which, according to his statement, Dr. Harley made before the British Association, that Dr. Basham was the originator of the oxidation-theory, claiming for himself priority. Unfortunately, Dr. Black misquotes dates in a most hopeless manner, none of his references being anywhere near correct. I have not seen the pamphlet in which Dr. Black claims to have advanced the theory, and cannot give a positive opinion; but, so far as I can make out, Dr. Basham at least thought out the theory for himself, and, what is more to the point, furnished the facts which give it some other than a speculative basis.

a series of analyses he has found that in uric acid diathesis not only is there a great increase of the urea, but that the uric acid, either free or combined, in the urine is greatly diminished. Dr. Basham, remembering that Mr. Schunck had proved that, under the oxidizing power of potash, uric acid outside of the body is converted into oxaluric acid, which in its turn is readily metamorphosed into oxalic acid and urea, carefully examined the urine of gouty patients taking the potash, and found that not only was the urea increased, but that oxalic acid also appeared as the uric acid decreased, and that, on standing, the urine deposited crystals of oxalate of calcium, although none of these could be found in it when first voided.

This research of Dr. Basham certainly seems to demonstrate that in uric acid diathesis the potash salt increases the oxidation and the ultimate metamorphosis of tissue. Where this occurs, whether in the blood, in the kidney, or in the urine itself, is not at present determined.

When a potassium salt is given in large doses for a long time, it produces a condition of dyscrasia, with impoverishment and excessive fluidity of the blood. How or why it has this action is unknown, as indeed is the exact nature of the changes. Very probably there is some connection between these changes and the oxidizing power of the drug; but any theory in the present imperfect state of our knowledge could at best be only a useless speculation.

Our knowledge of the physiological action of the potassium salts seems to show that the vegetable salts and the carbonates are equivalents, but that the mineral salts are more or less peculiar and individual; and clinical experience confirms this. There is, however, one exception: the bitartrate of potassium seems to act differently from the other vegetable acid salts, and, although direct proof is wanting, probably is not decomposed in the system. Potash itself is never used to affect the system, on account of its irritant properties; and its local action will be discussed under the headings of Escharotics and Antacids. I shall here group all the potassium vegetable salts, except the bitartrate, together: considering them first, then the bitartrate, and then the salts of the mineral acids.

POTASSII CARBONAS.

The potash of commerce, obtained from wood-ashes and other sources, occurs in the form of fused, stony masses, variegated in color, and of a caustic burning taste; when purified so as to form *pearlash*, it becomes of a bluish white, and constitutes the officinal *Impure Carbonate of Potash* (POTASSII CARBONAS IMPURA), which should contain seventy per cent. of the hydrated alkali, and one hundred grains of which should neutralize fifty-eight grains of officinal sulphuric acid. It contains phosphate and silicate of potash.

POTASSII CARBONAS.—*Carbonate of Potassium* ($\text{KO} \cdot \text{CO}_2 = \text{K}_2\text{CO}_3$) is prepared by dissolving pearlash in water, filtering, and evaporating. It occurs as a coarse, granular, whitish powder, very deliquescent, soluble in its weight

of water, insoluble in alcohol. It contains traces of the sulphate, the chloride, and more of the silicate of potassium.

POTASSII CARBONAS PURA.—*Pure Carbonate of Potassium* is a white, granular, deliquescent salt, made by heating the bicarbonate of potassium to redness. It should be absolutely pure.

POTASSII BICARBONAS.—*Bicarbonate of Potassium* ($\text{KO} \cdot 2\text{CO}_2 - \text{KHCO}_3$) is manufactured by passing carbonic acid gas through a solution of the carbonate in distilled water. It occurs in transparent, colorless crystals, not deliquescent, slightly alkaline to the taste and to test-paper. It dissolves in four times its weight of boiling water, but is insoluble in alcohol.

The bicarbonate and the pure carbonate contain no silicate of potash, and are therefore useful in making various preparations. Therapeutically they are of equal value, except that the carbonate is more irritant, and is therefore not so well borne by the stomach as is the bicarbonate. On account of its nauseous taste, even the latter salt is not so available as the acetate or the citrate. The full dose of the bicarbonate is half an ounce daily, given in diluted solution.

POTASSII CITRAS.—*Citrate of Potassium* ($3\text{KO} \cdot \text{C}_{12}\text{H}_5\text{O}_{11} - \text{K}_3\text{C}_6\text{H}_5\text{O}_7$). A whitish, granular, deliquescent salt, of neutral or very slightly acid reaction, freely soluble in water. It is the least offensive to the palate of all the potash salts, except the tartrates.

The following preparations of the citrate of potassium are officinal:

LIQUOR POTASSII CITRATIS.—*Solution of Citrate of Potassium* (Bicarbonate of Potassium, 300 grs.; Citric Acid, ʒss; Water, Oss).

MISTURA POTASSII CITRATIS.—*Neutral Mixture* (Lemon-juice, Oj; Bicarbonate of Potassium, enough to neutralize).

These preparations have been long used as diaphoretics in sthenic fevers. The dose is half a fluidounce to one fluidounce every one or two hours.

A very elegant method of exhibiting neutral mixture is in the form of *Effervescent Draught*. It is especially useful when there is any tendency to sick stomach. It should be prepared in two solutions: one consisting of lemon-juice and water, equal parts, or of citric acid ʒii, water fʒiv: the other of potassii bicarbonas ʒi, water fʒiii. An ounce of each of the solutions to be put together, and the whole to be drunk during effervescence.

POTASSII ACETAS.—*Acetate of Potassium* ($\text{KO} \cdot \text{C}_4\text{H}_3\text{O}_3 - \text{KC}_2\text{H}_3\text{O}_2$) is a perfectly neutral white salt, of a decidedly saline taste, extremely deliquescent, and soluble in half its weight of water. It is made by dissolving the bicarbonate in acetic acid, and evaporating. It occurs sometimes as soft, fibrous masses, at other times it has a foliated structure.

THERAPEUTICS.—The most important use of the carbonates and of the vegetable salts of potassium is in *acute inflammatory rheumatism*. It seems to me indisputable that the "alkaline treatment" is the best yet suggested for

thoroughly acute rheumatism: the medicine must be given freely, an ounce to an ounce and a half in the day, and be persisted in, opium, of course, being at the same time employed in as large doses as is necessary to relieve the pain; after a few days, when the violence of the symptoms has abated and decided anæmia appears, the exhibition of the drug should be discontinued and iodide of potassium, with tonics, be substituted. In cases subacute from the beginning I have found a combination of the iodide and acetate of potassium very efficient, ten grains of the former and thirty of the latter being administered three or four times a day. The potash probably does good in rheumatism by lowering arterial action, by favoring oxidation and elimination of partially effete materials, and by neutralizing excessive acidity. Be the method what it may, I have no doubt of the great clinical value of the remedy; its applicability being in direct proportion to the acuteness and violence of the symptoms.

As depuratives, the potash salts are very useful in various diseases. Attention has been especially called by Dr. Golding Bird to their value in that especial class of cases spoken of as "*chronic biliousness*;" in *chronic malarial poisoning*, in *catarrhal jaundice*, and in the *jaundice* of simple *hepatic torpor*, they are often of use.

In *uric acid gravel* and in *uric acid calculus* there can be no doubt of the value of potash as a prophylactic, as a preventive of the formation or deposition of the uric acid. The remedy has also been used to dissolve uric acid *calculi*; but the results offer such slight encouragement that it is only necessary here to give a reference to the work of Dr. Wm. Roberts (*On Urinary and Renal Diseases*, Am. ed., 1866).

ADMINISTRATION.—As usually given, the potash salts are exceedingly distasteful. There is no need of this whatever. The citrate may be given dissolved in lemon-juice, or, what is a still more pleasant method, a syrupy solution of the bicarbonate and the citrate may be made, of such a strength that every tablespoonful of it shall contain half a drachm of each salt. At the time of exhibition one or two tablespoonfuls of this may be put in a little water, and to it be added a large tablespoonful of lemon-juice, the whole to be drunk whilst effervescing. Of course, if the patient takes in the course of the day six of the largest doses mentioned, the whole amounts to an ounce and a half of the citrate of potash. When the remedy is used simply as a depurative, as in jaundice, such large doses are, of course, not proper; a teaspoonful of the alkaline solution, with a corresponding amount of lemon-juice, taken three times a day, will generally be sufficient.

POTASSII BITARTRAS — *Bitartrate of Potassium* ($\text{KO}, 2\text{C}_4\text{H}_2\text{O}_5\text{HO} - \text{KC}_4\text{H}_5\text{O}_6$).—*Cream of Tartar* is made from argol (see Tartaric Acid), and occurs in white crystalline crusts or masses, which are commonly pulverized before being offered for sale as cream of tartar. It usually contains some tartrate of lime, and is only sparingly soluble in cold water.

THERAPEUTICS.—The bitartrate of potassium appears to differ from its

congener in being more actively diuretic, and in acting more powerfully as a hydragogue cathartic. Half an ounce to an ounce of it given at once will very generally cause watery purging. In this city it is probably employed more frequently in *dropsy* than any other diuretic: the usual plan is to dissolve an ounce of it in a pint of infusion of juniper-berries, and have this all taken, in divided doses, during the twenty-four hours.

In acute *desquamative nephritis*, cream of tartar is a very useful diuretic: as, however, the avoidance of irritation of the kidneys is imperative in this disease, the alkaline diuretic should not be administered in infusion of juniper.

The *Tartrate of Potassium* (POTASSII TARTRAS) is officinal, but is rarely used in medicine. It is said to be actively purgative in doses of half an ounce.

POTASSII SULPHAS—SULPHATE OF POTASSIUM ($\text{KO},\text{SO}_3 - \text{K}_2\text{SO}_4$).

This salt occurs in the form of small, aggregated, transparent, very hard crystals, permanent in the air, having the shape usually of short six-sided prisms, and possessing a nauseous somewhat bitter taste. It is insoluble in alcohol, slowly soluble in nine and a half times its weight of cold and in less than four times its weight of boiling water. Sulphate of potash is said to be "a mild purgative, operating usually without heat or pain or other symptoms of irritation," in doses of four or five drachms; and in doses of one or two drachms acting as a laxative. It is, however, very rarely employed in this country as a purgative. Sulphate of potash, in doses not a great deal in excess of those which have been recommended by practitioners, acts as an irritant. Dr. Mowbray states that the salt is used in France as a popular abortifacient, and that he has seen very alarming symptoms produced by four drachms of it. Dr. Taylor records a case in which less than two ounces caused in a woman severe vomiting, purging, abdominal pain, and finally death. At the post-mortem the stomach and intestines showed very decided evidences of inflammation. The chief use of sulphate of potash in this country is as an ingredient of Dover's powder.

POTASSII NITRAS—NITRATE OF POTASSIUM ($\text{KO},\text{NO}_3 - \text{KNO}_3$).

Nitre is either obtained from certain saline earths, occurring chiefly in India but to a certain extent in other portions of the world, or else is artificially manufactured in nitre-beds formed out of animal and vegetable matter, wood-ashes, and calcareous earth, or, finally, is obtained from old plaster rubbish. In the "nitre-beds," as well as in the natural saline earths, which have undoubtedly in the beginning contained animal and vegetable matters in a state of decomposition, nitric acid is formed by the oxidation of ammonia, and unites with the bases in the soil. Most of the nitre used in this country comes from Calcutta, through Boston, packed in grass-cloth bags. *Chili saltpetre* is the *nitrate of calcium*, which impregnates certain soils in the

country whose name it bears. It is undoubtedly formed in these soils by a process precisely analogous to that in which the nitre of India is produced, excepting that, little or no vegetable matter being present to afford the potassium during the decomposition of the animal matter and the generation of nitric acid, the latter unites with the calcium of the soil.

Chili saltpetre is employed as a substitute for true saltpetre in the manufacture of nitric acid, and, it is said, may be made into the nitrate of potassium by means of crude caustic potash.

Saltpetre occurs in more or less perfect, long, striated, semi-transparent, six-sided prisms, with dihedral summits; of a sharp, saline, somewhat cooling taste, containing no water of crystallization, but decrepitating when thrown on the fire, from the evaporation of water mechanically retained in the crevices of the crystals; soluble in four or five times their weight of cold and in two-fifths of their weight of boiling water, sparingly soluble in proof spirit, insoluble in absolute alcohol. At a high heat they decompose, liberating a large quantity of nascent oxygen, and thereby greatly intensifying the combustion of surrounding objects.

The *Sal prunelle* of the shops is a saltpetre which has been fused and run into circular moulds.

PHYSIOLOGICAL ACTION.—Nitrate of potassium applied to any raw surface, or to a mucous membrane, acts as a violent irritant. As death has not infrequently resulted from its ingestion, whilst it has occurred very rarely from that of any of the other ordinary salts of the alkali, it has generally been thought that nitre possesses peculiar properties. It is not to be gainsaid, however, that the cause of death in nitre-poisoning is very generally the local inflammation of the stomach and intestines produced by it,—effects dependent simply upon its irritant properties, and not upon any constitutional action; a conclusion becoming doubly evident when it is remembered that, if the drug be given in weak solution, much larger amounts can be exhibited with only therapeutic effects, than would cause death if administered in solid form or in very concentrated solution.

Thus, in a case under the care of Dr. Wilks (*Guy's Hospital Reports*, vol. ii., 3d series, 1863, p. 173), a man suffering from renal dropsy took, between October 28 and December 26, 1862, one pound twelve ounces and six drachms of the nitrate of potassium, with benefit. As one ounce has caused death in three hours (Taylor, *Principles and Practice of Medical Jurisprudence*, 2d ed., vol. i. p. 237), this patient received in fifty-nine days the equivalent of twenty-eight fatal doses. Again, according to Prof. Stillé (*Therapeutics*, vol. ii.), Dr. Brocklesby habitually prescribed one ounce of the salt a day, and Dr. Martin-Solon even two ounces per diem.

The symptoms of poisoning by the nitrate of potassium are pretty constant, and yet, as in other irritant poisoning, vary within certain limits. Very generally there is first an intense burning pain in the stomach, coming on in a few minutes after the ingestion of the poison, and soon followed by violent

vomiting, and, it may be, free purging. In a little while collapse develops, with great muscular weakness, not rarely with local convulsive tremblings. The matters vomited, and even the stools, may be bloody (case, Th. Husemann, *Journal für Pharmakodynamik*, 1859, ii. p. 178).

Sometimes the nervous symptoms predominate, and the purging may be absent; collapse, with slight vomiting and with or without paralysis of the lower limbs, may alone exist. Suppression of urine has been noted in some cases (case, *Pharmaceutical Journal*, Feb. 1846, p. 356). After death, very grave lesions are found in the stomach and the intestines, such as intense redness and congestion, effusion of blood into the submucous coat, and sometimes into the stomach itself. Even ulceration and corrosion of the mucous membrane have been observed.

It is evident that the symptoms previously detailed as existing during life are in accord with the post-mortem results, and all point to the irritant action of the drug as the source of trouble.

The predominance of the nervous symptoms in some cases is no more than is exceptionally seen in other forms of irritant poisoning (see Antimony), and is no proof of a special action of the drug upon the nervous system. Sometimes, however, death has occurred, in poisoning by saltpetre, with great suddenness. In the only cases of this character the record of which I have met with, the dose has been very large, and it is possible that the death was the result of the action of the drug upon the heart, for, like the other salts of potash, it has a direct paralyzing influence upon the cardiac muscle.

Nitre has been supposed by practitioners to be especially sedative to the circulation; but there is no reason to believe that it is any more powerful as a cardiac sedative than the vegetable salts of the base. It certainly shows the diuretic properties of the latter, but appears to be more irritant to the kidneys, since it seems difficult otherwise to account for the suppression of urine already noted as occurring occasionally in poisoning by it.

It is possible that nitre may differ to some extent in its actions from the other salts of potash, on account of its being a nitrate and acting upon the blood. (See page 301.)

THERAPEUTICS.—Nitrate of potash has been especially used in *acute rheumatism*, and when given in large doses has some favorable influence upon the course of the affection. It is certainly, however, a more dangerous remedy than the vegetable salts of the base, and, according to my experience, much less efficacious. I can therefore see no good reason for continuing the practice. If given, not less than an ounce should be dissolved in a full *quart* of barley-water or other demulcent, and be administered in divided doses during the twenty-four hours.

In the treatment of poisoning by saltpetre, after the stomach and bowels have been emptied, the usual means for the relief of toxic gastro-enteritis should be resorted to.

POTASSII CHLORAS—CHLORATE OF POTASSIUM $\text{KO.CIO}_3 = \text{KClO}_3$.

This salt is said to be chiefly prepared by heating the solutions of the hypochlorite of lime and chloride of potassium; on cooling, the chlorate of potassium crystallizing out and the chloride of calcium remaining in solution. Chlorate of potassium occurs in white rhomboidal plates of a pearly lustre, of an acerb taste, dissolving in sixteen parts of water at 60°F. , and in two and a half parts of boiling water.

PHYSIOLOGICAL ACTION.—Upon mucous membranes and ulcerated surfaces this salt acts as a powerful irritant, being, I think, even more active in this respect than the nitrate of potassium. Owing to its irritant influence, when taken internally in sufficient quantity it acts as a poison, producing gastro-intestinal inflammation, and, it may be, death. Dr. Fountain, in order to prove the harmlessness of the remedy, took an ounce of it at a single dose: almost immediately copious diuresis occurred, and was followed after a time by persistent suppression of urine and marked symptoms of gastro-intestinal inflammation, with death on the seventh day (*The American Medical Times*, April, 1861, p. 245). Apparently opposed to this are the experiments of Dr. Tully (quoted by Stillé), who on several occasions took an ounce of the salt without injury. These diverse results were no doubt due to a difference in the concentration of the solutions taken, and are exactly parallel with those spoken of in the article on the Nitrate of Potassium.

To the therapist the chief interest in regard to the physiological action of the chlorate of potassium centres in its effects when given continuously. According to Dr. Stevens, when taken internally it causes the venous blood to acquire an arterial hue, and Dr. O'Shaughnessy has observed the same effect when it is thrown into the veins of animals. Probably led by these assertions, many of the profession have given credence to the idea that the chlorate yields its oxygen to the blood. It is, however, evident that even if it be granted that the observations noted are correct, they in no wise warrant such a deduction, since even partial mechanical asphyxia will produce the same phenomenon (see article on Prussic Acid). Viewed from a chemical stand-point, it is very improbable that the salt parts with its oxygen to the blood at the temperature of the body; and, as there is simply no proof whatever that it does so, the proposition must be looked upon as one of the numerous therapeutic myths engendered in the brains of those who find it more easy to speculate than to toil after facts.

As to the real action of the chlorate of potassium in the system, we have very little knowledge, but the salt probably is nearly equivalent to the nitrate of the same base. It certainly increases at times the flow of urine, and is eliminated unchanged and very largely by the kidneys,* but no doubt escapes with all the secretions, since Isambert has found it in the tears, the bile,

* See analyses of O'Shaughnessy, of Wöhler and Stehberger, and of Gustin, all quoted by Prof. Stillé.

the nasal mucus, the perspiration, and even in the milk of nursing women. The same observer noted that within five minutes after its ingestion it could be detected in the saliva, and in ten minutes after this in the urine, in which fluid it continued to be present for from fifteen to forty-eight hours.

In his experiments upon himself, Isambert found that when taken in large dose (two to five drachms a day) the chlorate of potassium caused salivation, free diuresis, increase of the appetite, and, when not well diluted, gastric irritation. The urine continued strongly acid, and contained an excess of rosacic acid, uric acid, and the urates.

THERAPEUTICS.—The chlorate of potassium has been very freely used, and with great asserted advantage, in all forms of disease believed to be due to blood-poisoning: such as *scarlet* and other *adynamic fevers*, *diphtheria*, *scurbutus*, *syphilis*, and even *hydrophobia*. As already stated, the theory upon which this practice rests has no foundation in reason or science, and my own empirical experience with the remedy has been in exact accord with the teachings of physiology. I have seen the chlorate repeatedly employed in various diseases of the class just spoken of, and have never seen it do a particle of good.

On the other hand, in various forms of *stomatitis* the remedy is undoubtedly of great value. Thus, there is a great deal of evidence of its usefulness in mercurial sore mouth; and when given in the form of a powder, with sugar, it almost always acts like a charm in the *follicular* or *aphthous stomatitis* of children. I do not believe, however, that its influence in these cases is other than local; yet, as the remedy is eliminated with the saliva, and, therefore, when given internally is constantly present in the mouth, the ordinary method of using it is probably the best.

The salt now under consideration has been used in *ascites* and other *drop-sical* affections, but is, undoubtedly, inferior to the other salts of the alkali.

When used locally, the chlorate of potassium acts as a stimulant to the various mucous membranes, and is often of excellent service in cases of *angina*, and is even said to have been used advantageously by enemata in *dysentery* and in *cholera infantum*. The most generally efficient gargle that I know of in ordinary *sore throat* may be made by pouring a pint of boiling water over a powder composed of an ounce of *sumach berries** and half an ounce of the chlorate of potassium, allowing to simmer in an earthen vessel, with occasional stirring, to three-fourths of a pint, straining, and using in the ordinary manner.

ADMINISTRATION.—For manifest reasons, when taken in large doses, the chlorate of potassium must be exhibited in dilute solution. The usual dose is from ten to thirty grains; as a lotion, from ten grains to half a drachm may be dissolved in the ounce of water.

* *Sumach berries* are included in the secondary list of the U. S. Pharmacopœia, under the name of the shrub that bears them, *Rhus Glabrum*. They contain very largely of tannic acid, and also of the bimaleate of lime: the malic acid seems to give them an especially beneficial influence upon mucous membranes.

STIMULATING DIURETICS.

BUCHU—BUCHU.

The leaves of the *Barosma crenata*, and of other species of the genus, natives of Southern Africa. These leaves, which are gathered by the Hottentots of the Cape of Good Hope, are an inch or less in length, from three to five lines broad, of various forms, but always notched on the edges, and having a strong, rather rank, yet somewhat aromatic odor, and a warm, bitterish taste. They owe their virtues, which they yield to water and to alcohol, to a volatile oil and to a bitter extractive.

THERAPEUTICS.—Owing to its bitter principle, buchu is perhaps slightly tonic; but its chief medicinal virtue is as a diuretic stimulant and alterative to the mucous membrane of the genito-urinary organs. It does not very largely increase the flow of urine, and hence is never administered in dropsy, but in all cases of subacute or chronic inflammation of the genito-urinary organs it may be employed with hope of success. Its oil is undoubtedly absorbed and is eliminated by the kidneys, to whose secretion it imparts its odor. In *chronic pyelitis*, *chronic cystitis*, and *irritation of the bladder*, it is one of our best remedies, especially when, as is frequently the case, these diseases are associated with a generally-lowered systemic tone. As compared with turpentine, buchu is much less stimulating, and has a far more soothing effect upon the mucous membranes of the genito-urinary tract.

In *irritated bladder*, when the urine is highly acid, and when there is a constant desire to urinate, with but little relief from micturition, buchu, in combination with salt of potash, and perhaps with the sweet spirit of nitre, often gives great relief.

ADMINISTRATION.—The following preparations are officinal:

Infusum Buchu—*Infusion of Buchu* (℞i to ℞j). Dose, one to two fluid-ounces.

Extractum Buchu Fluidum—*Fluid Extract of Buchu*. Dose, half a teaspoonful to a teaspoonful, well diluted with water.

PAREIRA.—*Pareira Brava* is the root of *Cissampelos Pareira*, a climbing plant inhabiting the West Indies and North America. The root occurs in pieces from half an inch to two inches in thickness and from two inches to two or more feet in length; it is cylindrical, simple, or forked, without rootlets, and tightly covered by a thin adherent bark. There appears to be in the root an alkaloid (see *U. S. Dispensatory*, 13th ed., p. 600), to which the name of *Cissampelina* has been given, but concerning which we have but little knowledge.

THERAPEUTICS.—*Pareira Brava* has been used with asserted advantage in chronic *cystitis*, in "*irritable bladder*," and in chronic *gonorrhœa*, and appears to exert a stimulant action upon the mucous membrane of the whole genito-

urinary apparatus. It is said to be also tonic, and slightly aperient, so that it is especially valuable in urinary diseases when there is feebleness of digestion and a tendency to costiveness. The remedy should always be given in the form of the official infusion (*Infusum Pareiræ*, ℥i to Oj) or fluid extract (*Extractum Pareiræ Fluidum*), of which the doses are respectively a wineglassful and a teaspoonful three or four times a day.

UVA URSI.—*Bearberry* is the leaves of the *Arctostaphylos Uva Ursi*, a low evergreen shrub, indigenous to northern maritime Europe, and also growing along our northern coasts as far south as New Jersey. They are from half an inch to an inch in length, wedge-shaped, thick, coriaceous, with a smooth, rounded margin. The odor is hay-like, the taste bitterish, astringent, and somewhat sweetish. *Uva ursi* contains gallo-tannic acid, besides several crystalline principles. *Ursin*, *Arbutin*, and *Urson*, the relations of which are not very well settled.

Mr. J. C. C. Hughes, the discoverer of ursin, claims that it is the diuretic principle of the drug, one grain of it acting powerfully upon the kidneys.

THERAPEUTICS.—*Uva ursi* is capable of acting as a weak astringent, but has been long used in medicine for its influence upon the genito-urinary mucous membrane, and at present is employed for no other purpose. It is stated that gallic acid may be detected in the urine of persons taking it, and it is very probable that other of its active principles are eliminated by the kidneys and exert an influence upon the mucous membranes over which they pass. In chronic *pyelitis*, *cystitis*, and other affections of the genito-urinary mucous membrane, when a slightly stimulant and an astringent action is desired, *uva ursi* may be employed in one of its official preparations, the decoction (*Decoctum Uvæ Ursi*, ℥i to Oj), or the fluid extract (*Extractum Uvæ Ursi Fluidum*), the doses of which are respectively a wineglassful and a teaspoonful three or four times a day.

PIPSISSEWA.—*Chimaphila* is the dried leaves of the *Chimaphila umbellata*, a little indigenous perennial, distinguished from its inert congener *C. maculata* by the uniform glossy green of its leaves. The latter are about an inch and a half long, wedge-shaped, notched, pointed, and coriaceous. They contain tannic acid, bitter extractive, and, according to Mr. Samuel Fairbank, a crystalline principle, *Chimaphilin* (see *U. S. Dispensatory*). Pipsissewa is probably about equivalent to *uva ursi* in its therapeutic value, though perhaps not quite so actively *diuretic*. Prof. Geo. B. Wood (*Therapeutics*, vol. i. p. 133) commends it very highly in external *scrofula*, asserting that he has had a large experience with the remedy, and that in power over the disease it stands next to cod-liver oil and the preparations of iodine and iron. He believes that it acts not only as a mild astringent and tonic but also as an alterative, and states that its exhibition should be long continued, the administration being temporarily suspended whenever there is much fever. The remedy should be administered in one of its official preparations: the decoction.

tion (*Decoctum Chimaphilæ*, ℥i to Oj), and the fluid extract (*Extractum Chimaphilæ Fluidum*), the doses of which are respectively a wineglassful and a teaspoonful three or four times a day.

JUNIPERUS.—*Juniper* is the fruit of the common juniper of Europe and this country. These berries are round, bluish bodies, about the size of a large pea, of a sweetish, terebinthinate, aromatic taste. They owe their properties to a *volatile oil* (*Oleum Juniperi*), which is officinal. They yield to boiling water and to alcohol. Juniper is gently stimulant and cordial to the stomach. Upon the kidneys it exerts a decided stimulant action, and the oil freely given is capable of irritating the renal organs above the secreting-point, and of producing lessened secretion, strangury, and even suppression of urine. The volatile oil is undoubtedly absorbed, and chiefly escapes from the system through the kidneys. As a diuretic, juniper has two distinct uses. The most usual employment of it is as an adjuvant to cream of tartar or the alkaline diuretics. On account of its stimulant local influence upon the alimentary canal, it renders the cream of tartar far more acceptable to the stomach, and at the same time aids its diuretic action. Sometimes juniper is employed for its stimulant action on the mucous membrane of the genito-urinary organs in *chronic pyelitis* and in *chronic catarrh of the bladder*. In the form of the *compound spirit* (*Spiritus Juniperi Compositus*), or its equivalent, *gin.* juniper is often useful in the subacute *congestion* of the *kidneys* often seen in old persons, and characterized by aching in the loins without other more serious symptoms. The officinal *infusion* (*Infusum Juniperi*) is made by macerating an ounce of the berries in a pint of boiling water for an hour, the whole to be taken in divided doses during twenty-four hours.

The dose of the oil is from five to fifteen drops; of the spirit, one tablespoonful.

CAROTA.—*Wild Carrot* of the secondary list of the U. S. Pharmacopœia is the seeds of the *Daucus Carota*, an indigenous umbellifer. These seeds are very small, and have a somewhat aromatic taste, due to the presence of a volatile oil. Carota acts upon the system as a feeble aromatic, and is said also to be possessed of a mild, stimulating, diuretic power.

ERIGERON.—*Fleabane*, according to the U. S. Pharmacopœia, is the dried herbaceous portions of *Erigeron heterophyllum* and *E. Philadelphicum*. These plants contain a very small proportion of volatile oil, and are possessed of but feeble medicinal properties. When given freely in an infusion made with hot water (℥i to Oj,—dose, a wineglassful every two hours), they are said to increase the flow of urine, and to act as stimulant alteratives to the genito-urinary mucous membrane.

Canada Fleabane (*ERIGERON CANADENSE*) contains a large proportion of a yellowish volatile oil of a rather pleasant odor and taste, which is officinal, and is the only form in which the drug should be exhibited. It has properties resembling those of turpentine, but is much less stimulating. It may be

employed in affections of the *genito-urinary organs* and in passive *hemorrhages*. It is especially valuable in *menorrhagia*. The dose is five to twenty drops every two or three hours, and is best administered on sugar.

TEREBINTHINA—TURPENTINE.

Sufficient has already been said concerning the natural history of white turpentine (see page 113); but *Canada Turpentine* (TEREBINTHINA CANADENSIS), or *Canada Balsam*, as it is more commonly called, was not spoken of, because it is never used as a stimulant. It is the product of the *Abies balsamea*, or Balm of Gilead, or American Silver Fir, as it is variously named, a beautiful evergreen indigenous to the extreme northern United States and to the British Colonies. The juice is said to collect in little receptacles under the bark, and to be gathered by cutting these open and allowing them to drain into vessels. Canada Balsam is a thick and viscid, but clear, yellowish liquid, containing, it is said, about twenty per cent. of volatile oil. The amount of the latter ingredient must vary greatly, since by age and exposure to the air the liquid balsam, losing its oil by evaporation and oxidation, becomes converted into a hard, brittle, translucent, resinous mass. Canada Balsam, although officinal, is very rarely, if ever, used in medicine.

Oil of Turpentine, the volatile oil of ordinary turpentine, has already been considered; but it seems necessary to add here a few sentences in regard to its diuretic action and use. When moderate doses (ten drops every three hours) of turpentine are taken, there are usually no renal symptoms produced, except a slight increase of the urine. Somewhat larger amounts, when exhibited, are apt to give rise to aching in the loins and to frequent micturition, with perhaps urethral pain accompanying the act. If still larger quantities are ingested, these symptoms are intensified, and at the same time the secretion of urine is diminished. After very large repeated doses of the drug, the aching in the loins is very great, often with spasmodic pain in the ureters; a constant desire to pass water struggles with the inability to micturate, caused by the urethral spasm; the urine is very scanty, albuminous, and even bloody; priapism may be present, and an intolerable irritation may affect all the pelvic organs. These symptoms produced by overdoses of the drug show that its action upon the urino-genital organs is essentially and powerfully irritant, and that under all circumstances the existence of active irritation or inflammation of these organs absolutely contra-indicates the use of oil of turpentine.

Oil of turpentine is never employed to increase the flow of urine for the purpose of affecting serous effusions. As a diuretic, it is used solely for its local influence upon the organs. *Excessive diuresis* sometimes is apparently dependent upon a relaxed condition of the kidneys, and under these circumstances oil of turpentine may be of great service. *Chronic pyelitis*, *chronic cystitis*, and *gleet* are all often very much benefited by the use of the drug.

In using the remedy in these cases, it should always be borne in mind that, with the exception of cantharides, it is the most actively stimulating of all the diuretics, and must be employed only when such a remedy is called for. In those comparatively rare cases of *urinary incontinence* which are dependent upon debility of the bladder, turpentine is sometimes of great service. When the same symptom is spasmodic, the remedy, of course, is harmful. In absolutely passive *hæmaturia*, in *impotence*, in certain conditions of *spermatorrhœa*, in *amenorrhœa* when great local debility exists, turpentine may be tried with fair hopes of its being useful. The dose of turpentine as a diuretic is ten to fifteen drops in emulsion, given from four to six times a day. If glycerine and oil of gaultheria be added to the emulsion in such proportion that half a teaspoonful of the one and one or two drops of the other be taken with each dose, they will almost completely disguise the taste of the remedy.

COPAIBA—COPAIBA.

The oleoresin of *Copaifera multijuga* and of other species of *Copaifera*, large trees growing in Brazil. Copaiba is a yellowish liquid of varying viscosity according to age, of a strong, terebinthinate, peculiar odor, and a bitter, burning, disagreeable taste. It mixes uniformly with absolute alcohol and volatile and fatty oils, and is readily dissolved by ether. It contains a volatile oil and resin.

In 1829, Schweitzer (*Poggend. Annal.*, Bd. xvii. pp. 487 and 1095) announced that he had found in copaiba a peculiar crystallizable acid, *Copaivic Acid*. It is not in the plan of the present work to discuss elaborately the chemistry of drugs, and consequently I must remain content with the statement that the researches of Prof. Bernatzik (*Prager Vierteljahrschrift*, Bd. c., 1868, p. 239) have shown that very frequently this crystalline acid does not exist in copaiba, and consequently that it is an unimportant constituent. Indeed, these researches seem to me to prove conclusively that copaiba is simply an oleoresin.

PHYSIOLOGICAL ACTION.—When given in therapeutic doses, copaiba has very little if any action upon the general system, and the influence even of very large amounts is often scarcely perceptible. In the researches of Bernatzik (*loc. cit.*, p. 251), eighteen grammes of the volatile oil were taken in three doses during twelve hours, and caused only an acceleration of a few beats per minute in the pulse-rate, and a rise of a fraction of a degree in the temperature, with, after a time, violent gastric and intestinal disturbance, evidently due to the local action of the drug, and characterized by vomiting and purging. Complete strangury was not produced, but there was some difficulty in passing the urine, which caused decided burning in the urethra.

On the other hand, the action of the drug is much more decided upon some very susceptible persons, so that full doses of it produce decided fever, with increased frequency of pulse, and hot skin, accompanied almost always by

decided symptoms of gastro-intestinal irritation. Sometimes, also, the urinary organs are more sensitive than usual to the action of the drug, so that strangury, and, as is stated by some authorities, even almost complete urinary suppression, may occur.

In 1841, Dr. G. O. Rey (quoted by Bernatzik) called attention to the fact that if nitric acid be added to the urine of persons taking copaiba, a precipitate is formed resembling that of albumen. This fact has been noted and commented on since its discovery by Dr. H. Weikart (*Archiv der Heilkunde*, 1860), by Dr. Rees (*Guy's Hospital Reports*, vol. xvii.), by Valentine (*Grundriss der Physiologie*), and by other observers. In order to produce the phenomenon with distinctness and certainty, it is seemingly necessary to use large doses of the drug, since Weikart failed to detect it after the exhibition of small amounts of copaiba oil. Various surmises as to the nature of this precipitate have been indulged in; but the experiments of Bernatzik (*loc. cit.*, p. 252) appear to show that it consists of the oxidized oil united to some urinary principles. Dr. Bernatzik found that the elimination of the oil goes on slowly, continuing for as much as four days after its ingestion when large doses are employed.

In his experiments with the resin of copaiba, the authority just noted exhibited fifteen grammes of it inside of five hours. It acted as an emeto-cathartic, causing a great deal of pain and irritation. The urine deposited very copiously on the addition of nitric acid, the resin seemingly being eliminated more abundantly than was the oil.

Copaiba does not increase to any great extent the amount of the renal secretion, and no evidence has, that I am aware of, been offered to show that it affects materially the solids of the urine.

The clinical employment of copaiba has shown that the drug exerts its peculiar stimulant alterative action on other mucous membranes than those of the genito-urinary apparatus, and it is very possible that a slight elimination of the volatile oil takes place through the lungs.

THERAPEUTICS.—The chief use of copaiba in medicine is in *subacute* and *chronic inflammations* of the *genito-urinary* mucous membrane. In its action upon this structure it is a decided stimulant, but is less irritating and less stimulating than the oil of turpentine.

Gonorrhoea is the disease in which it is mainly exhibited. It is especially useful in the advanced stages of this affection. If it be given in the beginning, before the inflammation has fully developed, it may sometimes succeed in aborting the attack, but, if it fail to accomplish this, may greatly aggravate the symptoms. During the height of the inflammatory stage, copaiba should not be employed.

In other inflammatory affections of the genito-urinary mucous membrane, such as *pyelitis*, and chronic *cystitis*, when the disease is of a subacute or chronic character, the remedy may be employed, it being borne in mind that in its action it is much more stimulant than buchu or pareira brava, but much less so than turpentine.

In chronic *diarrhœa* and *dysentery*, copaiba is sometimes of use, through its local action on the diseased surfaces.

The remedy has been highly recommended in the advanced stages of *bronchitis*; and in the chronic form of the disorder when attended by very free muco-purulent expectoration, I have occasionally employed it, with excellent results.

Copaiba is sometimes used internally in *skin-affections*, but more frequently is employed externally as a stimulant dressing.

ADMINISTRATION.—The dose of copaiba is ten to twenty minims, repeated from three to six times a day, according to circumstances; the best effects are probably to be attained by the frequent use of small quantities. The medicine may be given dropped upon sugar, or, what is much better, exhibited in an aromatic emulsion, made with syrup and mucilage of gum-arabic in such a manner that a tablespoonful shall contain the required dose. When patients object to the taste, the drug may be given in gelatine capsules, each containing ten drops. It is said that these capsules do not, however, agree so well with delicate stomachs as the emulsion. When copaiba is rubbed up with magnesia, the resin unites with the earth to form a solid mass, in which the oil is mechanically held. The officinal pills (*Pilulæ Copaibæ*) are made in this way; but, as they are disintegrated and absorbed with difficulty, the pills of copaiba are not an eligible preparation. Each pill contains four and a half grains of the copaiba.

The oil which is prepared by distillation is isomeric with oil of turpentine. It is officinal under the name of *Oleum Copaibæ*, and may be exhibited in emulsion or in capsules, in doses of eight to fifteen minims.

CUBEBA—CUBEBS.

The unripe berries of the Piper Cubeba, a climbing plant of Java and other portions of the East Indies. These berries are blackish-veined, about the size of a small pea, and have attached to them a short stalk three or four lines long. Their odor is aromatic and peculiar; their taste warm, camphoraceous, and peculiar. Cubebs is a somewhat complex body, but there can be no doubt that the ethereal extract, or oleoresin, as it is commonly called, represents its medicinal virtues. The extract appears to consist chiefly of three substances: a volatile oil and the brown resinous substance formed by its oxidation, a peculiar acid, and a neutral crystallizable principle, *Cubebin*. *Cubebic Acid* was first discovered by Monheim, but has been especially examined by Prof. Bernatzik (*Prager Vierteljahrschrift*, Bd. lxxxi., 1864, p. 9). It is nearly tasteless, forms salts with the bases, has a very faintly acid reaction, and dissolves in concentrated sulphuric acid with the production of (according to Bernatzik) a purple-violet color, changing on the addition of a little water to a cherry-red, and altogether disappearing when further dilution is practiced.

PHYSIOLOGICAL ACTION.—In some respects, cubebs, when taken in-

ternally, resembles black pepper in its effects. It is, however, much less stimulating than its congener. After the ingestion of the ordinary therapeutic dose of cubebæ, nothing unusual is experienced; but when very large amounts are taken, there are evidences of gastric excitement, such as sensation of warmth in the stomach, slightly-increased frequency of pulse and heat of skin, with perhaps some giddiness or headache. The urine is slightly increased in amount, and acquires a peculiar odor. When very large doses are taken, the symptoms of gastric irritation are more severe, and the subject suffers from gastric burning, nausea, vomiting, and colicky pains, with, in some cases, purging.

An eruption resembling urticaria has been occasionally noticed after the ingestion of cubebæ: it is exactly similar to the eruption sometimes caused by copaiba, and, like it, is probably due simply to gastric irritation.

That the active principles of cubebæ are eliminated by the urine is well established, as, after the exhibition of the drug, when nitric acid is added to the urine, a precipitate, resembling somewhat that of albumen, occurs.

The most elaborate physiological study of cubebæ that I have met with was made by Prof. Bernatzik (*loc. cit.*). This experimenter took himself, and gave to a student, ten grammes of the cubebate of magnesia. No decided symptoms were induced by this, further than some pulse-acceleration and gastric uneasiness; but the elimination of uric acid was greatly increased, and the cubebic acid was found in the urine. Half an ounce of the oil of cubebæ was taken in thirty-six hours, the last three doses, aggregating ten grammes, being ingested in six hours. This was followed by very decided gastric irritation; by the appearance in the urine of the oil, not as it was ingested, but oxidized and in the form of a resin; the eliminated uric acid was about one-third in amount of that excreted after the exhibition of cubebic acid. Of the powdered cubebæ, fifty grammes were taken in eight hours; the gastro-intestinal irritation was very marked; the nitric-acid precipitate was abundant in the urine; the elimination of uric acid was about midway between the extremes of the previous experiments. According to the researches of Prof. Bernatzik, *cubebin* is inert.

THERAPEUTICS.—Cubebæ has been used to some extent for its local stimulant action upon the alimentary canal, but for this purpose is very inferior to black pepper and other spices. It is at present almost exclusively employed to influence the genito-urinary mucous membrane, in precisely those cases in which copaiba is exhibited. The two drugs have very nearly the same range of action, but the cubebæ is less apt than is the copaiba to derange digestion. Very often the best effect in *gonorrhœa* and other genito-urinary disorders is obtained by giving the two remedies in combination.

Cubebæ is sometimes employed with asserted advantage in chronic *hæmorrhoids*, and also in those forms of *bronchitis* in which copaiba is useful.

It forms the basis of certain proprietary lozenges, much used by public speakers and others to relieve the relaxation of the larynx which follows slight

colds or over-use. For this purpose the berries themselves may be chewed, and are very effectual.

In *coryza*, the powdered drug used as a snuff has sometimes a beneficial effect. It should not be employed in the early stages before secretion has been established, but is indicated later in the affection, when the discharge is profuse.

ADMINISTRATION.—The dose of powdered cubeb is from half a drachm to three drachms, which may be exhibited in syrup or molasses three times a day. The volatile oil of cubeb (*Oleum Cubebæ*) is officinal, and may be given in emulsion three or four times a day, in the dose of fifteen drops, gradually increased to half a drachm, unless some effect is previously produced upon the urinary organs. This oil does not as thoroughly represent the crude drug as the oleoresin (*Oleoresina Cubebæ*), which is also officinal, and may be given in doses of from ten to fifteen minims, increased as necessary. It is best administered in emulsion, but may be exhibited in the form of a bolus, enough sugar having been added to make a plastic mass. The tincture (*Tinctura Cubebæ*) is officinal, but is an ineligible preparation.

MATICA.—*Matico*, an officinal drug, the dried tops of the *Piper angustifolium* of Peru, contains a volatile oil, resin, and, it is said, a bitter principle, *Maticin*. It is largely employed as a styptic, and as such probably acts chiefly mechanically, coagulating the blood in its interstices, adhering to the wound, and thus arresting the hemorrhage. It has also been employed in internal hemorrhages, and in gonorrhœa. In these affections it probably acts similarly to oil of turpentine, although much less of a stimulant, and much more feeble.

CANTHARIS.—*Cantharides* is considered elsewhere in detail (see *Epispastics*), and it is only necessary here to say a few words in regard to its use in diseases of the genito-urinary tract. The active principle of Spanish flies is certainly eliminated by the kidneys, and acts therefore locally upon these organs, as well as upon those over which their secretion flows. The influence exerted by this means is simply one of intense irritation, cantharides being an irritant to these organs in any dose sufficiently large to have an effect. Indeed, of all the officinal drugs, cantharides is the most actively irritant to the kidneys and subordinate organs. Consequently it is employed only when an intensely stimulant action is desired, as in obstinate *gleet*, in which affection it is often combined very advantageously with the tincture of the chloride of iron. In *pyelitis* and *cystitis* it is very rarely indicated, but may be cautiously employed in very chronic cases.

The tincture of cantharides is the only preparation used internally. For the dose and method of administration, see *Epispastics*.

CLASS IV.—DIAPHORETICS.

DIAPHORETICS are those medicines which are employed to increase the action of the skin. It is scarcely in place here to discuss the results of suppression of the functional activity of the skin or the importance of the surface-elimination to the system. It does seem well, however, to call attention to the fact that the perspiratory glands have a double function to perform,—that of elimination, already alluded to, and that of keeping down the temperature of the body during exposure to heat. When a man enters a Turkish bath whose temperature is perhaps 160° F., or when he works in the sun of a very hot day, there is, if he be used to such exposure, little or no rise in the temperature of the body, because the surface-glands secrete sweat so actively as to expose a great amount for evaporation, and by the conversion of so much water into vapor such an amount of heat is absorbed—*i.e.*, converted from heat into repulsive force—as to cool the body. The reason that even a moderate degree of heat in a moist atmosphere is intolerable is because evaporation cannot take place.

From what has already been stated, it is obvious that the use of dry external heat, or rather exposure to a hot atmosphere, is a powerful means of producing perspiration: it is, indeed, in healthy men the most powerful method at our command. It may be applied either in the form of the *Turkish bath*,* in which the air of the hot chamber is very dry, or in the *Russian or vapor bath*, in which the atmosphere is surcharged with hot vapor. Very wonderful therapeutic properties have been ascribed to the direct action of heat (Ureghart, *Manual of the Turkish Bath*, London, 1865)† when applied by the Turkish bath; but the remedy appears to me to act only as a powerful sudorific, perhaps also doing good in some cases of acute internal congestion by attracting the blood to the surface and thereby depleting the interior.

In private practice, or whenever a properly-provided bath cannot be com-

* The term Turkish Bath is here applied to the bath used in this country under that name. This bath appears not to be a copy of the Oriental bath, but merely a derivative from it. According to a writer in the *British and Foreign Medico-Chirurgical Review* (vol. xxxvii. p. 87), in the East, the sudarium, or sweating-chamber, rarely has a temperature of more than 98° F.

† Consult also *Bathing: How to Do It, When to Do It, and Where to Do It*, by E. Sheppard, London, 1865; *The Anglo-Turkish Bath*, by Y. J. Moore, London, 1865.

manded, a very efficient and readily-applied substitute consists of a large tin funnel furnished with a long bent beak, a stool with a hole in the centre of its seat, or else a few bricks, and a large spirit-lamp. The patient being closely wrapped up in bed, and the clothes being especially "tucked in" about the neck and shoulders, the funnel is placed upon the stool or the bricks in such a manner that the beak of it enters well under the bedclothes, coming close to but not in contact with the person of the sick man. The spirit-lamp, being then placed immediately under and close to the funnel, must, when lighted, send a column of hot air and vaporized water through the beak into the space around the body of the patient. When the lamp is sufficiently large, and a little care is taken to see that the nozzle of the funnel is not obstructed by the bedclothes, the process just detailed affords a very efficient method of giving a vapor-bath.

Hot-water baths offer another very successful method of inducing profuse perspiration. The patient should be placed in a bath of about 98° or 100° F., and remain there fifteen or twenty minutes, during which time, by the repeated addition of very hot water, the temperature should be raised to 110° F. or to such point as the patient can endure. Warmed blankets having been plentifully provided, the sick man should be lifted from the bath into them, be closely wrapped up, and so left for three or four hours before being transferred to the usual bed.

According to Dr. A. Steffen (*Jahrbuch für Kinderheilkunde*, Heft iii., 1871), after this use of the bath the body has been proven to undergo loss of weight continuously for one or two days.

Profuse sweating is always more or less exhausting, but is not nearly so much so as purging, and therefore may be practiced in dropsical patients too feeble to allow of the use of purgatives. The hot baths are not, however, altogether free from danger or objection. Sometimes in the Turkish and Russian baths the patient fails to sweat freely, and a feeling of distress, a bounding, rapid pulse, and perhaps severe headache, develop themselves: under these circumstances the bodily temperature rises, and a fever develops, which may go on to the production of a true "thermic fever," and perhaps terminate in sudden death. This is an exceedingly rare result, and one that never can occur if a little watchfulness be practiced and the patient removed from the hot chamber so soon as any unpleasant symptoms are manifested. Sudden death has, I believe, taken place once from "sun-stroke" in a patient whilst taking the "Turkish bath."

The use of hot baths of any kind is, of course, contra-indicated by the existence of fever; but, according to Dr. Steffen, the hot-water baths are pre-eminently contra-indicated by the existence of congestion or œdema of the lungs, or of a tendency towards these disorders, since under such circumstances the bath greatly increases the disease, or precipitates a perhaps fatal attack. My own limited experience, so far as it has gone, has corroborated these statements of Steffen. I have seen, under the conditions mentioned,

the most frightful dyspnoea result from the use of the hot-water bath, a dyspnoea which was apparently prevented from terminating fatally only by the removal of the patient from the bath-tub. If disturbance of the respiration come on during the bath, the patient should be immediately taken out, and, if the symptoms be urgent, cold water should be freely dashed over the head, neck, and chest.

Diaphoretics produce the desired result in various methods, which may be briefly considered under four headings, representing as many modes of action.

First. *By relaxing the skin.* As has already been sufficiently shown (see purgatives, p. 371), there is a form of secretion, or perhaps it would be more correct to term it of leakage, from mucous membranes, which is distinctly paralytic in its mechanism. The same assertion may be made in regard to the skin: the colliquative, so-called "night-sweats" of phthisis afford a familiar example of this, occurring as they do in profoundly debilitated subjects, and at such times as there is the greatest relaxation of the system,—i.e., during sleep. The profuse sweats of collapse also may be instanced as examples of the general truth just enunciated. Normal processes which produce great relaxation cause great sweating: thus, during vomiting, especially if it be accompanied by much nausea, the skin pours out its secretion. By virtue of this general law certain remedies act as diaphoretics. All of the diaphoretics which cause sweating by producing relaxation, and which are employed in medicine, are nauseants, constituting a distinct group,—the *Nauseating Diaphoretics*.

Second. *By reducing the force of the circulation.* There is undoubtedly a condition of over-action or over-rapidity of the circulation in which the affected glands are unable to perform readily their normal functions. Thus, it is well known that the first stage of inflammation is one of arrested secretion, and that in high fever there is a general drying-up of the secretions. The skin does not differ from other organs in this respect: consequently its functional activity may fail because of excessive arterial action. Hence there is a class of remedies which, although perhaps not actively sudorific in health, are in disease very efficient in reducing the circulation and restoring the functional activity of the skin. It is evident that there is a close connection between the present mode of influence and that noted in the previous section; and it is no less apparent that the nauseant diaphoretics act most powerfully in reducing the circulation. There are, however, certain diaphoretics which act in the present method but are not nauseants: these sudorifics form a separate class by themselves,—the *Refrigerant Diaphoretics*.

Third. *By entering the circulation and directly stimulating the glands of the skin.* It appears to be a general law that when any medicinal principle is eliminated by any excretory organ, the general activity of that organ is increased by the effort at elimination. Thus, the purging of arsenical poisoning, the vomiting of antimonial poisoning, the increased urinary secretion following the ingestion of a potassium salt, are apparently all of them the

results of attempted elimination. The skin undoubtedly eliminates medicinal substances, and is undoubtedly subject to the general law : consequently there is a class of remedies which increase its action by a direct influence.

It is manifest that a drug may relax the general system, may diminish the force of the circulation, and also may stimulate directly the skin : indeed, it is most probable that antimony does all of these ; and at least some of the refrigerant diaphoretics probably act in the last two ways. There are, however, certain substances which seem to cause sweating simply by stimulating the function of the skin. These are in this work grouped as *Simple Diaphoretics*.

Fourth. *By filling up the blood-vessels.* There is much reason for believing that precisely as under certain circumstances water, by increasing the amount of the blood, will provoke increased renal secretion, so, under other circumstances, will it cause increased dermal excretion.

The antagonism which exists between the skin and the kidneys in regard to the amount of their respective secretions has already been sufficiently dwelt upon (see p. 404). It seems well to reiterate, however, that warmth favors the action of the skin, whilst cold stimulates the renal activity. Thus, large draughts of water, if taken cold, the patient being kept cool, increase the urine, but, if they be taken hot, and the patient covered up warmly in bed, increase the perspiration.

Diaphoretics are employed in the practice of medicine to fulfil the following indications :

First. *To arrest forming diseases* of not very severe type, probably by causing a flow of blood to the surface, and thereby relieving slight internal congestions, and possibly by eliminating principles which have been retained in the blood instead of being excreted as they ought to have been. In *general cold*, in *muscular rheumatism*, *suppressed menstruation*, and other results of exposure to cold and of checked perspiration, the diaphoretics afford the most efficient means at our command for restoring the normal functions.

Second. *To favor absorption.* In *dropsy* the diaphoretics are of very great value, often aiding diuretics and purgatives in effecting a cure, and sometimes, when these fail, or when circumstances forbid their use, rescuing the patient from impending death.

None of the medicinal diaphoretics are of sufficient power to be relied upon in dropsy : in order to reduce an effusion, the Turkish, the Russian, or the hot-water bath must be vigorously employed, the medicinal diaphoretics being used merely to aid in their operation.

Third. *To aid in the subsidence of diseases* which naturally pass off with a sweat. The chief use of diaphoretics for this purpose is in *miasmatic fevers*, especially in the *remittent* form of the affection, when the sweating stage fails to develop itself thoroughly and the paroxysms run into one another. Even in the single paroxysm of *intermittent fever*, by hastening the closing stage, diaphoretics will often shorten the paroxysm.

Fourth. *To eliminate noxious materials* from the blood. The old humoral idea of the ground-work of such diseases as fevers, the belief in a distinct materies morbi which could be eliminated from the blood, has no sufficient demonstration to be accepted, and, although diaphoretics do good in *fevers*, yet it cannot be granted that it is in this manner. The very great power of increased diaphoresis in cooling the body through surface-evaporation has already been dwelt upon; and much of the good effected by diaphoretics in diseases of high temperature probably has its origin in this power. Modern science seems, however, clearly to point out that this class of remedies may aid in separating from the blood retained secretions, and may to some extent replace the action of the kidneys when these organs are disabled by disease.

In 1851, Dr. Schottin (*Archiv für Physiologische Heilkunde*, Jahrg. xi.) discovered urea in the sweat of patients suffering from the collapse of cholera. Not only has the discovery of Schottin been confirmed by the researches of G. O. Rees (*Encyclopædia of Anatomy*, vol. iv. p. 841), of Fiedler (*Dissert. Inaug.*, Leipsic, 1854), of Hirschsprung (*Gaz. des Hôpitaux*, 1865), of Kaup and Jürgensen (*Deutsches Archiv für Klin. Medicin*, Bd. vi., 1869, p. 54), of Leube (*Ibid.*, Bd. vii., 1869, p. 3), and of G. Deininger (*Ibid.*, p. 587), but it has also been abundantly proven that the skin excretes urea freely during the advanced stages of Bright's disease, and also during the partial suppression of urine of scarlatinal desquamation and nephritis. The urea in renal disease may even form a distinct crystalline powder on the skin; but it is most abundant about the mouths of the sweat-glands. I believe Landerer was the first to announce that urea was present in the sweat of healthy persons; and, although chemists of excellence have been unable to detect it, yet its presence at times can no longer be denied, since it has been found not only by Landerer, but also by Funke in 1858 (*Moleschott's Untersuchungen*, Bd. vi.), by Meissner (*De Sudoris Secretione*, Dissert. Inaug., Leipsic, 1859), and by Leube (*loc. cit.*); Fourcroy (quoted by Rees, *Encyclopædia of Anatomy*, vol. iv. p. 84) has also found it in the sweat of horses. By a series of elaborate experiments, Leube (*loc. cit.*) has rendered it probable, if he has not actually proven, that in health there is such a relation between the skin and the kidneys that when the former is very active the latter excrete less than the normal amount of urea.

When to the facts already narrated are added the observation of Griesinger, that in diabetes the perspiration contains sugar, and the well-known circumstances that in rheumatism the sweat contains lactic acid, and in jaundice biliary products, the value of diaphoretics as a means of getting rid of retained excretions becomes manifest. For this reason, in Bright's disease, especially of the acute form, they are of the greatest value, acting beneficially in three different ways: by drawing the blood to the surface, and thereby relieving any internal congestions of the kidneys or other organs that may exist; by promoting the absorption of dropsical effusions; and by eliminating retained secretions.

NAUSEATING DIAPHORETICS.

The most frequently employed medicine of this class is *tartar emetic*. This substance is so fully discussed in the earlier portion of this book that very little need be said about it here. It seems well to point out, however, that the antimonials act as diuretics, even when not given in sufficient doses to cause nausea, and that they do so apparently in two ways: by reducing the force of the arterial circulation, and by a direct action. They are probably eliminated to a very slight extent by the skin; although this has not, to my knowledge, been proven. Tartar emetic is to be employed as a diaphoretic in sthenic cases only, and is especially used in *inflammatory fevers*. The dose of it as a diaphoretic is from the sixth to the twelfth of a grain. If a diaphoresis is urgently demanded, the dose should be gradually increased until decided nausea is induced. An excellent combination in many cases is tartar emetic, neutral mixture; and a minute amount of morphia.

Ipecacuanha is another nauseating substance, which, like tartar emetic, seems to exert an influence upon the skin, even independently of its action on the stomach. Moreover, like the antimonials, it never in small doses by itself causes profuse sweating, but simply seems to aid in maintaining the insensible perspiration and in keeping the skin soft and pliable in inflammatory fevers. Partly for this reason it is frequently combined with mercury in such diseases as acute *peritonitis*. The diaphoretic dose of ipecacuanha is a grain every two hours. Some persons with delicate stomachs are decidedly nauseated by this amount; and to these only half a grain should be given at a time.

A very famous and efficient diaphoretic preparation of ipecacuanha is *Dover's Powder* (*PULVIS IPECACUANHÆ COMPOSITUS*), which contains one grain of opium, one grain of ipecacuanha, and eight grains of the sulphate of potassium. Dover's powders are employed in two distinct methods. In some diseases, as in acute *rheumatism*, an opiate is indicated to allay pain or for some other purpose, and at the same time a diaphoretic is needed to keep up the action of the skin. Under these circumstances, from three to five grains of the compound powder of ipecacuanha may be given every two, three, or four hours, *pro re nata*. The medicine is more apt to nauseate when taken in powder than when exhibited in pills; for which reason the latter form of administration is preferable in the class of cases now under consideration, especially as the powder is to most persons a disagreeable medicine.

In the second method of using Dover's powder, a single large dose is given for the purpose of producing profuse sweating. In the intense suffering which sometimes results from *suddenly suppressed menstruation*, the remedy is most efficient, alleviating the pain and aiding in the production of the desired diaphoresis. In breaking up a *cold*, or in *muscular rheumatism*, Dover's powder is the basis of the most commonly practiced method of causing sweating (see Alcohol as a Diaphoretic).

REFRIGERANT DIAPHORETICS.

Aconite, veratrum viride, and all the various remedies used to depress the cardiac action when excited, are, in the strictest sense of the term, refrigerant diaphoretics. Sufficient has, however, already been said in regard to their use.

The *citrate of potash*, whether in the form of *effervescing draught* or of *neutral mixture*, is constantly employed in sthenic fevers, and affords, I think, the best basis there is for fever-mixtures in such cases, the more powerful depressants being added to it as circumstances demand.

SIMPLE DIAPHORETICS.

LIQUOR AMMONII ACETATIS—SOLUTION OF THE ACETATE OF AMMONIUM.

Spirit of Mindererus is prepared by saturating dilute acetic acid with the carbonate of ammonium, and consequently consists of a solution of the acetate of ammonium, containing as much of free carbonic acid as the water will absorb. It is a colorless liquid, and should have no odor, or a very faint odor of acetic acid. Any specimen having an ammoniacal odor should be rejected, as containing an objectionable excess of the carbonate of ammonium. The taste is disagreeable and saline. As the solution upon standing undergoes decomposition after a time, it should be freshly prepared when needed for use.

THERAPEUTICS.—*Spirit of Mindererus* appears to be a feeble stimulant diaphoretic. Cullen has known eight ounces of it to be taken in a very short time without inducing any effect; yet the testimony is very strong as to its having some value in disease, and in *adynamic fevers* it may be employed as a diaphoretic. It is, however, much less efficient than the sweet spirit of nitre, and is probably to most persons more disagreeable than that favorite drug. Special value has been claimed for it in *dysmenorrhœa*, and even in *menorrhagia*; but I do not believe that it has any superiority over other diaphoretics in the former affection, and its being of any use in the latter disease seems to me apocryphal. The dose of it is one to two tablespoonfuls.

SPIRITUS ÆTHERIS NITROSI—SWEET SPIRIT OF NITRE.

This is at present officinally prepared by adding sulphuric acid to alcohol, then adding copper wire, and after this nitric acid, distilling at a temperature not exceeding 180° F., and adding alcohol to the distillate. The first reaction which takes place between the sulphuric acid and the alcohol results in the liberation of ether, the hydrated oxide of ethyl; the nitric acid, giving to the copper two of its equivalents of oxygen, is converted into nitrous acid, which combines with the oxide of ethyl to form a nitrite.

Spirit of nitrous ether is a volatile, inflammable liquid, of a pale-yellow color inclining slightly to green, having a fragrant, ethereal odor, free from pungency, and a sharp, burning taste. It slightly reddens litmus, but does not cause effervescence when a crystal of bicarbonate of potassium is dropped into it. When mixed with half its volume of officinal solution of potassa previously diluted with an equal measure of distilled water, it assumes a yellow color, which slightly deepens, without becoming brown, in twelve hours. A portion of the spirit in a test-tube half filled with it, plunged into water heated to 145° and held there until it has acquired that temperature, will boil distinctly on the addition of a few small pieces of glass.

Spirit of nitrous ether has the specific gravity 0.837, and contains five per cent. of its peculiar ether. It should not be long kept, as it becomes strongly acid by age.

PHYSIOLOGICAL ACTION.—Undoubtedly, the sweet spirit of nitre, if taken in sufficient amount, acts very decidedly upon the organism. Dr. Christison narrates the case of a woman who was found dead in her bed, and whose death was attributed to the inhalation of the fumes from a large quantity of the drug which had been spilled in her apartment. Mr. D. R. Brown reports in the *Pharmaceutical Journal and Transactions* for March, 1857, the effects of the inhalation of sweet spirit of nitre. The first symptoms are a bluish-purple, livid discoloration of the lips and fingers, and a peculiar pallor of the face. These increase, the face assumes a ghastly look, the extremities grow cold, and the pulse becomes very weak and frequent, but the breathing remains slow and regular; muscular weakness is extreme, and the least exertion causes hurried respiration, with painful oppression in the chest, and cardiac distress. Confusion of mind and giddiness may or may not occur, but headache always comes on sooner or later.

When given by the mouth in the ordinary therapeutic doses, sweet spirit of nitre exerts no marked effect, except in increasing the action of the kidneys and of the skin. It is generally believed, and probably correctly so, to exert a stimulant action upon the nervous system.

Clinical evidence shows at least that the drug often has the same action upon disturbed innervation as the so-called antispasmodics. This is especially visible in the case of children.

THERAPEUTICS.—Sweet spirit of nitre is one of the most popular of the diaphoretics. As it exerts a stimulant action, it is useful in *adynamic* rather than in *sthenic fevers*. In children with fever offering nervous symptoms, such as starting, jerkings, etc., it is especially useful.

If the patient be kept about and cool, instead of being warmly covered in bed, spirit of nitrous ether acts as a decided diuretic; but it is not sufficiently powerful to make it worthy of reliance in *dropsy*.

The dose of sweet spirit of nitre is: for an adult, a teaspoonful to a tablespoonful; for a child three years old, half a teaspoonful. If a diaphoretic action is required, very minute doses should be exhibited at short intervals.

Thus, for a child a year old, a teaspoonful should be put in five ounces of water, and a tablespoonful of this given every hour.

ALCOHOL.—As is known to every hard drinker, alcohol when taken in excess, and especially if drunk with hot water, is eliminated by the skin, and often causes profuse sweating. As a diaphoretic it is, however, used in medicine for only one purpose,—*i.e.*, in those cases, such as a forming “cold,” *subacute rheumatism*, and *suppressed menstruation*, when a single profuse sweating is desired. In these cases I have found the following plan most efficacious, provided the patient have not decided fever: a Dover’s powder is administered, and directly after this the patient goes into a hot-water or vapor bath, or else does what is known in common parlance as “soaking his feet,”—*i.e.*, takes a pediluvium. The proper method of doing the former of these acts has been sufficiently described. In taking a foot-bath the patient should use a tub of sufficient size to enable him to place in it his feet and legs up to the knees; the water should be as hot as can be borne, and, as the feet become a little accustomed to the temperature, hotter water should be added; the bath should continue fifteen or twenty minutes; during it, the patient, being in his night-clothes, should be well wrapped up in blankets, and at the close should be so transferred to the bed as not to get in any way chilled. After he has got to bed, and has been heavily covered with blankets, the patient should drink one or two tumblerfuls of a very hot and strong lemonade, containing one, or two, or three tablespoonfuls of whisky or brandy.

I think the popular belief that after a sweat there is a greater liability than usual to take cold is well founded: care must, therefore, be exercised to avoid any exposure for a day or two.

CLASS V.—EXPECTORANTS.

THESE are medicines possessed of the power of modifying the secretions and thereby influencing the inflammatory conditions of the respiratory mucous membranes. There are various, and even opposing, methods, in accordance with which drugs act upon the pulmonary surfaces. In certain states of the latter, as in the first stages of acute bronchitis, the irritation of the part is too great for secretion, the first stage of inflammation being, as is well known, connected with suspension of function: in this condition the so-called *sedative expectorants* are indicated. These are remedies which lower arterial action: they are all nauseants, and the increase of bronchial secretion by them is analogous to that which they produce in the skin. In order to get their full effect, they must be given in nauseating doses; and if these are gradually increased until emesis is induced, the fullest therapeutic influence will be obtained.

There are other expectorants which act directly upon the bronchial mucous membrane, some of them perhaps increasing secretion, but most of them rather modifying it, and some of them even diminishing it, by toning up a relaxed, over-secreting mucous membrane. Some substances which can hardly be called expectorants are nevertheless useful in bronchial diseases: thus, gallic acid will sometimes lessen the large quantities of mucus secreted in *bronchorrhœa*; or, when the mucus is tenacious in *chronic bronchitis*, alkalies will frequently be of great service by rendering it less viscid and thereby facilitating expectoration.

In many cases of disease occurring in the very young and in the very old, mucus may so accumulate in the lungs, owing to the inability of the enfeebled powers to force it up, as seriously to embarrass, or even fatally compromise, respiration. Very frequently, in acute cases of this character, such as the *suffocative catarrh* of infants, mechanical emetics are of the greatest service; but in chronic cases they induce so much disturbance of digestion as to render their use dangerous, and alcohol, carbonate of ammonium, oil of garlic or of turpentine, and other drugs capable of increasing the bronchial muscular power or activity, have to be relied upon.

Again, in certain conditions of the lungs, especially in *chronic catarrhal pneumonia*, iodine and other alteratives are of great value in facilitating the

absorption of exuded materials; yet these remedies can hardly be called expectorants.

It may be affirmed that the value of true expectorants in *pneumonia* of any variety is exceedingly problematic, and, except it be the muriate of ammonia, they far more often do harm in the chronic varieties by deranging digestion than they do good by influencing the lung-tissue.

There are several substances, chief among them being morphia, hyoscyanus, and hydrocyanic acid, which have the power of allaying cough and lessening irritation by an anodyne action. In regard to morphia, its property of checking secretion should not be forgotten, but should not prevent its use. These narcotics are especially useful when the cough is disproportionate to the amount of inflammation. When large quantities of mucus are being secreted in debilitated subjects, their use requires great caution, for fear of benumbing the nerves or nerve-centres and thereby increasing the danger of an accumulation of phlegm in the lungs.

Expectorants may be arranged under two heads: the nauseant or sedative expectorants, and the stimulating expectorants; and the general proposition may be laid down that expectorants of the first class are to be used in the first stages of acute bronchitis, whilst expectorants of the second class are to be employed in the advanced stages or in the chronic varieties of the disease. As these diseased conditions gradually merge into one another, so must the practitioner balance the ingredients of his expectorant mixtures, adapting their relations to the individual case.

The present seems a fitting place to say what is necessary in regard to the use of drugs by *atomization*. In this method of administration, a solution of the medicine employed is broken up by a mechanical contrivance into a fine spray and projected into the back of the mouth. There can be no doubt that when the operation is properly performed the spray reaches even the finest ramifications of the pulmonary bronchi. A discussion of this is at present scarcely necessary. Any reader having doubt upon the subject will find the evidence in the work of Dr. J. Solis Cohen (*Inhalation: its Therapeutics and Practice*, Phila., 1867).

The following rules should be attended to, to secure successful results:

1. Use the steam atomizers: all other forms of apparatus give irregular or too feeble currents, and should be discarded.

2. See that the points of the atomizing tubes are sharp and clean,—not foul, cracked, or with their edges worn.

3. See that the steam is generated equably and with sufficient force, and that the solution used is free from all solid particles, and, unless otherwise ordered, about the temperature of the body.

4. Never allow inhalations when the patient is excited, directly after eating, or immediately after exercise, unless especial circumstances, as in hæmoptysis, demand haste.

5. Have the glass mouth-speculum inserted well into the mouth, and the

line of its axis and of the propulsion of the spray coincident with that of the mouth.

6. When the pharynx, or even the larynx only, is to be reached, the operation is very simple, and respiration should be natural; but when it is desired to make applications to the ultimate bronchi, the respirations should be regular, slow, and as deep and full as possible, the lungs being well emptied at expiration.

7. Do not protract the sitting until the patient is fatigued. Five minutes is generally long enough to commence with. In hæmoptysis, this rule may sometimes be departed from with advantage.

8. Let the patient give his whole attention to the matter in hand.

9. In chronic disease, one, two, or three inhalations a day are usually sufficient. In acute disease, they may be required much more often, as every hour in diphtheria.

10. Never use atomization for the purpose of general medication: it is simply a method of applying substances locally to the respiratory organs.

The substances used by atomization may be conveniently arranged as follows:

DILUENTS.—The only diluent of any value is *warm water*. In acutely *inflamed conditions* of the *mucous membrane*, the very frequent application of water at about the temperature of 90° F. will often afford marked relief, especially in *laryngitis*.

ASTRINGENTS.—These are employed to arrest excessive secretion or hemorrhage. In their use, it must always be remembered that they are more or less irritant, and that whilst some persons bear them very well, in others they produce very harmful irritation; also, that the idiosyncrasies of patients vary, so that whilst one person will best bear a certain one of the astringents, a second may be less irritated by another drug. The rule is to try carefully until the pulmonary idiosyncrasies are known. In cases of excessive *bronchial secretion*, any irritation sufficiently severe to give origin to tightness in the chest, or to much coughing, calls for the withdrawal of the medicine. In *hæmoptysis*, a greater risk can be judiciously assumed. The astringent substances employed in this way are: First, *tannic acid*, one to twenty grains to the fluidounce of water. I have not used this; but it is said to be well adapted to cases of free secretion. Second, *alum*, varying in strength from a solution of five grains to the ounce to a saturated solution. This I have used with great satisfaction. In *hæmoptysis*, only the strongest preparation is of avail; in *bronchorrhœa*, a weak solution should be employed at first, and the strength increased *pro re nata*. One advantage this drug has over tannic acid is its compatibility with the sulphate of morphia, which should always be added to its solution when any tendency to irritation exists. In *chronic bronchitis*, with excessive expectoration, I have seen alum produce most gratifying results. Third, preparations of *iron*: of these the solution of the

perchloride has been used in Great Britain and on the Continent; but the Monsell's solution (*Liquor Ferri Subsulphatis*) is much preferable, as even more powerfully styptic and less irritating. Ten drops of the solution may be added to the fluidounce of water and be used for the trial dose. If the lungs will bear it and necessity exists for a stronger solution, the strength may be carefully increased up to half a fluidrachm to the ounce. *Acetate of lead* is at once astringent and sedative. It has been recommended by Beigel and by Fieber in the advanced stages of acute catarrhs, in the proportion of three to ten grains to the fluidounce.

NARCOTICS.—When there is great laryngeal or even bronchial irritation, especially in the *laryngitis* of advanced *phthisis*, half a grain of morphia, or a drachm of tincture of hyoscyamus, will often afford very great relief. It must not be forgotten that the constitutional effect of the narcotic may be produced in this way.

SOLVENTS.—As is well known, in certain respiratory affections it is very important to get rid of a false membrane, or exudation, which appears upon the mucous surfaces. Some substances seem to exert at least a degree of solvent power upon this material. Of these, lime-water appears to be the most effective and the least irritant. It should be used pure. (See article on Lime (*post*); also consult Meigs and Pepper, *Diseases of Children*, Phila., 1874, p. 680.)

ALTERATIVES.—In chronic *bronchitis*, and even in chronic *catarrhal pneumonia*, or *phthisis*, balsamic vapors have long been used, but atomization is not necessary in their application. *Muriate of ammonia* may, however, be so applied: in chronic bronchial *catarrh* it is sometimes advantageous, and in acute *laryngitis* inhalations of a warm saturated solution of it are often very useful. In chronic *laryngitis*, nitrate of silver may be exhibited by atomization, but it is, I think, much better to apply it directly by means of the laryngoscope and the brush or probang. In cases of *fetid expectoration*, carbolic acid in weak solution (gtt. i–ii to fʒi) may be used as an antiseptic and alterant.

NAUSEATING EXPECTORANTS.

The three nauseating expectorants are ipecacuanha, tartar emetic, and lobelia. As these substances are sufficiently discussed elsewhere in this work, it remains only to say a few words in regard to their comparative use as expectorants.

LOBELIA is used only when the inflammatory action is complicated with a tendency to spasm of the bronchial muscles. It is too powerfully depressant to be given to children with safety. The best expectorant preparation is the *Wine* (*Vinum Lobeliae*). The ordinary expectorant dose of this is twenty to thirty drops, usually given in combination every three hours. When a very decided impression is desired, as in some cases of asthma, one fluidrachm may


be exhibited every two hours, until the wished-for effect is produced. Under these circumstances, the patient should be closely watched, as lobelia in these large doses sometimes produces very alarming symptoms.

IPECACTANHA is the safest and most used, although perhaps the least powerful, of all the nauseating expectorants. It is the only one that should be given freely to children. In the early stages of all acute inflammatory conditions of the respiratory mucous membranes it is of great service. The preparation most generally used in acute *bronchitis* is the syrup, the dose of which is from twenty drops to a teaspoonful, every two, three, or four hours, according to the exigencies of the case. Any of the other liquid preparations of ipecacuanha may be employed in a corresponding dose. The official *Ipecacuanha Lozenges* (*Trochisci Ipecacuanha*) each contain about one-quarter of a grain of ipecac, and may be employed in catarrhal complaints. The troches of ipecacuanha and of morphia (*Trochisci Morphia et Ipecacuanha*) contain each the fortieth of a grain of morphia and the twelfth of a grain of ipecacuanha, and may be used when an anodyne effect is desirable.

TARTAR EMETIC is much more powerful than ipecacuanha as a sedative and nauseant, and must be used with more care. It is the most efficient of all the sedative expectorants, and affords in cases of urgency a very efficacious, although a very disagreeable, method of putting an end to an attack of acute bronchitis. I have known of a public speaker, who had risen in the morning completely overwhelmed with bronchitis, enabled to deliver his evening speech by the judicious use of tartar emetic. To effect this rapid relief, one-twelfth of a grain of tartar emetic should be taken every ten or fifteen minutes until it induces profuse vomiting. After the nausea is passed, the system may be toned up for exertion by a lunch of ale and oysters. Tartar emetic is a powerful remedy, and is especially injurious to young children, in whom it is very apt to induce collapse. It is therefore contra-indicated by infancy, and by the existence of gastro-intestinal inflammation. The dose as an expectorant is from one-twelfth to one-quarter of a grain, repeated according to circumstances.

STIMULATING EXPECTORANTS.

AMMONII CHLORIDUM—CHLORIDE OF AMMONIUM.



Syn.—*Muriate of Ammonia, Sal Ammoniac.*

This drug is prepared by heating with chloride of sodium the sulphate of ammonium, which is obtained from gas-liquor by the addition of sulphuric acid.

Gas-liquor is a water which has been used to wash ordinary burning-gas, and contains largely of the carbonate, hydrocyanate, hydrosulphate, and sulphate of ammonium. The *sulphate of ammonium* is not itself employed as

a medicine, but is officinal, because the other preparations of ammonia are made from it.

The muriate of ammonia is a white, translucent, fibrous, and tough salt, free from odor, but having a sharp, saline taste. It occurs in large concavo-convex plates, dissolves in three parts of cold and in one of boiling water, and at a red heat sublimes without decomposition.

PHYSIOLOGICAL ACTION.—When applied in a solid form, or in a concentrated solution, the chloride of ammonium acts as an irritant upon raw surfaces and upon mucous membranes. This influence seems, however, not to be sufficiently powerful to enable the drug in any dose to produce lethal poisoning: at least, Oesterlen (*Heilmittellehre*, Tübingen, 1851) states that one of his patients took two ounces of the salt without suffering any more severe results than colicky pains and some diarrhœa.

If some of the older experimenters are to be credited, its influence upon the lower animals is more powerful. In the experiments of Dr. Smith (quoted by Stillé), two drachms of the salt applied to the wounded thigh of a dog caused death in from twelve to thirty-six hours, and, according to Orfila, the same quantity dissolved in two ounces of water, and introduced into the stomach of a dog, caused violent convulsions, with great tetanic rigidity, and finally death. Arnold (Wibmer, *Die Wirkungen der Arzneien und Gifte*, Munich, 1831, Bd. i. p. 143) found that thirty grains will kill a rabbit in ten minutes. On the other hand, in the recent studies of Dr. Rabuteau (*L'Union Médicale*, 1871, p. 330), half a drachm injected into the veins of a moderate-sized dog had no apparent effect, whilst one drachm only produced muscular weakness, deepening into temporary paralysis of the hind legs, vomiting, without diarrhœa, and general prostration for four or five hours.

The chief interest of the clinician in the physiological action of the muriate of ammonia centres in its effects when given continuously for some time. The older writers upon the subject assert that its influence on the heart is a sedative one, but that it increases the capillary circulation (Sundelin, *Heilmittellehre*, Band i. p. 150, Berlin, 1853). This opinion appears to me to be founded chiefly upon speculation and inference, and not to have any sufficient basis. Whatever may be the action of very large amounts, I have never been able to perceive that the muriate of ammonia, given in ordinary doses, has any decided influence upon the circulation.

According to Sundelin and other authorities, the drug, when given freely and continuously, produces a profound impression upon the blood itself, lessening its plasticity and impairing its constitution. One case of profound and otherwise inexplicable prostration and typhoid condition, which occurred in a patient who was taking nearly a half-ounce of muriate of ammonia per diem, has come under my own notice; but, although I have given the drug very largely and freely, I have never witnessed any other evidences of its affecting the general system. On the other hand, the very elaborate analyses

of F. W. Böcker (*Beiträge zur Heilkunde*, Bd. ii. p. 170, Crefeld, 1849), although somewhat discordant, seem to indicate that sal ammoniac does impoverish the blood, since in some instances there was a decided decrease in its solid constituents. Arnold (*loc. cit.*) also noticed that in dogs poisoned with the muriate of ammonia the blood contained less than the normal percentage of solids.

In accord with this reputed action on the blood is the effect of the drug upon the urinary secretion. In a very elaborate series of experiments, Böcker (*loc. cit.*, p. 158) found that, given to a healthy man, sal ammoniac increased very notably all the solids of the urine, except the uric acid, which was very slightly diminished; and Rabuteau, in an investigation in which, by identity of diet, etc., all sources of fallacy were as far as possible excluded, found that the excretion of urea was very decidedly increased.

The opinion has long prevailed that the muriate of ammonia especially affects the mucous membranes, and Böcker believes that in them it hastens very greatly the nutritive changes and the exfoliation of epithelium. Clinical experience has, I think, demonstrated that the drug does act especially upon the mucous membranes; although it is very difficult to bring forward any definite tangible proof of this.

The muriate of ammonia, when ingested, probably enters into all the excretions, since Rabuteau (*L'Union Médicale*, t. xii. p. 329, 1871) has found it in the saliva. The same observer has shown that the chief elimination takes place through the kidneys, since he found in the urine almost all of the salt that had been taken.

THERAPEUTICS.—In the last century the chloride of ammonium was very extensively used in *intermittent fever*; but at present the salt is rarely or never employed, although Aran (*Bulletin Thérapeutique*, t. xii. p. 344) has attempted to revive the practice, and asserts that he has had very good effects resulting from it,—in thirteen cases of intermittent fever, curing, by the exhibition of two drachms a day, seven at once, four after the second paroxysm, one after the third, and one after the fourth.

The chief present use of the muriate of ammonia is as a remedy in *acute* and in *chronic bronchitis*: in the first variety of the disease the drug should not be exhibited in the first stages; but after active inflammatory action has been subdued by the sedative expectorants, it is very useful. It is very largely employed in the *catarrhs* of young children, and is often of great service.

In Germany the remedy has been extensively exhibited in *gastric* and *intestinal catarrhs*; and recently (*Indian Medical Gazette*, Aug. 1, 1872) Dr. W. Stewart, as the result of a large experience, has highly commended its use in *chronic torpor* of the *liver*, *chronic hepatitis*, and even in *hepatic abscess*. In the first of these affections I have employed it to a limited extent, with very good effects. Dr. Stewart gives twenty grains of it three times a day, and continues its use for weeks or even months.

In various *neuralgias*, especially in the *ovarian* variety, but to a less extent in *migrainæ* and other forms, *sal ammoniac* has been largely employed for the purpose of relieving pain. Thirty grains of it are given in combination with from two to five drops of tincture of aconite root, and the dose is repeated in half an hour, if necessary.

ADMINISTRATION.—The usual dose of the muriate of ammonia in catarrh is from five to ten grains three or four times a day. This dose is given in a tablespoonful of water, to which, for the purpose of concealing the taste, from five to ten grains of liquorice may be added.

SENEGA—SENEKA.

The root of *Polygala Senega*, a small, herbaceous perennial, indigenous to the Middle and Southern United States. This root is several inches long, very much contorted, of a peculiar feeble odor, and a taste at first sweetish but afterwards acrid. It is distinguished by a keel-like line, shorter than the root, and presenting the appearance as if a string were drawn tightly under the bark from end to end. Quevenne discovered in *seneka* a peculiar principle which he named *Polygalic Acid*, although he thought that it was closely allied to saponin. According to Husemann (*Die Pflanzenstoffe*, p. 751), the more recent researches of Bolley (*Ann. Chem. Pharm.*, xc., xci.) have shown that it is identical with saponin.

Saponin is a glucoside, first found in the root of *Saponaria officinalis*, but now known to exist in a great many plants. It is a white, amorphous powder, of a neutral reaction, odorless, but very irritating to the nostrils, and having an acrid, burning taste. It is very soluble in water, to which, even when in no greater quantity than one part in a thousand, it imparts the property of foaming like soap-suds, when shaken.

THERAPEUTICS.—*Seneka* is chiefly used as a stimulant to the mucous membrane of the lungs in *chronic bronchitis* and in the very advanced stages of the acute disorder. It is believed to be one of the most stimulant substances of its class, and is therefore contra-indicated by acute pulmonary inflammation and is indicated by a relaxed state of the bronchial mucous membranes. If its stimulant effect be modified by combination with tartar emetic, it may be given in acute bronchitis at an earlier stage than it could be used by itself. *Seneka* is locally irritant, and, when administered in large doses, produces vomiting, which is probably reflex in its origin. In overdoses it not only vomits, but also purges: it is, however, never used for these purposes. *Seneka* has been employed as an emmenagogue in *amenorrhœa*, and as a diuretic in *dropsy*, but has achieved no permanent reputation in these disorders. The powdered drug is sometimes given in doses of ten to twenty grains; but one of the following official preparations is preferable:

Decoction Senegæ—*Decoction of Seneka* (3i to Oj). The decoction is much improved by the addition of liquorice, but at best is a very poor preparation. Dose, one to two fluidounces.

Syrupus Senegæ.—*Syrup of Seneca*. An efficient preparation. Dose, a teaspoonful.

AMMONIACUM—AMMONIAC.

The concrete juice of an umbelliferous plant, *Dorema Ammoniacum*, a native of Persia. It is said to exude from punctures made in the plants, and also to be obtained by a process similar to that employed in the case of *assafoetida*.

Ammoniac is a gum resin, containing a little volatile oil, and occurs in commerce in two forms: as irregularly globular yellowish tears, from the size of a pin to that of a large chestnut, hard and brittle when cold, and breaking with a smooth, resinous fracture; and in irregular mottled masses, composed in part of tears, and containing many impurities. The odor is faint and disagreeable, the taste bitterish, sweetish, and somewhat acrid.

THERAPEUTICS.—The influence of ammoniac upon the general system is very slight. It is a local irritant, and therefore in large doses is capable of producing vomiting and purging. It was formerly employed as a nervous stimulant, but its internal use is now restricted almost exclusively to *chronic bronchitis* with either deficient or excessive secretion. The dose of it is twenty to thirty grains; of the officinal *Mixture* (*Mistura Ammoniaci*), a tablespoonful.

There are two officinal plasters of ammoniac: the *Emplastrum Ammoniaci*, which is used as a mild counter-irritant and as a local stimulant to promote the resolution of *scrofulous tumors* or *enlarged joints*; and the *Emplastrum Ammoniaci cum Hydrargyro*, which contains mercury and sulphur, and is employed as a local stimulant, alterative, and discentient in the same affections as the first-named plaster, and in *syphilitic nodes* and *tumors*: its external use is said to have caused salivation.

BENZOINUM—BENZOIN.

The concrete juice of *Styrax Benzoïn*, a large tree, native of Peru. The drug is said to be obtained by incising the tree and allowing the juice to harden as it exudes. The finest specimens of benzoïn consist of tears agglutinated together; the poorest, of brown or blackish masses without tears. The fracture is resinous, the surface of the tears smooth and whitish, the odor fragrant, the taste at first very slight, afterwards somewhat acrid. The chief constituents of benzoïn are resin and benzoic acid; cinnamic acid is also frequently present, and is said to be especially found in the white tears.

Benzoic Acid (ACIDUM BENZOICUM, U.S.P.) is obtained by sublimation of gum benzoïn. As thus prepared, it is in white feathery crystals, of a silky lustre and a fragrant vanilla-like odor, due to the presence of a volatile oil. Pure benzoic acid is inodorous. The taste is warm, acrid, peculiar. It melts at 250°, and volatilizes without change; is soluble in two hundred

parts of cold and in twenty-four parts of boiling water; is soluble in alcohol, fixed oil, and alkaline solutions. It is a feeble acid, but forms neutral salts with the alkalies. Benzoic acid is widely distributed through the vegetable kingdom, constituting the peculiar principle of all true balsams, and is occasionally present in the urine of grass-eating animals. It is a normal constituent of castor, and has been detected by Seligsohn (*Chemische Centralblatt*, 1861) in the suprarenal capsules of an ox.

Benzoic acid is used considerably in the arts, and for this purpose is prepared from the allied hippuric acid of horse-urine, and also, it is said, from naphthalen.

PHYSIOLOGICAL ACTION.—Locally applied, benzoic acid is a decided irritant, and its vapors, when inhaled, produce bronchial catarrh.

Especial interest attaches to the elimination of benzoic acid, because, as was first discovered in dogs by Wöhler, and afterwards in man by Ure, it appears in the urine, united with nitrogenous atoms, as hippuric acid. It becomes a matter of great interest to know where this change occurs; but this has not as yet been certainly determined.

Kühne and Hallwachs thought that the conversion occurred in the liver; but the more recent researches of Meissner and Shepard (*Untersuchungen über das Entstehen der Hippursäure in thierischen Organismus*, Hanover, 1866) appear to show that it really takes place in the kidneys. Certainly it does not happen in the intestines or in the blood, since after the exhibition of large doses of benzoic acid it alone can be detected in the blood; and after the administration to rabbits by the mouth of large amounts of hippuric acid, only traces of the latter, with large quantities of benzoic acid, can be found in the blood, although the hippuric acid reappears in the urine; further, moderate amounts of hippuric acid injected into the blood cause severe symptoms of poisoning, which is not true of benzoic acid. When very large amounts of benzoic acid are introduced into the blood, a portion escapes through the kidneys unchanged. According to Meissner and Shepard, sometimes the benzoic acid is converted into succinic acid instead of hippuric acid in man, and in chickens it is habitually changed into two new products, one of which is nitrogenous.

The point as to whence the nitrogen necessary for the formation of the hippuric acid is obtained has not yet been clearly made out. The testimony as to the effect of the ingestion of benzoic acid upon the urea and uric acid of the urine is singularly contradictory. Thus, Ure (*Medico-Chirurgical Transactions*, xxiv. p. 30, 1841), Leroy d'Etiolles (quoted by Stillé), and Debouy (quoted by Stillé) all affirm that the uric acid is very much diminished or altogether absent, whilst Garrod (*Memoirs of the Chemical Society*, i., 1842, and *London Lancet*, Nov. 1844) and Keller (*Ann. der Chem. und Pharm.*, xliii., 1842) assert that its quantity remains normal. Again, Garrod affirms that the urea is very much diminished in quantity, whilst Keller and Meissner and Shepard (*loc. cit.*) declare that it is not affected. The only

logical conclusion would seem to be that the effect of benzoic acid upon the urine is variable. All authorities appear to agree, however, in asserting that the acidity is increased.

Upon the general system benzoic acid exerts no apparent influence, except it be taken in enormous doses; and even then its action is very slight. Thus, Schreiber took in two days about half an ounce of the acid, and suffered only a feeling of abdominal warmth, spreading over the whole body, and accompanied by an increase of the pulse-rate amounting to thirty beats per minute, by increased secretion and excretion of phlegm, with slight disturbance of digestion. Owing to its irritant effect, large doses of benzoic acid not rarely nauseate or derange the stomach.

THERAPEUTICS.—Benzoic acid, or gum benzoin, was formerly very freely exhibited in *chronic bronchial catarrh* as a stimulant expectorant; at present it is not much employed. The chief use of it is in urinary affections. Dr. Ure first suggested and commended its employment in uric acid *gravel* and *calculus*, because, as he thought, it diminishes the excretion of uric acid: as its possession of this power has been strenuously denied, the theory of its use in uric acid gravel is exceedingly doubtful. No less an authority, however, than Dr. Golding Bird* asserts that clinical experience has shown its value in *uric acid diathesis*. In the *phosphatic urine* of *vesical catarrh*, benzoic acid may be employed with much advantage. By rendering the urine more acid it increases its power of dissolving the phosphates, and at the same time it exerts a stimulant action upon the mucous membrane of the bladder.

Benzoic acid has the property of preventing animal fats from becoming rancid, and is therefore much used as an addition to ointments. Moreover, it exerts a peculiar, often very beneficial, stimulant action upon the skin, and is very useful in such conditions as *chapped* hands, lips,† or nipples, and even in *fissure* of the *anus*.

ADMINISTRATION.—The dose of benzoic acid is from ten to thirty grains. Gum benzoin is never used itself, but the following preparations of it are official:

Tinctura Benzoini—*Tincture of Benzoin* (℥vi to Oj). Dose, f℥ss to f℥i.

Tinctura Benzoini Composita—*Compound Tincture of Benzoin*. Used in *chronic bronchial catarrh*. Dose, f℥i to f℥ii.

Unguentum Benzoini—*Benzoin Ointment* (Tinct. Benzoini, f℥ii to ℥xviii).

BALSAMUM PERUVIANUM, or *Balsam of Peru*, is obtained from the *Myrospermum Peruiferum*, a tree of Central America, by making incisions

* *Urinary Deposits*, Phila., 1859, p. 160. He states that he has found the following formula of great service in chronic uric acid gravel:

℞ Sodæ carbonatis, 5iss; acidi benzoici, gr. xl; sodæ phosphatis, 5iii; aquæ ferventis, f℥iv; solve et adde aquæ cinnamomi, f℥viiss; tinct. hyoseyami, f℥iv. S.—Two tablespoonfuls three times a day.

† Prof. Stillé commends a mixture of one part of the compound tincture and four parts of glycerine.

in such places as have been previously beaten with clubs, slightly burning them, catching the juice in old rags, and finally boiling these in water and skimming off the balsam as it rises to the surface.

This balsam is a viscid, honey-like, fragrant, brownish fluid, of a warm, bitterish taste. According to Frémy, it contains not benzoic, but cinnamic acid. It has been used in chronic catarrhs of the respiratory and the genito-urinary system, in doses of half a fluidrachm.

BALSAMUM TOLUTANUM, or *Balsam of Tolu*, is obtained from a tree very closely allied to that which yields the balsam of Peru, if indeed it be not identical with it. The incisions, however, are not burnt, and the juice is simply caught in vessels.

Balsam of Tolu is at first a thick, viscid fluid, but by time it is converted into a hard, translucent, resinous solid. Its odor is highly fragrant, and its taste vanilla-like. It contains *cinnamic acid* and a volatile oil, and its medical properties are the same as those of the balsam of Peru. On account, however, of its grateful taste, it is preferred to the latter, and is very much used to flavor medicines, especially cough-mixtures. In large doses (grains xx to xxx every three hours) it may be of some value in chronic catarrh, but as generally used its preparations are simply agreeable vehicles. The dose of the tincture (*Tinctura Tolutanæ*) is half to one fluidrachm; of the much more frequently used syrup (*Syrupus Tolutanæ*), half a fluidounce.

ALLIUM, or *English Garlic*, the clove of the *Allium sativum*, contains a volatile oil, which is a stimulant in small doses to digestion, and is also a stimulating expectorant often of very great service in certain stages of *bronchitis*. It is in lingering, deep-seated "colds" that I have derived especial benefit from its use. It is also very valuable in the *acute bronchitis* of infants, when the powers of the system begin to give out. The oil of garlic is further believed to have the ability to stimulate the expulsive function of the small bronchial tubes, and is certainly a powerful rubefacient and a decided nervous stimulant. For these reasons, garlic poultices are a favorite application in the *acute suffocative catarrh* of infants, and are not rarely applied to the spine, legs, and feet in general *infantile convulsions*. They are made by simply reducing the garlic to a pulp by pounding. When a continuous application is desirable to the delicate skin of an infant, as in catarrh, it is generally necessary to reduce their strength with flaxseed meal.

SCILLA, or *Squill*, is one of the most used and most efficient of the stimulating expectorants, coming especially into play in the advanced stages of ordinary *bronchitis*. The syrup (*Syrupus Scillæ*) is the favorite expectorant preparation. As it contains acetic acid, it is incompatible with the carbonate of ammonium. The *Compound Syrup of Squill* (*Syrupus Scillæ Compositus*; squill, seneka, āā ꝑiv; tartar emetic, grs. xlviii to Oij) contains one grain of tartar emetic to the ounce, and is therefore sedative to the cir-

eculation although stimulant to the bronchial mucous membrane. It is, of course, in large doses a powerful emetic, and, under the name of *Coxe's Hive Syrup*, is very frequently used in the domestic treatment of croup. The dose of the simple syrup is one-half to one fluidrachm; of the compound, twenty to forty drops; as an emetic to children, ten drops to a fluidrachm, according to age, repeated every twenty minutes until it operates.

PIX LIQUIDA.—*Tar* is a black semi-liquid substance, of peculiar odor and taste, obtained by the destructive distillation of various species of pine. The tar used in this country is almost exclusively the product of the *Pinus palustris* of North Carolina and other of the Southern States. In composition it is very complex, containing pyroligneous acid, creasote, empyreumatic oil, and a number of more or less peculiar principles. When distilled, it yields an oily liquid, known as *oil of tar*, and a solid, black residue, *pitch*. It is freely soluble in alcohol, ether, and the fixed and volatile oils, and also to a slight extent in water.

PHYSIOLOGICAL ACTION.—As tar contains a notable proportion of creasote, if taken in sufficient quantity it is capable of exerting the peculiar influence of that agent upon the system. But creasote is not the only active principle in it: hence tar differs from that drug in acting more than it does upon the mucous membranes. That tar is capable of acting as a poison is shown by the case reported by Taylor (*Principles and Practice of Medical Jurisprudence*, 2d ed., vol. i. p. 334), in which death resulted in a man from taking by mistake the *oil* of tar. To cause death, tar itself would have to be ingested in enormous quantity, since a sailor (according to Stillé) is said to have recovered after taking between a pint and a quart of it.

Applied to any part, tar acts as a very decided stimulant.

THERAPEUTICS.—Tar is used internally almost solely as a stimulant expectorant in the advanced stages of obstinate *acute bronchitis*, or in chronic *bronchitis*. Its chief use in medicine is in chronic diseases of the skin, as a stimulant application in the form of the officinal ointment (*Unguentum Picis Liquidæ*, equal parts). In many cases this is too severe, and the strength must be reduced. Prof. Hebra states, in his work on diseases of the skin, that if it be applied too freely, enough of the tar may be absorbed to darken the color of the faeces and the urine, and even to cause gastric irritation and black vomit. *Tar Water* (*Aqua Picis Liquidæ*) may be used internally, in doses of half a fluidounce to one fluidounce. But a better, although not officinal, preparation is the *Syrup of Tar*, which may be administered in half-ounce doses. This syrup may be made as follows: Tincture of tar (℥ii to Oj), f℥ii; carbonate of magnesium, ℥i or q. s.; white sugar, lb. i. Rub the tincture thoroughly with the carbonate, add half a pint of water gradually, then filter, and when the liquid ceases to pass pour water into the filter till the product measures half a pint; lastly, dissolve the sugar by means of a gentle heat.

CLASS VI.—EMMENAGOGUES.

EMMENAGOGUES are medicines which are employed to promote the menstrual flux. As the stoppage, scantiness, or non-appearance of this secretion arises from very different causes, and as these causes are of diverse or even opposite natures, and may often be removed by drugs, it is obvious that very many remedies of very different character are indirect emmenagogues. Thus, amenorrhœa may depend upon plethora, or it may be the result of anæmia; and whilst in the one case depletory medicines are indicated, in the other case tonics are no less essential. Besides these indirect emmenagogues, there are other substances which appear to act directly as stimulants to the uterine mucous membrane; and indeed it is probable that many of the indirect emmenagogues possess more or less of this power. The emmenagogues may be conveniently arranged in three groups: the tonic emmenagogues, the purgative emmenagogues, and the stimulant emmenagogues.

TONIC EMMENAGOGUES.

IRON is the most prominent member of this section of the emmenagogues. By far the larger number of cases of *amenorrhœa* are associated with, if not dependent upon, anæmia, and are benefited by the use of iron. It should be given in full tonic doses until the anæmia is relieved or the powerlessness of the remedy to effect such change is demonstrated. It is very rarely proper to rely solely upon the iron, which in the great majority of instances should be combined with more decidedly active emmenagogues.

MYRRH has some reputation as a tonic emmenagogue, but, as it is always employed in combination with other more active medicines of its class, the rôle it plays is somewhat uncertain. It should be employed in atonic uterine conditions, and is said to be especially valuable when chronic pulmonary complications exist. The preparations of it most used in amenorrhœa are the compound pills of iron, the compound mixture of iron, and the pills of aloes and myrrh.

PURGATIVE EMMENAGOGUES.

ALOES is believed by some to act as an emmenagogue solely by virtue of its stimulant action upon the rectum, but it very probably directly affects the uterine mucous membrane. Be this as it may, it is a stimulant emmenagogue, especially useful when *atonic amenorrhœa* exists with constipation. Ordinarily it should be given in repeated doses (three times a day) of such size as will produce daily one or two soft and semi-liquid stools. At the menstrual period advantage may be sometimes derived from the administration of a full purgative dose. It is almost always given in combination, especially with iron, whose tendency to constipation it obviates. In *plethoric amenorrhœa*, when torpidity of the bowels is present, salines, and not aloetic purgatives, should be employed.

BLACK HELLEBORE has been used by some as a purgative emmenagogue, but is now very rarely if ever employed. From twenty drops to a fluidrachm of the tincture may be given three times a day.

STIMULATING EMMENAGOGUES.

With the exception of guaiac, of parsley, and of one drug of animal origin, the medicines of this class depend upon a volatile oil for their virtue, and are very closely allied in their physiological and therapeutic action.

SABINA—SAVINE.

The dried tops of *Juniperus Sabina*, a juniper, native of the south of Europe and the Levant, but very similar to our native species, the *Juniperus Virginiana*, or red cedar. The active principle is a pale or dark yellow, when highly rectified colorless, volatile oil, which has a strong terebinthinate odor and burning taste, and is officinal (*Oleum Sabinæ*).

PHYSIOLOGICAL ACTION.—The oil of savine is a powerful local stimulant, causing burning and redness when applied to the skin, and is capable of producing fatal gastro-intestinal inflammation. Taken internally in minute doses, its effects are confined to a sense of warmth, with perhaps some ill feeling in the stomach, and slight acceleration of the pulse. After larger amounts, the arterial excitement is more pronounced, and is accompanied by an increased frequency of urination, and sometimes, also, by an actually increased flow of urine. The symptoms induced by poisonous doses are: severe abdominal pain; incessant vomiting and bloody purging; diminution or even suppression of the urine, which is often albuminous and bloody; disordered respiration; symptoms of disturbed innervation, such as unconsciousness, stertorous breathing, convulsions or convulsive tremblings; the scene closing by death in collapse. In pregnant females, abortion, accompanied by violent flooding,

almost always occurs before the fatal issue. After death, signs of gastrointestinal inflammation are generally present, but in some instances these are wanting, and in one case reported by Dr. Letheby (*London Lancet*, 1845) pulmonary apoplexy and congestion of the brain were the chief lesions.

THERAPEUTICS.—Savine is a powerful stimulant to the uterine system, and may be used in small repeated doses in atonic *amenorrhœa*. Its powers in *menorrhagia*, dependent upon a relaxed state of the uterine tissues, are even more pronounced. Its use as an *abortifacient* is accompanied by the gravest dangers to life. In uterine disease of a sthenic type savine is strongly contra-indicated. The oil is the only preparation of it which should be used. The dose of it in *amenorrhœa* or in uterine hemorrhage is from five to ten drops, repeated every three or four hours.

RUTA—RUE.

The leaves of *Ruta graveolens*, or common garden rue, an undershrub of the south of Europe. Rue has a strong peculiar odor, a warm, bitter, acrid taste, and is dependent for its medical properties upon a peculiar volatile oil, although it also contains a crystalline neutral body, *Rutin*.

PHYSIOLOGICAL ACTION.—The influence of rue upon the system is similar to, but less decided than, that of savine. Locally it is an irritant, producing, when applied to the skin persistently or in a concentrated form, such as the oil, burning, redness, and vesication. According to M. Hélié, taken internally, in large doses, it causes violent gastric pains, excessive and sometimes bloody vomiting, profuse salivation and swelling of the tongue, great prostration, confusion of mind, and convulsive twitchings, with, in pregnant women, abortion. Rue has been, and probably still is, used in Europe for the production of criminal abortion, and, although its use for this purpose certainly endangers life, I have met with no record of a fatal case. Indeed, the only death from rue that I am cognizant of occurred in a man weakened by dysentery (case, Dr. G. F. Cooper, *Medical Examiner*, N. S., vol. ix. p. 720).

THERAPEUTICS.—Rue possesses, to a considerable extent, the properties of the aromatics, and has been used considerably as a carminative. Like the preceding drug, it is useful both in *amenorrhœa* and in *menorrhagia* when dependent upon uterine atony; and especial advantage has been claimed for the combination of it with savine. The oil is the only proper preparation, and may be used in doses of from three to six drops, similarly to the oil of savine.

PETROSELINUM—PARSLEY.

The root of the *Petroselinum sativum*, or common parsley. It contains a peculiar, non-nitrogenous, liquid, neutral principle, *Apiol*, which resembles somewhat the fixed oils, but is not saponifiable; a glucoside, *apiin*, and a volatile oil.

PHYSIOLOGICAL ACTION.—The volatile oil of parsley has probably the same physiological and therapeutic value as the more ordinary essential oils, and according to Mitscherlich very large quantities of it (half an ounce) will produce death in the rabbit, largely, no doubt, owing to its local irritant action. The chief interest of parsley to the physician centres in apiol. According to the discoverers of this principle, MM. Joret and Homolle (*Journal de Pharmacie*, 3e série, xxviii. p. 219), one gramme of it will produce in man a cerebral excitation very similar to that induced by coffee, without other symptoms. In doses of from two to four grammes it causes a species of intoxication, with vertigo, ringing in the ears, and severe frontal headache,—a group of symptoms very similar to those seen in cinchonization.

THERAPEUTICS.—Apiol was introduced by its discoverers as a remedy in intermittent fever, over which, they asserted, it exercised a control secondary only to that of quinia. A commission of the Paris Society of Pharmacy reported that it would cure about half the cases of quotidian and tertian, but was powerless against the quartan; also, that a return of the paroxysm was more common after apiol than after quinia. Clinical experience subsequent to these experiments has, I think, demonstrated that the drug has some power over *malarial disease*, but is very inferior to quinia. In one or two cases in which the latter could not be employed on account of the idiosyncrasies of the patient, I have used apiol very successfully.

Joret and Homolle also employed the drug in *intermittent neuralgia*, and in *amenorrhœa* as well as in *dysmenorrhœa*. Joret (*Bulletin Thérapeutique*, Feb. 1860) has recommended apiol very highly in the last two affections, and his results have been confirmed by Marotte (*Ibid.*, t. lvi., 1863) and by some other writers. When there was very decided plethora, the apiol was thought by Marotte not to be so efficacious as at other times. In any case of amenorrhœa dependent upon or associated with anæmia or other systemic vice, the continuous administration of iron, tonics, or other suitable medicines must not be neglected. The apiol is given not between the menstrual periods, but just before the latter. Joret and Homolle believe that small doses (three grains twice a day) of it should be exhibited for the week preceding the expected return of menstruation. If any symptoms of the menstrual molimen appear, fifteen grains of it may be administered in the course of a few hours; or they may be given daily for two or three days at the expected time. In intermittent fever, the same quantity may be exhibited four or five hours before the expected paroxysm. On account of its exceedingly disagreeable taste, apiol is always administered in capsules, one of which, as imported from France, usually contains the fourth of a gramme (grs. 3.9).

CANTHARIDES is a very decided uterine stimulant, and is much used as an ingredient of emmenagogue mixtures. From three to five drops of the tincture may be given three times a day: if no unpleasant symptoms arise, the dose may be cautiously increased to eight drops, the production of *strangury* being of course sedulously avoided.

GUAIAIC, as an emmenagogue, is much less stimulating than cantharides, and is believed by some to be especially useful in *rheumatic dysmenorrhœa*. In this affection, full doses of the ammoniated tincture should be given. The following formula, adapted from one of Prof. Dewees, and known as *Dewees' Emmenagogue Mixture*, I rely upon almost exclusively in the treatment of simple atonic *amenorrhœa*. The proportion of the various ingredients should be varied to suit the exigencies of individual cases. Thus, the amount of iron should be altered according to the extent of the anæmia; of the aloes, according to the state of the bowels; of the cantharides, according to the susceptibility of the urinary organs:

℞ Tincturæ ferri chlorid., fʒiii; tincturæ cantharidis, fʒi; tincturæ aloës, fʒss; tincturæ guaiaci ammoniatæ, fʒiss; syrupi, q. s. ad fʒvi. S.—Tablespoonful three times a day.

CLASS VII.—OXYTOCICS.

OXYTOCICS are those remedies which are employed during or directly after parturition, to increase the uterine action. It has been asserted that the sulphate of quinia is a member of this class; but, as this subject has been already fully discussed, no more will be here said about it. The power of stimulating the uterine pains has been claimed for various drugs, but at present there is only one substance really used for this purpose; and instead of discussing under the general heading the indications for and the dangers attending the employment of oxytocics, I shall do it in the article upon Ergot.

ERGOTA—ERGOT.

Ergot is a blackish body, one to two inches in length, irregularly cylindrical, grooved along one side, and very generally curved. When examined under the microscope, it is seen to be composed of very thick-walled cells, which contain oil-drops but no starch. Various opinions have been advanced in regard to the nature of this body; but as by the researches of Tulasne (*Mémoire sur l'Ergot des Glumacées, Annales Scien. Natur., Botan.*, 3e série, t. xx., 1853) it has been definitely determined exactly what it is, I shall not occupy space with a discussion of the older views.

Among the lowest of vegetable organisms, and distinguished from all other plants by the absence of chlorophyll, are the fungi. There are in most cases two distinct states or stages in the life of a fungus: in the first of these, the vegetative period, it exists as a *mycelium*, a usually filamentous mass or flocculus, whose sole function is to grow and increase; in the second stage the *thallus*, or ordinary fungus or mushroom, is formed; to it is assigned the function of developing reproductive bodies, after whose formation it perishes. Between these stages there is in some fungi an intermediate one, in which the plant exists as a *sclerotium*. The genus *Claviceps* comprises a number of parasitic fungi, which develop in the pistils of the various species of Gramineæ. The officinal ergot is the sclerotium of the *Claviceps* (*C. purpurea*, Tulasne) which infests the grain of the *Secale cereale*, or common rye. The first appearance of the fungus is during the earliest life of the pistil, at the base of which there arises a minute flocculent mass of mycelial filaments. These filaments, continually growing and invading all parts of the tissue of the pistil, at last

form of it an irregular whitish body, at the base of which after a time appears a dark-colored body, the sclerotium, which continues to grow, lifting up the diseased and withering mass formed out of the original pistil, and finally developing into a perfect ergot.

If a fresh, living ergot be placed in a damp, warm place, after a time little cracks will appear in its surface, and out through these cracks little round bodies will project, and finally be raised up on stalks and constitute perfect thalli,—minute fungi, which finally produce spores.

The ergot is an exceedingly complex body, containing nearly thirty-five per cent. (M. Legrip) of a fixed oil, and, according to recent analyses, three alkaloids and a peculiar acid, *Ergotic Acid*.

Propylamia, to which very probably the drug owes its peculiar odor, is a volatile alkaloid, which was first found in ergot by Dr. F. L. Winckler. It exists also in herring-pickle, and probably contributes but little to the medical properties of the drug.

In 1864, Mr. Wenzell (*American Journal of Pharmacy*, May, 1864) announced the existence of two alkaloids in ergot, *Ecbolina* and *Ergotina*; and his discovery has been confirmed by the German chemists (see Dr. T. C. Hermann, *Büchner's Repertor. für Pharm.*, 1871). As we have almost no knowledge concerning the action of these alkaloids upon the system, and as they are separated from the drug with such difficulty as to cause them to be simply chemical curiosities, I shall say very little about them. Mr. Wenzell believes that ergot owes its ecbolic properties to ecbolina; but Eugene Haudelin (*Schmidt's Jahrbücher*, Bd. clv.) denies that these alkaloids are the active principles of the drug, because he has found a watery solution efficient after precipitation with tannic acid or corrosive sublimate.

PHYSIOLOGICAL ACTION.—In ordinary therapeutic doses, ergot causes no immediate perceptible symptoms; but when a sufficient amount is exhibited, it acts as a poison both upon man and animals. Before considering its action when given in small quantity, I shall discuss the toxic effects of the drug.

According to Diez (quoted by Stille), the principal effects of poisonous doses of ergot are in the lower animals profuse salivation, vomiting, dilatation of the pupils, hurried breathing, frequent pulse, cries, trembling, staggering, paraplegia, sometimes diarrhœa, sometimes constipation, prostration, urgent thirst, convulsions,* and death. Mr. Saml. A. Wright, in a series of experiments (*Edinburgh Medical and Surgical Journal*, Oct. 1839, vol. lii.), noted, when the medicine was given by the mouth, symptoms similar to those just spoken of: the paralysis was much more marked than the spasms. Late in the poisoning, the heart's action became irregular and intermittent, and the pulsations, which had been rapid, grew slow and feeble. In some cases the special senses seemed to be destroyed, and coldness of the surface was a very prominent symptom.

* Pereira (*Materia Medica*, 3d American edition, vol. ii. p. 137), on the authority of Phœbus, states that in the experiments of Diez convulsions were not present.

Mr. Wright also injected strong infusions of the drug directly into the torrent of the circulation. Death was in some cases produced in nine minutes, the symptoms being immediate dilatation of the pupils, great increase in the rate of the cardiac pulsations, paralysis, and convulsions.

When the fatal result was not brought about in so short a space, great anaesthesia of the surface was noted a considerable time before death; coldness of the surface and paralysis of the special senses were also present in some cases. In Dr. Kersch's experiments (*Betz's Memorabilien*, vol. xviii.*), the concentrated infusion was injected into the jugular vein; the coldness of the surface was especially noted, and also great muscular rigidity. Upon rabbits, according to the researches of Wright, ergot acts very feebly. Upon birds, as represented by chickens, turkeys, and pigeons, it causes symptoms analogous to those produced in mammals, as is testified to by Tessier and by Gross, both quoted by Stillé, and by Bonjean (*Traité de l'Ergot de Seigle*, Paris, 1845).

Enormous doses of ergot are required to produce toxic symptoms in animals, since in one of Wright's experiments an amount equivalent to two drachms for every pound weight of the dog, failed to kill.

Upon man the toxic influence of ergot is also very slight, and, although I have given the fluid extract in ounce doses, I have never seen it cause any distinct symptoms. There are at least two cases on record (Neubert, *Journal für Pharmakodynamik*, Bd. ii. p. 483, 1860; Tardien, *Ann. d'Hyg.*, 1855, vol. i.) in which fatal abortion has been produced by ergot; but I know of but a single instance of decided poisoning in a non-pregnant person. In this (Dr. G. S. Oldright, *Canada Medical Journal*, 1870, p. 404), two hours after taking the drug (amount not stated) there were developed tingling in the fingers and feet, cramps in the legs, arms, and chest, with dizziness and weakness; the pupils were dilated, the pulse was very small, and a feeling of coldness was complained of. These symptoms were relieved by the administration of stimulants and the use of external heat; after a time they recurred with greater violence; finally, under the reinstitution of the measures previously employed, the face became intensely congested and purplish red; pain in the head was felt, the patient seemed much excited,† and convulsions were feared, but did not occur; there was some diarrhoea, with dark-gray stools. In the case recorded by Neubert (*loc. cit.*), the great coldness of the surface was especially noted; and as this symptom has been so commonly remarked in ergotized animals, it probably is characteristic of poisoning by the drug.

The above summary of the general symptoms caused by poisonous doses of ergot shows that the phenomena are mainly paralytic in their nature; but, although an enormous amount has been written about the drug, we have very little knowledge as to the immediate causes of the paralysis.

* Unfortunately, I have not had access to Kersch's paper, and know it only by an abstract in *The Medical Times of Canada*, vol. i.

† These symptoms were very probably caused by the large quantities of alcohol taken.

Since Wright found (*loc. cit.*, pp. 320, 321) that the muscles responded well to galvanism after death, it would seem that the nervous system must bear the brunt of the poison. Further, Eugen Haudelin is said to have shown that the peripheral nerves are not affected:* if this be true, ergot must influence the nerve-centres.

The chief interest to the therapist in regard to the physiological action of ergot centres upon its influence on the circulation, especially on the blood-vessels, and upon its action on the impregnated uterus.

I shall discuss these points *seriatim*. Before doing so, it would be, perhaps, best to speak of the chronic poisoning by ergot; but, as any deductions from the symptoms of ergotism as to the nature of the physiological action of the drug would be at best only inferences, I shall defer the consideration of the subject to the section on toxicology.

Action on the circulation.—Although the heart is profoundly affected in acute poisoning by ergot, yet death is probably not due to this cause, since Wright (*loc. cit.*, p. 320) found that after death, even though the heart was quiet, it commenced to beat so soon as the congestion was relieved by an incision, and continued to pulsate for fifteen minutes.

The only careful study of the action of ergot upon the heart is that of P. Eberty (abstracted, *Schmidt's Jahrbücher*, Bd. clviii. p. 127). He found that in the frog the injection of a gramme of ergotin caused a diastolic arrest of the heart, and that the viscus was unable to respond at all to stimuli. It is inconceivable that this can be due to other than a direct action of the drug upon the cardiac muscle; yet Eberty seems to believe it is caused by an influence exerted through the pneumogastrics, and it is said that after division of these nerves even very great quantities of ergotin are powerless to produce cardiac arrest.

In man, full doses of ergot unquestionably diminish the frequency of the pulse, since the phenomenon has been independently noted by Parola, Gibbon, Arnal, Hardy, Beatty (all quoted by Prof. Stillé), and by Profs. Bailly and Sée (*Bulletin Thérapeutique*, t. lxxviii. p. 435); but the method in which this reduction is brought about is uncertain. The amount of the reduction varies from ten to thirty-five beats per minute; but very rarely is the pulse reduced below sixty even by the largest doses.

According to Eberty's experiments, therapeutic doses of the drug produce in mammals, as in man, slowing of the heart's beat. It was found that in frogs the pulsations of the heart were still affected after destruction of the medulla, but that in mammals, after paralysis of the peripheral vagi by atropia, ergot was powerless to alter the cardiac rhythm. These experiments, if correct, appear to prove that ergot acts as a stimulant to the peripheral cardiac nerves, and that the reduction of the number of beats is due to this, and is independent of the nerve-centres.

* *Inaug. Dissert.*, Dorpat, 1871. I have seen only an abstract, contained in *Schmidt's Jahrbücher*, Bd. civ. p. 142.

As early as 1827, M. Courhant advanced the opinion that ergot produces a spasm of the blood-vessels; but, although his theory was very generally adopted, only within a very few years has any earnest attempt been made to prove or disprove its correctness.

In 1870, Dr. Ch. L. Holmes (*Archives de Physiologie*, t. iii., 1870) found that when the blood-vessels of the frog's web were watched under the microscope and the animal poisoned either with the aqueous extract of ergot or with the powdered drug, the vessels, both venous and arterial, could be seen to undergo a very great contraction. Dr. A. Wernich (*Virchow's Archiv*, 1872, Bd. lvi. p. 510) observed that when rabbits in which the arteries of the thigh, back, pia mater, etc., were exposed, received full doses of ergot, these arteries could be seen to undergo a very remarkable diminution in their calibre.

These observations have been confirmed by other observers, among whom may be mentioned Vogt, Dr. S. Kersch (*loc. cit.*), and Boldt (*Schmidt's Jahrbücher*, March, 1872). The latter observer also affirms that there ran through the ergotized capillaries wave-like, peristaltic spasms. Patrick Nicol and J. Mossop (*Brit. and For. Medico-Chir. Rev.*, vol. l., 1872) have noted with the ophthalmoscope the contraction of the retinal vessels after the exhibition of ergot in man.

If general vaso-motor spasm be produced by ergot, unless the heart's action be greatly weakened by the drug there must be a very decided rise in the arterial pressure. The question as to the existence of this rise was first investigated by Dr. Ch. L. Holmes (*loc. cit.*). This experimenter injected the aqueous extract directly into the jugular vein of the dog, whilst the manometer or the kymographion was connected with one of the main arteries. The phenomena induced were immediate, rapid, and very decided depression of the arterial pressure, with violent cries and efforts on the part of the dog, followed by a period of quiet and a rise of the arterial pressure above the normal point. Twenty-three experiments were performed: in two of them the first depression of the arterial pressure was wanting; in six of them the arterial pressure did not rise above normal.

As the experiments are not reported in detail, it is not possible to point out in all the cases the causes of these exceptional phenomena; but Dr. Holmes mentions sickness of the animal, death during the stage of depression, or some other obvious reason for the departure from the usual results, as being present in nearly all the cases, so that the exceptions are, I think, at present best disregarded in the interpretation of the phenomena.

The unexpected result in the investigation of Dr. Holmes was the primary depression of the arterial pressure, a depression which at first sight seems incompatible with the idea that ergot contracts the vessels. It is, of course, possible that the fall of pressure may be due to an intense action of the drug upon the heart; but clinical evidence, and the secondary rise and other results of Dr. Holmes's experiments, are opposed to the idea that the drug in the

doses used has any such power. Dr. Holmes tried the effect of the injection of the ergot after section of the cardiac nerves, but obtained results so varying and contradictory that he could come to no other conclusion than that the alterations of the arterial pressure produced by ergot are not dependent upon its cardiac action. In the absence of details of these experiments, we can but accept this conclusion. To account for the phenomena following the injection, Dr. Holmes propounds a theory which is very plausible, but which he certainly does not prove to be true. His idea is that the first fall of the arterial pressure is due to a spasm of the pulmonic capillaries, hindering the blood in its passage to the left heart, and thereby causing venous repletion and arterial depletion. It is to be remembered that he injected the ergot directly into the jugular vein, so that the whole force of the remedy fell directly upon the pulmonic circulation. It is evident that this matters very little if ergot produces vaso-motor spasm only through an influence upon the nerve-centres, but that it does make a material difference if the contraction of the vessels is the result of a local action. Holmes asserts that after section of a sympathetic nerve and injection of the ergot the vessels supplied by the divided nerve can be seen to contract; and Wernich confirms this observation. If ergot does thus cause vaso-motor spasm, it is plain that in Dr. Holmes's experiments this spasm must have been very intense in the lungs before the systemic capillaries were affected. Our investigator proved by experiment what *a priori* seems necessarily true, that if blood be prevented from passing freely to the lungs the arterial pressure falls very greatly. In this connection it is worthy of notice that the period of arterial depression following the injection of the drug into the jugular vein was very brief. In the only experiment detailed in full by Dr. Holmes, the pressure began to rise in three minutes after the completion of the injection; in five minutes it was nearly normal, and in eleven minutes was above the normal point.

It is evident that if the asserted facts are all true, the theory of Dr. Holmes is probably correct. It is, however, very doubtful whether the ergotic spasm is local in its origin. Evidence derived from judging by the eye as to whether a vessel does or does not contract must, I think, always be taken *cum grano salis*, and the observations of Holmes and of Wernich are directly contradicted by some apparently careful and elaborate experiments of Dr. Paul Vogt (*Berliner Klinische Wochenschrift*, 1869, No. xii.), in which the dilated vessels of the ear of a rabbit whose cervical ganglion had been extirpated could not be made to contract by the hypodermic injection of ergot.

Seemingly the most elaborate and thorough examination yet made as to the action of ergot upon the blood-vessels, as evinced by its influence on arterial pressure, is that of P. Ebert (*Inaug. Dissert.*, Halle, 1873). His results are in accord with those of Vogt, and disagree with those of Dr. Holmes, especially in their bearing upon the question whether the contraction of the arteries is centric or peripheric in its origin. He finds that the arterial pressure rises directly and enormously after the injection of ergotin. This rise,

which he acknowledges to be chiefly due to the contraction of the vessels, occurs in the veins as well as in the arteries, and in the frog as well as in the dog and the rabbit. According to his experiments, it must be, at least in the batrachian, centric, since in the frog it does not take place after destruction of the medulla. He also found that after the inhalation of nitrite of amyl the ergot caused rise of the arterial pressure; but this certainly does not prove that the vaso-motor spasm is centric, as the abstract (*Schmidt's Jahrbücher*, Bd. clviii. p. 127), with which alone I am acquainted, makes him assume. It is a fair inference that if the vaso-motor spasm be centric in the frog it is also so in the mammal; but the only real proof of the point would be an experimental test as to its action in the higher animals after section of the cord.

Until further studies are made, no positive opinion can be arrived at as to whether ergotic vaso-motor spasm is peripheric or centric; but the probabilities seem to me to be very strong in favor of the latter origin.

The rise of arterial pressure produced by ergot, and the existence of vaso-motor spasm, must, I think, be considered as proven facts; yet Haudelin is stated to have found that the arterial pressure falls after the exhibition of the poison. Brown-Séquard has insisted (*Archives de Physiologie*, 1870, t. iii. p. 434) that in ergotic poisoning there are two periods: first, vaso-motor spasm; second, vaso-motor paralysis. It is very possible that he is correct, and that Haudelin's results were simply due to his having used such enormous doses of ergot as at once to overpower the vaso-motor apparatus.

Action on the intestines and on the uterus.—The fibres in the coats of the blood-vessels are certainly not the only non-striated muscles upon which ergot acts: indeed, the probabilities are strong that the drug influences muscular fibre of this character wherever it exists in the body.

There is considerable evidence to show that it causes increased intestinal peristalsis. In Dr. Wright's experiments (*loc. cit.*, p. 320) the intestines were found in very active peristalsis at the post-mortem examination of the poisoned animal. Dr. Wernich (*Virchow's Archiv*, 1872, Bd. lvi. p. 515) noticed that very violent peristaltic movements followed the injection of ergot, as was seen not only in rabbits whose abdomens were opened, but in some cases even through the uninjured walls; and Haudelin has confirmed his observations.

Upon the uterus of parturient women ergot exerts a very pronounced and fixed influence, increasing the length and force of the pains, and, if it be given in sufficient dose, causing after a time violent tetanic cramp of the whole organ. The drug certainly acts in this respect upon the lower animals as it does upon man, since Youatt states that in a large experience, both with monogastric animals and ruminants, he has never known the drug to fail in its action on the uterus of the parturient female.

The action of ergot in producing contraction in the impregnated but not parturient womb is by no means so certain. Upon animals, Dr. Wright found it to fail in all of a number of trials, as did also Bonjean in a single experi-

ment. On the other hand, Diez (Stillé, *Therapeutics*, 2d ed., vol. ii. p. 585), Oslere (*Ibid.*), and Percy and Laurent (*Ibid.*) found it to cause abortion in guinea-pigs, sows, rabbits, cows, and cats; and M. Bodin has reported an epidemic of abortion occurring among cows near Trois Croix, which he attributes to feeding upon ergotized grasses (*Journal des Connaissances Médicales*, 1842). The evidence of those who have used ergot for the induction of premature labor in woman tallies very closely with that which is brought forward in regard to the lower animals. As the matter is so fixed, I will not enter into an elaborate discussion of the effects of ergot upon pregnant women. To show that the fungus very often will act as an abortifacient, it is only necessary to quote Prof. Ramsbotham, who states (*The Principles and Practice of Obstetric Medicine and Surgery*, Phila., 1860, p. 318) that he has made a "great number of trials," and found that "expulsive action soon followed its exhibition, with very few exceptions." It cannot be gainsaid, however, that very often the drug has failed: sometimes, no doubt, because of poor quality or because given in insufficient dose, yet sufficiently often to show that its abortifacient action is uncertain.*

Whether the uterine disturbance is of centric origin, or is due to a direct action of the drug upon the uterus and its nerve-fibres, is not determined. Dr. Wernich, in two experiments, found that no vermicular movements were produced in the unimpregnated womb by ergot after section of the spinal cord; but in one of these experiments the animal was so feeble as to destroy any possible force the result might have, and I do not think much weight is to be attached to the evidence.

THERAPEUTICS.—As a therapeutic agent, ergot is employed both by the obstetrician and the physician; and I shall consider these uses of it separately.

Owing to the power that ergot possesses of intensifying labor-pains, it has been long used in *uterine inertia* during parturition. Indeed, it was for this purpose that the drug was first employed in medicine, and thereby acquired the name of *pulvis parturiens*. The literature of the subject is immense, and all imaginable opinions as to the effects of the drug when given in labor, and as to the advisability of its employment, have been advanced; but, without discussing these, I shall here simply point out the clearly-established rules for its use and the clinically-determined dangers and advantages of its employment. If ergot be given in very small doses during labor, the natural pains are simply intensified; but if the dose be large enough to have a decided effect, their character is altered: they become not only more severe, but much more prolonged than normal, and finally the intervals of relaxation appear to be completely abolished and the intermittent expulsive efforts are changed into one violent, continuous strain. It is evident that, if the resistance be sufficiently great, this may endanger the safety both of the mother and of

* For a more elaborate setting-forth of the matter, the reader may consult Prof. Taylor's *Principles and Practice of Medical Jurisprudence*, 2d ed., 1873, vol. ii. p. 193.

the child. The dangers to the mother are twofold: there is a possibility of the uterus rupturing itself by its efforts; and, when the head comes down upon the perineum, if the soft parts be rigid there is a very strong probability that they will be lacerated. The danger of uterine rupture is, I think, a remote one; for although several alleged cases have been recorded, yet in very few is the accident clearly traceable to the asserted cause (see *Stillé, Therapeutics*, 2d ed., vol. ii. p. 591). The fatal character of the accident is such, however, that the possibility of its occurrence should always prevent the reckless use of the drug.

The improper use of ergot is far more serious in its effects upon the child than upon the mother. During a violent uterine contraction, the passage of the blood from the placenta to the child must be interfered with, or, in other words, the respiration of the fœtus is temporarily stopped, so that its life depends upon the aeration of the blood during the intervals. If the latter be very much shortened, the life of the child is greatly imperilled; and if they be abolished, it must be destroyed, unless delivery occurs in a very few moments. These considerations are, I think, sufficient, without further discussion, to show the imperativeness of the rule *never* to give ergot in uterine inertia when there is much *resistance*, either in the bony or the soft parts of the mother. In primiparæ such resistance is always to be looked for, and its degree often difficult to judge of beforehand; and in such women ergot should not be used at all for the purposes of expulsion. Even under the most favorable circumstances for its employment—when the woman has previously borne children, when the bony pelvis is capacious, and the soft parts are relaxed and dilatable—its use should be entered upon with caution; and if the accoucheur be skillful in the application of instruments, cases must be rare in which the latter are not preferable to the ecbotic.

In women of lax fibre, with roomy pelvis, ergot may be used in uterine inertia if instruments are not at hand, or if they are objected to, or if the obstetrician is timid in their application.

At the close of parturition, ergot is very commonly employed to prevent *post-partum hemorrhage*; and in this case there is no objection to its use, and the remedy is invaluable. But, as it requires from fifteen to twenty minutes for its action when given by the mouth, ergot exhibited in this way cannot be relied upon to arrest flooding when it has already set in. To prevent the occurrence of the latter, it is an excellent rule to give a full dose of the ecbotic when the child's head is well down upon the perineum and beginning to emerge at the vulva. After labor, if a tendency to bleeding is manifested, the exhibition of ergot may be added to the other measures employed. Its hypodermic use under these circumstances would be eminently proper.

For the induction of *premature labor*, ergot has been and still is to some extent used; but it is uncertain in its action, and offers no advantages over instrumental methods.

The success of ergot in arresting hemorrhage after labor soon led to its use in uterine hemorrhages in other than parturient or pregnant women; and the next step beyond this was its employment in other hemorrhages. In all forms of hemorrhage in which no direct local application can be made, ergot is to-day probably the most generally used remedy; and even when local applications can be made, if the bleeding be obstinate, ergot is very generally exhibited. It is thus employed in *menorrhagia*, in *hæmoptysis*, *hemorrhage* from the *gums*, *epistaxis*, etc., etc. Ergot acts with especial rapidity and efficiency in these cases, if given hypodermically: used in this way, the suddenness of the result in hæmoptysis is sometimes surprising.

The value of the drug in hemorrhage is no doubt dependent upon its power of contracting the small vessels. As has already been insisted, colligative sweating is due to a relaxation affecting the capillaries, and therefore, *a priori*, ergot should be of value in the *night-sweats* of phthisis and similar diseases. Dr. Christmann states (*Centralblatt f. d. Med. Wissenschaften*, Nov. 1869) that in these cases he has seen very good effects from the use of ergot in doses of two or three drachms a day.

It was the physiological action of ergot upon the muscular fibres in the walls of the vessels that led Prof. Langenbeck to try the effect of the hypodermic injection of the remedy in *aneurism*. The result obtained, as reported in the *Berliner Klinische Wochenschrift* (vi., 1869), was surprising. How far surgeons have followed up this matter I do not know; but, according to Dr. Paul Vogt (*Berliner Klinische Wochenschrift*, March, 1872), Schneider has cured a case of femoral aneurism, and Dutoit one of the subclavian. Vogt has himself employed the remedy successfully in a very severe case of varicose veins of the legs, of many years' standing. The remedy is injected into the immediate neighborhood of the aneurism or varix, one or two grains of ergotin, or the aqueous extract of ergot dissolved in water and a little alcohol, being employed every day. A good deal of local swelling and hardness is induced, and it is certainly a question whether the good achieved has not been simply the result of the local inflammation set up.

Very many years ago, Dr. F. E. Barlan-Fontayral proposed (*Journal des Sciences médicales pratiques de Montpellier*, tomes vi., vii.) the use of ergot in *chronic dysentery* and *diarrhœa*, on account of its power of causing contraction of the capillaries; and Massolaz, in an epidemic of chronic diarrhœa among the French troops serving in the Orient, found that the suggestion was well timed. Although Barlan-Fontayral afterwards published a book (*Le Seigle ergoté et de l'Application de l'Ergotine à la Cure de la Dysentérie et de la Diarrhée chroniques*, Montpellier, 1858) upon the subject, yet it attracted little or no attention. In 1871, Prof. A. Luton, of Rheims (*Gazette Hebdomadaire*, October, 1871), stated, as something new, that he had used ergot with remarkable success in a violent and protracted epidemic of dysentery; and Dr. A. Palmberg (*Schmidt's Jahrbücher*, Dec. 1871) reports cases of chronic diarrhœa cured by the use of ergotin after the failure of other

remedies. Of course, further clinical experience is wanting before any definite conclusion can be arrived at as to the value of ergot in these diseases; but its use certainly seems to be a rational one.

Another employment of ergot for the purpose of restraining excessive secretion is in *galactorrhœa*, in which affection it has been used with success by Dr. Le Gendre (*Bulletin Thérapeutique*, t. lxxvii. p. 282), who was led to employ it by an observation of Drs. Poyet and Commarmond (*Annales de la Société de Médecine de Saint-Etienne et de la Loire*, April, May, June, 1863) that wet-nurses fed upon ergotized bread lost their milk.

The use of ergot in diseases of the nervous system has recently attracted a good deal of attention. It would appear to be especially indicated in congestion of the nerve-centres, and in *congestion* of the *cord* has been especially commended by Brown-Séquard, Hammond, Beard, and others. I have employed it with most excellent results in several cases, and there can be no doubt as to its great value when given in sufficiently large doses. In other diseases of the nervous system its use is not so clearly established. Dr. Hampel (*London Practitioner*, vol. i. p. 263) recommends it in *whooping-cough*.

Mr. Woakes believes the pain of *neuralgia* (*Ibid.*, p. 257) to be due to congestion and serous effusion into the nerve-sheaths, and has used ergot with very good results. Dr. Daniel H. Kitchen (*American Journal of Insanity*, July, 1873) has employed it in *headaches* of the most diverse and even opposite origins, and has always found it to do good.

In 1872 (*Berliner Klinische Wochenschrift*, June 17), Prof. Hildebrandt announced that in nine cases of *fibroid tumors* of the uterus he had used with the utmost advantage hypodermic injections of ergotin; and this practice has been imitated by Dr. W. V. Keating and Dr. John Ashburt (*American Journal of the Medical Sciences*, July, 1873) and others.* I have known of several unreported cases in the practice of my colleagues in which the symptoms produced by uterine fibroids have been greatly benefited by ergot. It may be that the drug simply lessens the uterine congestion, and does good precisely as it does in *chronic subacute metritis* and in *subinvolution* and *hypertrophy* of the *uterus* (Meadows, *The Practitioner*, vol. i. p. 166); or it may be that it strangles the growth by causing uterine contractions. If the latter be the case, good, as is suggested by Prof. Goodell (*Proceedings of the Medical Society of Pennsylvania for 1873*), is to be expected from the remedy only in mural and submucoid tumors.

An objection to the method of Hildebrandt is the great pain and local inflammation which often result; and Goodell proposes, as a substitute, the use of enemata or suppositories containing the drug.

Very recently, led by a fancied resemblance between the physiological action of ergot and that of quinine, Dr. Duboué has proposed the fungus as

* See *American Practitioner*, May, 1873; the *Clinic*, April, 1873; the *Lancet*, May, 1873.

a succedaneum to the alkaloid in *malarial fevers*, and in support of his views has published a book entitled *Recherches sur les Propriétés thérapeutiques du Seigle ergoté*, Paris, 1873. He brings forward some evidence of the value of the remedy; but probably the latter will share the fate of the older substitutes for the South American specific.

TOXICOLOGY.—Enough has already been said in regard to the acute poisoning by ergot, except it be to state that, when abortion is threatened from its ingestion, in the maintenance of perfect quiet and in the free exhibition of opium are to be found all the measures of relief at our command.

Since the days of Galen, there have swept over larger or smaller districts of Europe epidemics of diseases which have been attributed to ergot. In many parts of Europe, rye bread forms the great staple article of food of the lower classes. It always contains a small quantity of ergot, but not enough to have any deleterious effect upon the health.

When the summer is wet and cold, the rye becomes very extensively ergotized, so that the fungus constitutes a large proportion of the materials entering into the bread. It is under these circumstances that there occur those epidemics of *ergotism* or chronic ergotic poisoning which have been recorded from time to time since the days of Galen and of Cæsar. It is not always the rye which causes these frightful losses of life, as Heusinger (*Journal für Pharmakodynamik*, Bd. i. p. 405) has traced one epidemic to diseased oats.

Before going farther, it seems proper to state that no less an authority than Trousseau and Pidoux asserts that these epidemics are not dependent upon any specific action of ergot, but are either epidemics of blood-diseases or simply the results of improper and insufficient food,—merely the outcomes of poverty, wretchedness, and famine. It seems to me indisputable that some of the various epidemics which have been recorded were of this character, but certainly it is no less indisputable that others were not. Moreover, quite numerous scattered cases are on record in which a few persons or a family have been affected with ergotism unmistakably traceable to the use of bread largely composed of the fungus.

The scope of the present work is such as to forbid my entering into an elaborate discussion of the epidemics of ergotism, especially as the subject has no practical bearing so far as the American profession is concerned, since the absence of deep poverty is so complete in our country that no one would feed on largely ergotized bread, and, in fact, no case of ergotism has as yet been recorded as occurring in the United States.*

There are two varieties of ergotism, the spasmodic and the gangrenous.

Gangrenous ergotism has been especially observed in France, and is believed to be the same as the *Ignis Sacer* or the *Ignis Sancti Antonii* of the Middle

* Any one especially interested in the subject will find the literature very well represented in the references in Prof. Stillé's work on Therapeutics, to which may be added Duboué's book, quoted on this page. For the reader of German, a very excellent résumé exists in Husemann's *Handbuch der Toxicologie*.

Agres,—an affection which in 922 killed forty thousand persons in South-western France, and in 1128–29 fourteen thousand in Paris alone. It generally commences with itching and formications in the feet, severe pain in the back, contractions in the muscles, nausea, giddiness, apathy, with abortion in pregnant women, in suckling women drying of the milk, and in maidens amenorrhœa. After some time, deep, heavy, aching pains in the limbs, an intense feeling of coldness with real coldness of the surface, deep apathy, and a sense of utter weariness, develop themselves. Then a dark-red spot appears on the nose or on one of the extremities; all sensation is lost in the affected part; the skin, perhaps over a large surface, assumes a livid red hue, and in the foci of local changes bullæ filled with serum appear. The adynamic symptoms, in severe cases, deepen as the gangrene spreads, until finally death puts an end to the scene.

Very generally the appetite and digestion are preserved to the last, and not rarely there is an almost ferocious hunger. The gangrene is generally dry, the parts withering and mummifying; but sometimes it is moist, and pyæmic symptoms may even be developed. Of course a very large number of cases do not terminate in death; but the part immediately affected is generally lost. In these cases the toes most generally are the portion destroyed, but it may be any one or all of the extremities; and the nose, lips, ears, and even the buttocks, sometimes bear the brunt of the disorder.

Spasmodic ergotism may in the lightest cases be manifested only by itchings, formications, numbness, or complete anæsthesia of the fingers and toes or of the buttocks, and by gastro-intestinal irritation, as shown by colic, vomiting, diarrhœa, or constipation, and withal a ravenous hunger. In more severe cases these manifestations are intensified, and spasmodic symptoms appear, violent and painful tonic contractions affecting especially the flexors of the extremities, interrupted at times by intervals of quiet, but gradually growing into severe general tetanic paroxysms, with opisthotonos and emprosthotonos. In the intervals there are very generally muscular tremblings, and as the case progresses there are developed cerebral manifestations, such as disturbances of vision, photophobia, chromopsis, hemiopia, and periodic amblyopia and amaurosis, giddiness, cataleptic and epileptic paroxysms with or without loss of consciousness, delirium, and idiocy. Gastro-intestinal symptoms are always very marked, but with them is a characteristic ravenous hunger and a longing for sour food and drink. The skin is earthy or yellowish in its tint, often spotted with boils or pustules or semi-gangrenous vesicles.

Death is apparently caused by exhaustion; and in those that recover, various local paralyses, habitual spasms, amaurosis, mental aberrations, or even idiocy, often remain through life.

In a few cases the symptoms are still more violent, and the spinal and cerebral disturbances soon lead to death.

In some epidemics of ergotism the cases have been a mixture of the spasmodic and gangrenous form of the disease.

ADMINISTRATION.—Ergot may be given in powder, but at present this method is very rarely used: the dose is half a drachm to two drachms. The U. S. Pharmacopœia recognizes a *Wine* (*Vinum Ergotæ*, $\S$ i to f $\S$ vii) and a *Fluid Extract* (*Extractum Ergotæ Fluidum*). The first of these is a good preparation for internal use, and may be administered, as an ecboïc, in doses of half an ounce to two ounces. The fluid extract is prepared by exhausting ergot with a menstruum composed of alcohol, water, and glycerine, adding acetic acid, and evaporating. If carefully made, it is an efficient preparation, and closely represents the drug, drachm for drachm. The usual ecboïc dose is one to two drachms. In nervous diseases much larger doses are required: thus, I usually commence in congestion of the spinal cord with half an ounce, and increase it to an ounce three times a day. The fluid extract, containing so much acid and not being thoroughly soluble in water, is not so good for hypodermic use as the so-called *Ergotin*, a non-official preparation, made by exhausting the ergot with water, evaporating to the consistency of a syrup, precipitating the albumen, gum, etc., with an excess of alcohol, decanting the clear liquid, and evaporating to the consistency of a soft extract. This ergotin is believed to be about ten or eleven times as strong as ergot, five or six grains of it representing about a drachm of the drug. The following formula of Dr. Keating probably offers as good a preparation of ergot for hypodermic use as can be had: R Ergotinæ, gr. xlv; glycerinæ, aquæ destillatæ, āā ℥v. Twenty to thirty drops of this may be given at once. It is stated that the danger of severe local trouble is much lessened by plunging the nozzle of the syringe deeply into the muscular tissues.

CLASS VIII.—SIALAGOGUES.

SIALAGOGUES are medicines which increase the flow of saliva and of the buccal mucus. Various substances, such as mercury, when taken internally, affect the mouth and its tributary glands in such a way as to produce salivation: these substances are, however, never employed in medicine for this purpose, so that practically sialagogues are local remedies acting by the induction of a local impression on the mouth. The influence which they exert is a stimulant one, and most of them are more or less irritant. They are used to effect two distinct purposes. Some of them dissolved in the saliva pass over the mucous membranes not only of the mouth but also of the fauces and of the epiglottis, and exert upon them a direct stimulant influence. Other sialagogues, by exciting a very great flow of saliva, seem to lessen the congestion of the part.

In reference to the first of these modes of action, sialagogues are employed in relaxed conditions of the mucous membrane of the fauces, and even of the larynx. Chief among the substances so used is cubeb, which when slowly chewed in the mouth exerts a very decided local influence, and is useful in relaxation of the fauces, of the uvula, and even of the upper portions of the larynx. Either in form of the berries or made into lozenges, cubeb is much used by public speakers; and in the *hoarseness* from relaxation following over-use or slight inflammation, it is often very efficient.

Through their depletory influence, sialagogues are sometimes useful in allaying *rheumatic toothache*, or other rheumatic irritations about the jaws. The only remedy of the class employed for the purpose and requiring a separate notice here is pellitory.

PYRETHRUM, or *Pellitory*, is the product of the *Anacyclus Pyrethrum*, a small herbaceous perennial, growing in the neighborhood of the Mediterranean. It is a small root, about the size of the little finger, wrinkled longitudinally, light brown externally, with bright, shining spots on the surface, hard, brittle, with a resinoid radiated fracture. It is inodorous, and when chewed is at first almost tasteless, but soon becomes acidulous, saline, and acrid, and produces a very persistent burning, tingling sensation, which is accompanied by a profuse flow of saliva. Half a drachm to one drachm of it may be chewed at a time in painful *rheumatic affections* of the face, in *toothache*, in *relaxation of the uvula*, and in similar disorders.

CLASS IX.—ERRHINES.

THESE are substances employed to act upon the mucous membrane of the nose. Strictly speaking, the term should be applied only to those drugs which are used to excite secretion in the nasal mucous membrane. Such remedies are, however, so rarely used as to be by themselves scarcely worthy of notice. The employment of irritating vapors to arouse the nerve-centres by stimulating the nerves distributed in the nasal mucous membrane, is a very old and a very popular custom. *Smelling-salts*, or preparations of hartshorn, so much used by ladies as a slight stimulant, and by others in reviving those who are suffering from or threatened with fainting, act in this manner. The ammonia held close to the nostrils brings about the reaction, not by any direct stimulating action on the circulation, but by irritating the nasal mucous membrane, as is proven by the rapidity of its influence and by the exceedingly minute quantity which will sometimes act efficiently. In the use of hartshorn, especially with young children, it is necessary to exercise care, lest injury should be done to the delicate mucous membrane. The only errhine used for the purpose of influencing affections of the nasal passages themselves which is worthy of mention here is *cubebæ*. This freely snuffed up in powder is very useful in *acute coryza*, after the first stage of congestion and dryness has passed away.

CLASS X.—EPISPASTICS.

Counter-irritation.—Almost from time immemorial, physicians have believed that morbid processes in deep-seated or superficial organs could be modified by irritations artificially induced in distant parts. To the drugs used for producing these remedial irritations the name of revulsives or counter-irritants has been given, the process being called revulsion, or counter-irritation.

Latterly, the value of these remedies in disease has been questioned, chiefly because not only were the accepted theories of their action deemed untrue, but any explanation of how they do what is claimed for them was asserted to be, in the present state of our knowledge, inconceivable.

Evidently, in studying the matter, the inquiry should be divided into two parts, and fact should be separated from theory; the effort being made to ascertain, first, whether experience does or does not demonstrate that it is possible by an irritation to affect a distant part which has no apparent connection with the seat of the new irritation; second, whether the facts taught by experience are in truth irreconcilable with reason.

In regard to the first part of this inquiry, it seems to me indisputable that experience does teach, in the most unequivocal manner, that an organ may be affected through a distant part. There are physiological proofs of this, which it is only necessary to allude to: such is the relation of the uterus and the mammary glands. The proofs which may be drawn from disease are, however, much more numerous and striking. Thus, it is well known that in mumps there may be relief of an existing irritation of the salivary gland by a new irritation of the testes; in gout, the swelling of the toe will relieve the disordered digestion, etc. If it be affirmed that these phenomena, happening during the existence of a blood-disease, are *sui generis*, the objection cannot be made to the paraplegia produced by the irritation of a calculus in the kidney, or to the headache due to the irritation of the gastric mucous membrane by acid, or to the shoulder-pain of diseased liver, or to the amaurosis caused by the irritation of a decayed tooth. A well-known experiment of Brown-Séquard's illustrates the point so well that it may be quoted. In it he found that if one sciatic nerve of a guinea-pig be cut, epileptic attacks may be produced by gently rubbing the back of the ear upon the same side.

A very curious instance of an external irritation affecting a deep-seated part is the duodenal ulcer produced by burns, especially of the abdomen.

The pathological evidence of the truth of the present proposition is simply overwhelming, and facts might be brought forward almost indefinitely to show that irritations are capable of affecting the functions and nutrition of distant parts. This being true, surely it is in the highest degree reasonable to suppose that artificial irritations can in a greater or less measure be controlled so as to affect the distant organ for good and not for evil.

Clinical experience has certainly demonstrated that this can be done. The value of any individual counter-irritant in this or that disease is not the present question; but certainly no physician who has had any practice can have failed to see instances of relief from the use of counter-irritants. A case of obscure brain-trouble recalls itself at present writing, in which stupor and a clear intellect alternated at will, according as the drastic cathartic was given or withheld. The relief of abdominal pain, or "stomach-ache," by a mustard plaster, is a daily nursery experience.

From what has been already stated, it may be laid down as proven beyond cavil—first, that we have power to influence internal morbid processes by creating external irritations; second, that the fact of counter-irritation exists, whether we can or cannot explain its rationale.

Physiological knowledge is not yet sufficiently extensive to enable us to perfect a theory of counter-irritation. The action of these remedies is complex, but I think can be explained at least in part. There is only a certain amount of blood in the body. If it be accumulated in one place, it cannot be in another. Thus, the difficulty of studying after a hearty dinner probably depends, as do the cold feet so common in feeble persons under such circumstances, upon the accumulation of blood, and probably also of nervous energy, in the digestive organs. Now, by artificial interference, by determined study, by violent exercise, we can often draw the blood away from the alimentary apparatus into the cerebrum, or into the motor system, and produce indigestion. Clinical experience proves that we can also reverse this process. The brain is excited, the blood is concentrated in it, congestion exists, inflammation is threatened; a drastic cathartic is given, the blood is drawn into the intestinal canal, and by revulsion the brain is relieved. Certainly this is not mysterious, not inexplicable.

All forms of counter-irritation cannot, however, be explained on the above principle. It is a probable, but not a positive, teaching of modern physiology that there are nerves which preside over nutrition,—the so-called trophic nerves. If this be so, it is to be expected, *a priori*, that peripheral irritations will cause reflex alterations of nutrition, precisely as they cause reflex disturbances of the motor functions. Further, whether these trophic nerves do or do not exist, there are vaso-motor nerves, and the duodenal ulcer of burns is a positive proof that, either through the trophic or through the vaso-motor nerves, external irritations do produce internal reflex alterations of nutrition.

The sympathetic ophthalmia caused by a morbid eye in its healthy fellow, or induced by a diseased tooth, is another instance of this reflex alteration of nutrition. As this is true, it seems to me absurd to state that it is impossible to conceive how an external counter-irritant can affect the nutrition of a deep-seated organ.

It is evident that in all the cases which have been mentioned of external irritation causing disease in a distant organ, there is no direct communication between the part irritated and the organ which is secondarily affected. And clinical experience confirms the evident deduction from this,—*i.e.*, that it is impossible to determine, except by experiment, where the counter-irritant should be placed to affect most powerfully any given organ.

It has, however, been clinically demonstrated that the general law for deep-seated parts is that the revulsant should be put directly over the part. When a superficial action is desired, other directions are needed.

We are indebted to Dr. Anstie for pointing out what appears to be another law,—namely, that when a superficial part supplied by the anterior branches of a spinal nerve is to be affected, the counter-irritant should be placed over the posterior roots of the nerve. Not only can obstinate neuralgia often be relieved by this reflex action, but also the inflammatory changes so often coincident with intercostal neuralgia. The law seems also to apply to cervical nerves, since the proper position for the blister in facial trigeminal neuralgia is back of the ear or on the nape of the neck.

Counter-irritants may be conveniently arranged under two heads: 1st, those which produce a decided structural alteration of the skin, including *epispastics*; 2d, those which do not provoke decided alterations of dermal structure, the *rubefacients*. The indications for the use of these substances can best be considered under their respective headings.

As is well known, any sthenic inflammation, if of sufficient extent and intensity, may excite the general system even to the point of high fever. In this respect inflammation of the skin does not differ from that of other organs. Hence dermal irritants have a direct tendency to arouse or excite the system, and may be used as general stimulants. It will be seen at once that it is the nervous and arterial systems which alone feel their influence. Hence the irritants should not be relied upon in cases of exhaustion, for the only possible source of absolute increase of power to the system is in food; and in exhaustion those stimulants should be employed which increase the power of assimilating food. For this reason, external irritants are useful as stimulants in conditions of depression rather than of exhaustion. Such conditions of depression exist in *acute collapse* from any cause, in "*shock*" following injuries, in the first stage of *pernicious malarial fever*, in *snake-bite*, and in other cases when the powers of the system are seemingly overwhelmed by some depressing agency. The rubefacients are preferable to blisters for this purpose, because their local after-effects are comparatively so trifling that they can without injury be applied to a very large extent of the surface.

EPISPASTICS, *vesicatories*, or, more colloquially, *blisters*, are substances which are used by the physician to produce that peculiar inflammation of the cuticle and outpouring of serum known as a blister.

The immediate effect of a blister is more severe and more permanent than that of a rubefacient. Blisters are especially useful in inflammations of serous membranes, such as *pleuritis* and *peritonitis*; are very strongly recommended by some practitioners in parenchymatous inflammations, such as *pneumonia*; and are often of service in *neuralgia*, and in other forms of nervous irritation, such as the *maniacal delirium* of fevers, when dependent upon the irritant action of a blood-poison, and not upon exhaustion.

The amount of serum which is poured out from a blister is sometimes quite large, and vesicants have even been used to relieve *dropsy*. In general dropsy, their use is simply unjustifiable; but in *local dropsies*, as, for example, serous effusion into the pleural sac or into the pericardium, dependent upon local disease, they often do good not only by affecting favorably the disease-process, but also by hastening the removal of the effusion.

In some chronic affections, long-continued severe counter-irritation is required: in such cases a blister may be "kept open" by the use of stimulating ointments, such as the *unguentum mezerei*.

In *chronic inflammation* of the *joints*, repeated blistering is very often of service. When the inflammatory action is rheumatic, in my experience better results are obtained by repeated blistering than by keeping a blister sore by means of irritants.

In *neuritis*, whether rheumatic or otherwise, blisters are often of service: they should be applied as a long narrow strip along the course of the nerve.

In obstinate local *neuralgia*, very mild blistering over the seat of pain, or in accordance with Anstie's law, is sometimes advantageous.

The *contra-indications* to the use of blisters are high arterial and febrile excitement and a decided want of vital power. In the former case, the irritating influence which they exert upon the general system may increase the constitutional disturbance to such an extent as to do far more injury than any local benefit derived from them can do good. When the vitality is very weak, blisters may give rise to sloughing ulcers, which, refusing to heal, may waste very seriously the already exhausted system. Hence, in all acute diseases of such type that the nutritive forces are exceedingly depressed, blisters must be avoided, or must only be used with great caution. For the same reason, great care must be exercised in their employment in the very young or the very aged. Very rarely indeed is a blister called for in the case of a young infant, and, when employed, it should be allowed to remain in contact with the skin only just long enough to produce slight redness, and the complete vesication should be obtained by the use of the poultice.

There are various substances which are capable of producing vesication, but

the only one in ordinary use is cantharides. In cases of emergency, a blister, it is said, may be raised in a very few minutes by the use of the *stronger water of ammonia*, a little of which is to be kept in contact with the skin by means of an inverted watch-glass. It is necessary to watch the process closely, and to remove the irritant so soon as vesication has occurred, as the ammonia is very capable of causing sloughing.

CANTHARIS—CANTHARIDES.

The dried bodies of the *Cantharis vesicatoria*, a beetle inhabiting Southern Europe, and coming into commerce in Spain, Italy, Sicily, and the southern provinces of Russia. *Spanish flies* are from half an inch to nearly an inch in length, and two to three lines in breadth, and have a large heart-shaped head, and brilliant metallic-green elytra or wing-cases. Their odor during life is very strong and fetid, but is almost entirely lost in drying; their taste is urinous, very burning, and acrid. They are taken in May and June, when they swarm on the trees which they affect, by beating the branches early in the morning, when the insects are torpid from the cold, catching them upon linen sheets, and plunging them into hot vinegar and water, or exposing them to the fumes of boiling vinegar. In some places they are gathered by smoking the trees with the fumes of burning brimstone. When ground, Spanish flies afford a grayish-brown powder, full of minute greenish spangles, the remains of the feet, head, and wing-cases. The active principle of cantharides is *Cantharidin*, which occurs in white crystalline scales, is inodorous, tasteless, insoluble in water, nearly so in cold alcohol; soluble in ether, benzole, the oils, and very freely so in chloroform. Notwithstanding the insolubility of their active principle, Spanish flies yield their virtues to alcohol and to water, because, as is generally believed, a yellow coloring-matter is so united with the cantharidin as to render it soluble.

PHYSIOLOGICAL ACTION.—When a minute therapeutic dose of cantharides is taken, no perceptible immediate result is produced, and after a somewhat larger quantity the only symptom is usually some burning and pain in urination. Doses more than just sufficient to induce this should not be employed in medicine, as the symptoms produced by large amounts of the drug are exceedingly severe and distressing. Cantharides is very irritating, and, when applied to the skin, causes at first redness, with burning, then free vesication and severe pain, and, if the contact be longer maintained, deep inflammation and sloughing. Upon the mucous membranes it produces a no less intense effect; and consequently gastro-intestinal inflammation forms a prominent symptom of poisoning by it. Further, the active principle or principles are undoubtedly absorbed and are eliminated by the kidneys, coming in contact with almost the whole genito-urinary mucous membrane: hence intense irritation and inflammation of these organs always result from the ingestion of an overdose of *Spanish flies*.

Very soon after a toxic dose of cantharides has been taken, the sufferer is

seized with burning in the pharynx and œsophagus, and a sense of stricture in the throat. The pain soon spreads to the stomach, and vomiting comes on. The symptoms rapidly increase in severity; the abdominal pain becomes very severe, and, in the majority of cases, purging takes place. The matters rejected by the stomach are first mucous (with, if the drug have been taken in powder, little greenish specks through them), then bilious, and finally bloody. The stools are mucous, then fibrinous, bloody, becoming often very scanty, but excessively numerous, and in their passage accompanied by great tenesmus. Probably in most cases very early in the attack, severe salivation is developed; and it is very frequently accompanied by great swelling of the salivary glands.

Sometimes death occurs in a very short time, from collapse produced by the intense gastro-intestinal inflammation; but more generally it is postponed for some hours, and a new train of symptoms arises. Aching pains in the back, and very frequent micturition, indicate the commencing urino-genital irritation. These symptoms increase in intensity until there is a constant irresistible desire to urinate, with violent tenesmus of the bladder, and yet an inability to pass more than a few drops of urine, which is albuminous, and not rarely bloody. In some cases there is a violent erotic excitement, an unquenchable lust, accompanied in man by numerous seminal emissions; violent priapism, swelling and heat of the organs, and even severe inflammation of the parts, indicate the intensity of the local action of the poison; sometimes gangrene ultimately occurs.

Neither amatory desire nor true priapism is, however, a constant symptom in cantharidal poisoning (cases, *Journal de Pharmacie et de Chimie*, June, 1871): indeed, the former is probably absent in the majority of cases. Consciousness and general power are often long preserved amid intensely severe local symptoms and agony, but, if the dose have been large enough, sooner or later collapse comes on with the usual symptoms, and the prostration deepens into complete powerlessness, stupor, coma, and finally death. In some cases violent hydrophobic delirium and severe tetanic convulsions are said to have occurred (Tardieu).

Paraplegia has been noticed in several cases by Dr. Pallé (*Journal de Pharmacie et de Chimie*, June, 1871): it was probably reflex in its origin, and due to the intense irritation of the urino-genital organs.

In animals, cantharides produces very much the same symptoms as in man. In dogs, according to the experiments of Orfila and of Beaupoil, the symptoms of gastro-intestinal inflammation are more prominent than those of irritation of the genito-urinary tract. It has been asserted that the lack of erotic excitement in these cases shows that the medicine acts differently upon man and animals. As already stated, however, erotic delirium is very generally absent in fatal poisoning by man, whilst Schroff states that ten drops of the tincture of cantharides will frequently produce great sexual excitement in man, and the whole drift of the evidence is that libidinous desires are much

more apt to be caused by amounts of Spanish flies but slightly toxic, than by fatal doses. Indeed, the irritation caused by the latter would seem to be too intense, the general perturbation too great, for erotism to be induced. There appears to be the same difference in the effects of different doses of the drug upon animals. Fatal doses very generally do not excite sexual desire; but Schubarth (quoted by Stillé) found that small doses do cause evident salaciousness and irritation of the genital organs, whilst, according to Husemann (*Handbuch der Toxicologie*, 1862, p. 264), the peasants of Northern Germany habitually give cantharides to cows when backward in coming into heat at the proper season.

THERAPEUTICS.—Cantharides is employed internally only for the purpose of influencing the genito-urinary organs; and sufficient has already been said in regard to this use under the headings of Diuretics and Emmenagogues.

The external use of cantharides is simply as a vesicant; and the employment of blisters has been sufficiently considered in the general discussion of the class. Two points, however, seem worthy of notice here: first, that this drug affords the only practical means of blistering at our command; secondly, that when it is freely employed there is always some danger of the absorption of a sufficient amount of the active principle for strangury to be induced. In susceptible persons, therefore, care has to be exercised in the use of epispastics; and whenever active irritation of the kidneys exists, cantharidal blisters should not on any account be applied.

TOXICOLOGY.—Sufficient has already been said about the symptoms produced by cantharides. The minimum fatal dose is not certainly determined, and probably varies very much. According to Stillé, twenty-four grains of the powder, taken in two doses, have caused fatal abortion, and an ounce of the tincture has destroyed life after the lapse of a fortnight. After death, intense injection, swelling, patches of exudation, loss of epithelium, and other results of inflammation are found along the whole tract of the alimentary canal; intense hyperæmia of the kidneys, with contraction and injection of the bladder, also usually exists.

There is no known antidote to cantharides, and the treatment of the poisoning must be conducted upon general principles. The stomach, if not already thoroughly emptied, should be evacuated at once by a stimulating emetic if the stomach-pump be not at hand. Large quantities of mucilaginous or albuminous drinks should be taken; and all oily substances should be avoided, as favoring the solution, and consequently the absorption, of the poison. Opium should be freely exhibited, especially by the rectum, to allay pain and relieve the strangury. For the latter purpose warm sitz-baths or general baths should be given. In some cases leeches to the epigastrium are advisable. When the suffering is very intense, the cautious use of anæsthetics seems to me not only justifiable, but imperative. In the stage of prostration, the measures to be adopted are those commonly practiced in collapse from poison.

ADMINISTRATION.—The preparation of cantharides most commonly used for the production of a blister is the *Ceratum Cantharidis*, which is best spread upon sticking-plaster in such a way as to leave a margin about an inch in width, which shall adhere to the skin and hold the plaster in its place. In the last edition of the U. S. Pharmacopœia the *Cantharides Paper* (*Charta Cantharidis*) was introduced: it is thought to be more elegant than the plasters made of the cerate, and is said to be efficient. In order for a blister to “draw” thoroughly, it usually has to be left on some eight hours; but in most cases the same result can be achieved with less suffering by allowing the blister to remain only six hours, or until decided redness and slight vesication have been induced, and then applying a flaxseed poultice. In certain localities vesication requires a much longer application than that just spoken of: thus, upon the shaved scalp a blister will rarely act efficiently in less than twelve hours, and often not in that time. In maniacs, in the delirious sick, in children, and in other unruly patients, it is often necessary to put on a blister in such a way that the sick person has no control over it. For this purpose the officinal *Cantharidal Collodion* (*Collodium cum Cantharide*) may be used. It is ordinary collodion impregnated with cantharides, and on evaporation leaves an adhesive blistering film: two or three coats of it should be applied by means of a camel’s-hair brush.

When there is any especial danger to be feared from absorption of the active principle, the use of the poultice, after a brief application of the blister as described above, should always be practiced.

CLASS XI.—RUBEFACIENTS.

RUBEFACIENTS are those remedies which are employed for the purpose of producing, not any permanent inflammation of the skin, but a general intense irritation, redness, and congestion, which shall exert a temporary influence, whose power is the result of the large surface affected, and not of any permanent impression upon the nutritive acts of that surface. Most, if not all, rubefacients are capable of causing disorganizing inflammation if allowed to remain for too long a time in contact with the skin.

The superiority of rubefacients over blisters when it is desired to arouse or stimulate the system has already been pointed out (see p. 476). It remains to speak of the use of rubefacients in local diseases. They are especially useful in sudden cases of severe pain due, it may be, to acute congestion of a part, or to some internal irritation like that of gout. Thus, in the ordinary intestinal pain caused by irritant articles of food, or more commonly by a rheumatic, gouty, or other irritation following exposure to cold or wet, rubefacients are most useful. In this as in all other cases of what may be termed temporary functional derangement, when a counter-irritant is desired, rubefacients are superior to blisters, because their effects are not nearly so lasting, and also because, for the time being, they seem to impress more powerfully the nervous system, breaking up, as it were, the concentration of nervous energy, or calling off the irritation, or impressing the nervous system in some way which in our present ignorance it is difficult to find terms to express. A correct idea of the difference in the use of the two classes of counter-irritants can perhaps be conveyed by saying that when profound local alterations of nutrition are to be dealt with, blisters are to be used; when functional disturbance is to be met, rubefacients are to be employed. Yet this rule cannot be applied with strictness.

SINAPIS ALBA—WHITE MUSTARD.

SINAPIS NIGRA—BLACK MUSTARD.

The seeds of the *Sinapis alba* and *Sinapis nigra* respectively,—European crucifers, cultivated in the temperate regions of the world.

These seeds are minute, globular bodies, yellowish within: they are to be distinguished one from the other by the smaller size, external brown color,

and more fiery taste of the black mustard, and the light-yellowish exterior of the white mustard.

Black Mustard contains *Myronic Acid* in combination with potash, and also a peculiar albuminous principle, *Emulsin*. When to these substances water is added, a reaction occurs, resulting in the production of a volatile oil out of the myronic acid. *Volatile oil of mustard* is a colorless or yellowish fluid, of an intensely pungent, or even corrosive, odor and taste. A momentary contact with it suffices to redden and blister the skin, and mucous membranes are said to be rapidly destroyed by its vapors.

White Mustard does not yield on distillation with water a volatile oil, but contains an acrid fixed principle. The chemistry of white mustard seeds appears not to have been certainly determined. In 1825, Henry and Garot discovered a substance in them,—*Sulpho-sinapisin*,—which, according to Husemann, has been variously designated as *Sinapin*, *Sinapisin*, *Sinapinic Acid*, etc., but has been demonstrated by Babo and Hirschbrunn to be an alkaloid, which also exists in the seeds of the black mustard. Robiquet and Boutron believe that the acrid fixed principle of white mustard is formed by a reaction between this and water in the presence of the emulsin.

PHYSIOLOGICAL ACTION.—The oil of mustard is a most intense irritant, and if taken internally would act as a very powerful irritant poison. The ground mustard, of course, shares its properties.

THERAPEUTICS.—Mustard affords a most excellent material for the practice of mild revulsion. One great advantage it possesses is the ease with which it can be controlled,—all grades, from the mildest impression up to severe blistering, being at the will of the practitioner. It should be remembered, however, that the blister produced by it discharges but little, and is exceedingly sore and painful as well as very slow and difficult of healing: so that, as an epispastic, mustard is in every way inferior to cantharides, and should not be employed.

The black mustard is much stronger than the white, and must usually be diluted at least one-half (by the addition of flour or of flaxseed meal). The white variety may sometimes be employed pure, but generally it also may be reduced in strength.

In many cases it is desirable to maintain for hours a mild, equable counter-irritant impression; and this may be done by adding from one to three teaspoonfuls of mustard, more or less, to a poultice of flaxseed. A mustard poultice half-and-half black mustard, three parts to one white mustard, and flour, may generally be left on from twenty minutes to half an hour without danger of blistering. Weaker preparations may be used longer. A mustard plaster may be prepared like an ordinary poultice; but a very convenient method is to take a newspaper folded to a little larger than the desired size, and tear open the front piece so that it can be folded back like a flap, leaving one edge attached; next, to spread upon the thick portion the mustard, leaving the edges free, and then to close the flap upon it and fold the edges back to

the desired shape: when done with, this plaster can be thrown away, and no rags are lost. The mustard draws well through the single layer of newspaper covering it, but is, I think, less apt to leave troublesome after-soreness than when employed in the usual manner. In the last edition of the U. S. Pharmacopœia a new preparation of mustard was introduced, under the name of *Charta Sinapis*, or *Mustard Paper*. It consists of black mustard mixed with solution of gutta-percha and spread upon stiff paper four inches square. These mustard papers may be useful for travelers, but in my experience do not act so well as the home-made mustard plaster. Their size is not so easily regulated, and their action is often too violent.

CAPSICUM and the *stronger spices* afford excellent materials for rubefaction. The Cayenne pepper is probably as strong as mustard, but is much less pleasant to handle, on account of the readiness with which it is diffused, and is much less frequently employed. *Spice plasters* are useful when it is desired to make a steady, continuous mild impression, as in certain abdominal complaints.

They may be made by the apothecary by means of the following recipe: Take of powdered ginger, ℥ii; powdered cloves and cinnamon, each, ℥i; Cayenne pepper, ℥ii; tincture of ginger, f℥ss; honey, q. s.; mix the powders, add the tincture, and sufficient honey to make of proper consistence for a stiff cataplasm.

The domestic spice-plasters are much more elegant and cleanly than those made on the above plan. They are to be prepared as follows. Take equal parts of ground ginger, cloves, cinnamon, and allspice, and one-fourth part of Cayenne pepper, and thoroughly mix them; then put the resulting dry powder into a previously-prepared flannel bag of the desired size, distribute the powder equably through the latter, and quilt it in,—i.e., run lines of stitching across the bag, so as to confine the powder in little compartments. When used with common whisky or with alcohol, a plan which has seemed to me still more pleasant is to put two ounces of *unground* ginger, an ounce of unground cloves, cinnamon, and chillies, or African peppers, in a pint bottle, and pour the whisky upon them. After this has stood awhile, the liquor is to be put upon a piece of flannel of the proper size, and the latter is to be laid upon the part and covered with a larger piece of oiled silk, or else a piece of spongio-piline may be employed. If the strength of the preparation is too great, it can readily be reduced by dilution; if it is too little, it can as readily be increased by adding more of the spices, especially of the peppers. In many cases, when the tenderness is very great, the weight of the spice-plaster is objected to. Under these circumstances the substitute here proposed is especially valuable.

The Oil of Turpentine is a very powerful rubefacient, capable, if applied to the skin for too long a time, of destroying the epidermis. It produces, when properly used, simply an intense diffused redness. The most frequent mode of application is in the form of *stupes*, which should be made by dipping a

piece of flannel, previously wrung out with warm water, into a cup of the turpentine which has been warmed by setting it in hot water, and then wringing out all excess of the turpentine, and applying. These stupes may be left on from ten to thirty minutes, according to the severity of the impression desired and the susceptibility of the patient's skin. On some persons the least contact of turpentine, or even of its vapors, produces a most painful furuncular eruption. Where this idiosyncrasy exists, of course the remedy should never be used. The official liniment (*Linimentum Terebinthinæ, Kentish Ointment*) is used as a stimulant application to burns. According to the U.S. Dispensatory, it should be applied as soon as possible after the reception of the burn, by covering the injured surface with pledgets of patent lint saturated with it, and should be allowed to remain on until the peculiar inflammation excited by the fire has subsided.

Ammonia is a most efficient rubefacient, which in its general relations has been sufficiently discussed elsewhere. When great haste is required, it may be employed as an epispastic by applying a piece of common lint, saturated with the strongest water of ammonia, and covering it with some impervious coating. Great care must be practiced lest the ammonia act as an escharotic, since a too prolonged application may produce a deep slough. To raise a blister requires from five to ten minutes. On account of its cheapness and efficiency, ammonia is very largely used in extemporaneous liniments. In prescribing, it must always be borne in mind that there are two official waters of ammonia: *Aqua Ammoniæ Fortior*, with a specific gravity of 0.900, containing twenty-six per cent. of the gas; and *Aqua Ammoniæ*, with a specific gravity of 0.960, containing about ten per cent. of the gas. The rubefacient action of ammonia is less permanent than that of turpentine. There is an official liniment (*Linimentum Ammoniæ*), containing one part of the simple water to two parts of olive oil. It is, of course, an ammoniacal soap.

PIX BURGUNDICA, or *Burgundy Pitch*, is a concrete juice obtained by wounding the *Abies excelsa*, or Norway spruce, and *Abies picea*, or European silver fir,—lofty forest-trees of Middle and Northern Europe,—melting the product of the exudation with hot water, and straining. It is hard, opaque, brittle, of a feeble terebinthinate odor and taste, and contains resin and a minute amount of volatile oil. It is a mild rubefacient, which, in the form of plaster, may be kept applied for a long time in *chronic bronchitis* and in *rheumatic affections* of the trunkal muscles. The official plaster (*Emplastrum Picis Burgundicæ*) contains one-twelfth of its weight of wax.

The *Warming Plaster* (*Emplastrum Picis cum Cantharide*) contains one part of the cerate of cantharides to twelve parts of Burgundy pitch, and is a very decided counter-irritant, whose prolonged use will sometimes blister susceptible skins.

PIX CANADENSIS, or *Canada Pitch*, is the concrete juice of the *Abies Canadensis*, or the indigenous hemlock spruce of this country, purified by melting with water and straining. It is hard and brittle in winter, soft in summer; of a yellowish-brown color, deepening to black on exposure. It contains resin and a minute quantity of volatile oil, and is used for the same purpose as the Burgundy pitch. The officinal plaster (*Emplastrum Picis Canadensis*) contains one-twelfth of its weight of wax.

CLASS XII.—ESCHAROTICS.

ESCHAROTICS are drugs which are used to destroy diseased or sound tissue. Many of them exert a purely chemical influence, whilst others seem to destroy life by directly affecting the vitality of the part, and are said to act dynamically. Those which act chemically do so in several ways: some, like bromine, probably produce an intense corrosive oxidation, whilst others, like sulphuric acid, abstract the water.

Escharotics are used for various purposes. Formerly they were employed to open abscesses; but in the very few cases in which the knife is not allowable, as in abscess of the liver, aspiration affords, without doubt, a much superior and, in *hepatic abscess*, much safer method. They are constantly employed to destroy unsound, harmful tissues and growths. Thus, they are used to remove the specific tissue of a *chemere*, or to kill a *malignant* or *semi-malignant tumor*.

Another purpose which they fulfil is the destruction of *poisoned wounds*. In these cases they may in some instances destroy the poison itself, but in other cases they simply prevent the absorption of the toxic agent by putting an end to the life-actions of the tissue containing it. It is hardly necessary to mention all the various cases in which caustics are employed to overcome the effects of poisoned wounds. *Hydrophobia* is a perfectly uncontrollable disease; but the thorough destruction of the wounded tissue at any time before the manifestation of the symptoms will probably prevent its occurrence, as it will certainly do if performed early. In *malignant pustule*, life depends upon the free early use of escharotics.

Escharotics are employed to produce ulcerations which shall be the bases of *issues*; also, by destroying the exuberant granulations or the indolent surfaces of *ulcers*, to remove at the same time diseased tissue, afford protection to the parts below by forming an impermeable surface, and exert such alterative action upon the part as shall modify for good the life-processes.

It is evident that the choice of the caustic should depend upon the object to be attained. When large tumors are to be killed, or when it is all-important completely to destroy a poisoned wound, a powerful deep-reaching escharotic must be employed; but when the surface of an ulcer is to be filmed over, a caustic which acts superficially and forms a dense albuminous coating, as does nitrate of silver, is to be chosen.

All of the more powerful of the escharotics, when taken internally in sufficient amount, act as violent corrosive poisons, producing agonizing pain in the œsophagus and hypogastrium, violent bloody vomiting, often purging of similar character, and finally collapse, deepening into death, which is sometimes preceded by convulsions. When the dose is not so large, the system may rally from the immediate effects of the poison to succumb finally to the local lesions produced, or to struggle through a protracted convalescence to health, perhaps only to die years afterwards from organic stricture, caused by the ulcerations of the œsophagus or other of the digestive tubes. The first indication in poisoning by one of these substances is to neutralize or chemically antidote the poison. Many, if not quite all, of the escharotics have some chemical antidote: with the alkalis, dilute acid, generally convenient in the form of vinegar; with the acids, alkalis, generally at hand in the shape of whitewash or of soap; with others, specific substances, which, as antidotes, should be at once exhibited. Opium should always be freely given, and the symptoms during and after the first poisoning be treated as they arise, upon general principles.

POTASSA.—*Caustic Potash* is officinally prepared by boiling liquor potassæ until ebullition ceases and the potassa melts, when it is run into cylindrical moulds. It occurs in grayish semi-translucent sticks, about three inches long and as thick as a large goose-quill, very deliquescent when exposed to the air, and extremely soluble, except impurities (lime, oxide of iron, and carbonate of potassa), in both water and alcohol.

When it is placed upon the skin it soon melts, and, as it does so, gives rise to a pain which increases until it becomes very intense, and continues until the power of the alkali is so lost that it can no longer reach through the tissue it has killed to the sound flesh below. Under the action of the escharotic the skin becomes a dirty ashen gray, and finally a slough is formed, with inflammation of the surrounding parts, and ulceration and detachment of the dead tissue in from six to ten days. The potash appears to act chiefly by abstracting the water, and, to some extent, by combining with the fatty and other portions of the tissues.

Its slough being perfectly permeable, and its power being but slowly expended by its own action, potash is one of the most thorough of the escharotics: it is, therefore, to be preferred when a very deep and decided influence is required, as after the bite of a *rabid* dog. It is somewhat uncontrollable in its action, and requires care in its use. The best method of application is as follows. Take a piece of heavy adhesive plaster, and cut a hole in it of such size that, when the piece is warmed and properly placed upon the skin, the part to be acted upon will be exposed whilst all around it will be protected. Then apply the plaster, and grease the outer surface of it, without allowing any of the oil to come in contact with the exposed central skin. Then lay the caustic potash upon the latter, and, when the action is believed to have extended deep enough, wash the part with dilute vinegar.

POTASSA CUM CALCE.—*Vienna Paste* is a grayish-white powder, composed of equal amounts of caustic potash and of caustic—*i.e.*, unslaked—lime. It is less active than the simple caustic potash, but is preferred by some on account of its being less apt to spread and diffuse itself. When applied, it is mixed with sufficient alcohol to form a paste, and then used in the manner described when speaking of the simple alkali. M. Piedagnel affirms (*Journal de Pharmacie et de Chimie*, 3e sér., t. xxxiii.) that this caustic may be rendered nearly or entirely painless in its action by mixing one part of the muriate of morphia with three parts of the powder, and then by the addition of chloroform forming a paste that may be spread upon lead-plaster and so applied. In five minutes the skin under the application becomes of a dead white color, and at the end of fifteen minutes is brown and carbonized. If the application be persisted in, the thickness of the eschar becomes finally about equal to that of the layer of the paste employed.

ACIDUM ARSENIOSUM.—As a caustic, *arsenic* is energetic and powerful, but somewhat slow, in its action, and causes intense pain, with violent inflammation of the neighboring parts. It is stated to affect more rapidly morbid than normal structures, and hence is especially used for the destruction of malignant growths. It appears to act chiefly upon the vitality of the part, acting, when sufficiently diluted, as a powerful irritant, and when in a concentrated form producing an irritation so intense that life cannot endure it. Hence, probably, the reason of its affecting more rapidly morbid growths, which, as is well known, have a lower vitality than sound tissues.

The great objection to the employment of arsenic is the possibility of its absorption in sufficient amount to cause constitutional symptoms: even death has resulted from its external use. Absorption takes place, of course, much more rapidly in a healthy than in an intensely inflamed or dead tissue. For this reason, whenever arsenic is employed it should be used freely, and under no circumstances should it be applied to a large healthy surface, such as that of a fresh wound. The rationale of these precepts is very evident. Arsenic, if employed at all as an escharotic, must be applied in quantities which would, if absorbed, endanger life, but when a very large amount is used it causes death so rapidly as to prevent absorption. In the case of a fresh wound, the poison may be taken up before it has an opportunity to produce local death.

Used in any way, arsenic is a hazardous caustic, and it ought to be employed only with the knowledge and distinct remembrance of this fact. *Cancer*, and perhaps some forms of semi-malignant ulceration, such as *lupus*, appear to be the only diseases which justify its use. *Sir Astley Cooper's Arsenious Ointment* consists of one drachm of arsenious acid, one drachm of sulphur, and an ounce of spermaceti cerate, which is to be allowed to remain in contact with the morbid growth for twenty-four hours.

The *Arsenical Paste of Frère Cosme and Rousselot*, which is officinal in

France, is composed of one part of arsenious acid, two of dragon's blood, and two of porphyryzed cinnabar, made into paste with mucilage when applied. There is no reason for believing that any of the almost innumerable substances which have been proposed as a basis for arsenious pastes possess any marked advantages: the only needful direction is to mix the caustic with from eight to ten times its bulk of inert material of such a nature as to make either an ointment or a paste, as the whim of the surgeon or the circumstances of the case may dictate, and to allow this to remain on for from eighteen to twenty-four hours.

ZINCI CHLORIDUM.—*Chloride of Zinc* is made by the action of muriatic acid upon zinc. It occurs in broken fragments of a grayish-white color, translucent and waxy in appearance, of an acrid corrosive, or, when diluted, acrid astringent, metallic taste. It is extremely deliquescent, fusible, volatilizable at a high temperature, and very soluble in both water and alcohol.

Chloride of zinc is a very powerful caustic, producing, when applied in a concentrated form, intense pain lasting for six to eight hours, and a whitish eschar, which usually separates in from six to twelve days. Its penetrating powers are said to be a little less than those of caustic potash, and its action is certainly much less rapid and diffusive. It is a superior caustic, because its action is more readily controlled than that of potash, because its absorption does not endanger life, as is the case with arsenious acid, and because it leaves a slough which is free from odor.

Canquoin's Paste is made by mixing the chloride of zinc with flour and water. The strength varies according to the purpose, the weakest paste containing only one part of the caustic in six parts: the strongest, one part in three. When used, ten or fifteen drops of water are added to the paste, which is applied in layers, successive applications being required when a large tumor is to be destroyed. Anhydrous sulphate of lime has been especially commended by Dr. A. Ure, as forming a drier paste with the escharotic and imitating its action more definitely to the site of application than any other substance. Concentrated alcoholic or watery solutions of the chloride of zinc are often used as caustics in cases of *chancres* and other small *specific ulcers*, and are reputed to be efficient. They should be applied by means of little pledgets of lint. As the action of the chloride upon the skin is slow and very painful, whenever the cuticle over the part to be destroyed is sound it should be removed by means of blisters. By some surgeons the escharotic is introduced directly into the tumor to be destroyed. Thus, Maisonneuve makes a paste of one part of the chloride with three of flour and a little water, then cuts these into pointed strips or "arrows" and dries them. He then thrusts these hardened bodies into the tumor,—if necessary first making incisions with the bistoury,—in such a way that they lie close together and form a ring around the base of the tumor. A continuous slough is thus created, which cuts off the remainder of the mass from the sound tissue

and causes its death. Sometimes Maisonneuve simply thrusts these arrows into the body of the tumor, and destroys it directly. There is an official *Solution (Liquor Zinci Chloridi)*, which is used as a disinfectant.

HYDRARGYRUM CHLORIDUM CORROSIVUM.—*Corrosive Sublimate* is an escharotic of moderate power, which shares the dangers of arsenic, since death has followed its external use. In saturated solution it is much used as a caustic in *chancres*, but is scarcely equal to the acid nitrate of mercury. In these cases it should be applied by means of a camel's-hair brush. Prof. George B. Wood recommends very highly in *onychchia maligna* the use of a powder composed of equal parts of corrosive sublimate and sulphate of zinc intimately mixed. This powder is to be sprinkled thickly over the diseased surface, and a pledget of lint thoroughly wet with laudanum is to be laid thereon. There is severe pain for half an hour to an hour; but the dressings are not to be removed until eight or ten hours have elapsed. When the slough which is thus formed separates, a healthy granulating surface is left behind.

LIQUOR HYDRARGYRI NITRATIS.—Solution of the acid nitrate of mercury is a nearly colorless, highly corrosive, acid liquid, having a specific gravity of 2.165, and made by dissolving mercury, or its red oxide, in a large excess of nitric acid. It contains free nitric acid and the binitrate of the deutoxide of mercury. I do not know of its external use ever having produced death; but it has caused salivation, and it is perfectly conceivable that its careless employment should lead to much more serious results: indeed, in a recent number of the *London Lancet* (Jan. 3, 1874, p. 40) is reported a very serious case of poisoning by the application of the pernitrates to a space not bigger than a half-crown. It is rarely used, except for the purpose of destroying *specific* or *cancerous ulcers*. It is especially useful in *chancres*, to which it should be applied with a glass rod. In obstinate *acne*, a minute drop applied by means of a glass brush to the top of an indolent tubercle is said to destroy it without producing a scar. It has been largely employed by gynecologists in *ulcerations of the cervix uteri*. Its action in all these cases is very prompt and is moderately deep; the pain is severe, but transient.

ACIDUM SULPHURICUM.—*Sulphuric Acid* is a powerful escharotic, and has been used extensively for the purpose of destroying even large growths. For this purpose the strongest acid is mixed with charcoal, so as to make a thick, manageable paste. Before the application of this, the skin should be removed by a blister. At present this escharotic is not much employed.

ACIDUM NITRICUM.—*Nitric Acid* is a powerful caustic, which is never employed to destroy large growths, but is a favorite application to *chancres*, to *syphilitic*, *phagedenic*, and other unhealthy *ulcers*, as well as to *condylo-mata* and other small *dermal growths*. It is applied by means of a glass rod or a splinter of wood. A drop or two is usually amply sufficient; and

when the action has gone far enough, the part should be washed with soapsuds, which at once arrests action and causes the severe pain to cease.

ACIDUM MURIATICUM.—*Muriatic Acid* is capable of acting as a caustic, but is less powerful than either sulphuric or nitric acid, and is rarely used.

ACIDUM CHROMICUM.—*Chromic Acid* occurs in anhydrous acicular crystals, of a deep-red color, and an acid, metallic, corrosive taste. They are very deliquescent, melting down, when exposed to the air, into a deep-red solution. Chromic acid is a very active oxidizer, and when mixed with organic matter rapidly alters it, and if in slight excess will dissolve almost any form of tissue. It is no doubt in this way that it acts as an escharotic. It is very much used to destroy *condylomata* and other *dermal growths*, and no doubt would be efficient in cases of larger tumors. In superficial affections it is best applied by means of a glass rod, the liquid formed by the spontaneous deliquescence of the crystals being used.

BROMINIUM.—*Bromine* is a dark-red liquid which has a very powerful, disagreeable, chlorine-like odor, and at ordinary temperatures emits exceedingly acrid, pungent fumes. It is sparingly soluble in water, more soluble in alcohol, and still more so in ether. Alcoholic and ethereal solutions lose their color and become acid in a few days, from the generation of hydrobromic acid. When bromine is brought into contact with organic matter, it oxidizes and completely destroys it with great rapidity. On account of its liquid form and of this property, bromine is one of the most severe, thorough, and rapid of all the caustics. It has not been much employed to destroy morbid growths, but during our late war was found to be the most efficient of all the applications tried in *hospital gangrene*. After most of the slough had been cut away, the caustic was applied pretty freely to the living tissue by means of a glass rod. The pain was very severe; but it ceased after a few moments when the caustic was washed off.

Sulphate of Zinc and *Sulphate of Copper* are feeble escharotics, never used to destroy sound tissue. *Burnt Alum* belongs in the same category, but is probably a little more feeble. All three of these drugs are used to destroy *exuberant granulations* in ulcers.

CLASS XIII.—DEMULCENTS.

THESE are bland substances, which form more or less gummy or mucilaginous solutions in water, capable of exerting a calming or soothing influence upon inflamed surfaces. Without doubt water itself is the demulcent *par excellence*; but the remedies here discussed do seem to enhance its power. It has been claimed for these medicines not only that they soothe surfaces to which they are immediately applied, but also that taken internally they relieve irritation in distant organs. As, however, all of them are complex vegetable products, as many of them are staple articles in the world's food, and as none of them have been detected in the blood or the secretions, it cannot be allowed that they reach distant parts through absorption into the blood. What is certainly true of some of them is probably true of all,—*i.e.*, digestion of them occurs in the *primæ viæ*. The relief which undoubtedly follows their use in certain affections of parts which they can reach only through the circulation is probably in great part, if not altogether, due to the large quantities of water with which they are administered, which, passing through the body, lessen the concentration, and hence the acidity, of the urine and other secretions.

Clinically, demulcents are useful as local applications in all forms of acutely inflamed surfaces, and they are taken internally in acutely *inflammatory* conditions of the *alimentary canal*. In slight *bronchial irritation* they are often of service, especially when allowed to dissolve slowly in the mouth: used in this manner, they not only exert an influence upon the mucous membrane of the mouth, but very probably find their way into the respiratory passages.

ACACIA—GUM ARABIC.

A gummy exudation from *Acacia vera* and other species of *Acacia*, small trees growing in Northern Africa, Senegambia, Guinea, etc., the Cape of Good Hope, and Australia. Gum arabic occurs in roundish or irregular pieces, more or less transparent, hard, brittle, varying in color from white or yellowish white to red, or even deep orange brown. It consists of a peculiar principle, *Arabin*, united with about three per cent. of lime, potash, and magnesia. According to Husemann, pure *arabin* is an amorphous substance,

glassy and transparent when dry, but milk-white when moist, and having a feeble acid reaction, with the power of uniting with bases.

In the plant, arabin, like other gums, appears to be formed by a retrograde metamorphosis of cellulose. Indeed, Wigand declares that the flesh and even the hard endocarp of the plum can undergo this metamorphosis. The same investigator affirms that *bassorin* is a first product of the change, and is, by a continuation of the process, converted into arabin.*

THERAPEUTICS.—On account of its solubility and pleasant taste, gum arabic is often used as a demulcent in irritation of the fauces and in *angina*. It is also sometimes used as an addition to drinking-water in fevers, and is believed to have some slight nutritious properties. Its chief use is, however, to aid in the suspension of oils and other substances in emulsions, in the making of pills, and in other pharmaceutical operations. The *Mucilage* (*Mucilago Acaciæ*) and the *Syrup* (*Syrupus Acaciæ*) are official.

TRAGACANTHA.—*Tragacanth* is the concrete juice of *Astragalus verus*, a small shrub of Asia Minor. Tragacanth occurs in large, whitish, horny, waved flakes, or sometimes in filamentous pieces. It is odorless and nearly tasteless. Introduced into water it does not dissolve, but swells up into a soft paste. One hundred parts of it contain, according to Guérin, 53.3 parts of arabin, 33.1 parts of bassorin, and 2.5 parts of inorganic ash. Bassorin is a gummy principle, at once distinguished from arabin by its not dissolving in water, but simply swelling up into a pasty mass. Tragacanth is used in medicine only in the manufacture of troches and in suspending heavy powders, for which purposes the difficulty of its solution and the extreme viscidility of its mucilage especially fit it. Its *Mucilage* (*Mucilago Tragacanthæ*) is official.

ULMUS FULVA.—*Slippery Elm* is the inner bark of *Ulmus fulva*, a large indigenous tree. The bark is of a yellowish-white or tan color, fibrous, yet when dry somewhat brittle, and occurs in long, flat strips or pieces one or two lines thick. It is pleasantly mucilaginous when chewed. It contains a large quantity of a peculiar mucilage, which it yields freely to water. Its infusion is sometimes taken in large quantities in *inflammations* of the *intestines* as a demulcent laxative; but its chief use is as an external application. When ground into powder, slippery elm makes a very excellent soothing poultice. The *Mucilage* (*Mucilago Ulmi*) is official.

* The reader interested in the method of formation of the gums is referred to the following papers:

Kützing, *Archiv d. Pharmacie*, 1851, Heft i. Translated in the *American Journal of Pharmacy*, vol. xxv. p. 37.

Mohl, *Botanische Zeitung*, 1857, p. 33.

A. Wigand, *Jahrbücher für Wiss. Botanik*, 1861, iii. p. 117.

And for a general discussion by Dr. J. Sachs, to *Hofmeister's Handbuch der physiologischen Botanik*, Bd. iv. p. 367.

CETRARIA.—*Iceland Moss* is the fronds of a lichen, *Cetraria Islandica*, growing on rocks in Iceland, and in most of the northern portions of the world. It is said to be abundant in the mountains of New England. The foliaceous, dry, shining, lobed, and lacinated fronds are about four inches long, of various intermixed colors, gray, brown, and red, of a mucilaginous, bitter taste. Iceland moss contains a peculiar starch-like principle, lichen starch, and a bitter principle. It is inodorous, and has a mucilaginous, bitter taste. It yields to cold water its bitterness; to boiling water all of its virtues.

Cetrarin, or, more correctly, *Cetraric Acid*, is the bitter principle, which may be obtained as a snow-white mass of interlaced acicular crystals. It unites with alkalies to form salts. With it in the lichen is associated in small quantities *lichstearic acid*. *Lichenin*, or *Lichen Starch*, the mucilaginous, nutritive principle of Iceland moss, differs from ordinary starch in not being deposited in granules within the cells, but in layers or irregular masses between the cells, or indeed forming the walls themselves of the cells (De Bary, *Hofmeister's Handbuch der physiologischen Botanik*, Bd. ii. p. 255). In cold water it swells up without dissolving; in hot water it dissolves, and on cooling condenses into a jelly. With iodine it strikes a yellow, green, or sometimes rather faint blue color. It is found in very many lichens; also in many species of sea-weed, notably in the so-called *Corsican moss*.

Iceland moss has enjoyed some reputation as a demulcent in pectoral complaints. From its bitter principle it is probably somewhat tonic, and its lichenin is probably about equal to ordinary starch as a nutrient. When prepared as an article of diet, in the form of jelly, the bitter taste should be removed by soaking for some hours in a very weak cold alkaline solution, and afterwards for a little while in cold water.

CHONDRUS.—*Irish Moss*, or *Carrageen*.—The fronds of *Chondrus crispus*, a sea-weed growing on the coast of Ireland, and also on the northern coast of the United States, where it is now gathered in large quantities. The fronds are purplish red,—but, as kept in the shops, bleached by washing in fresh water, whitish, and translucent,—cartilaginous, slender, much branched, swelling up but not dissolving in water, and having a slightly saline taste. Their virtue depends chiefly upon a starch- or gum-like principle, *Carrageenin*, which is distinguished from starch by not turning blue with iodine, from gum by not precipitating from its watery solution on the addition of alcohol. *Chondrus* also contains a notable proportion of a vegetable albumen, very rich in nitrogen.

Carrageen, being demulcent and nutritious, is employed as an article of diet in those cases requiring food of such character, and may be used instead of arrow-root. It is to be prepared by first soaking for ten minutes in cold water, and then boiling from half an ounce to an ounce of it (according to the desired consistency) in a pint and a half of water down to a pint, sweetening and flavoring to taste. Milk may be substituted for water, and the preparation thus rendered more nutritious.

GLYCYRRHIZA.—*Liquorice Root* is the root of *Glycyrrhiza glabra*, a native herb of Southern Europe. It occurs in long cylindrical pieces, from a few lines to more than an inch in diameter, brownish externally, and yellowish within. Its fracture is fibrous, its taste sweet and mucilaginous, its odor none. Its active principle is *Glycyrrhizin*. This is a sweet, neutral substance, differing from the sugars in not being converted by nitric acid into oxalic acid, and by its inability to undergo the vinous fermentation. Liquorice root is very largely used as a demulcent in pectoral complaints, and, on account of its pleasant taste, as a means of disguising or of flavoring medicines. In the form of glycyrrhizin it is said to conceal almost entirely the bitter taste of quinine and similar substances. It is used almost exclusively in the form of the *Extract* (*Extractum Glycyrrhizæ*), known as *Liquorice*. The *Mistura Glycyrrhizæ Composita*, or *Brown Mixture*, contains paregoric, antimonial wine, and sweet spirit of nitre, and is much used as a domestic remedy in "colds" and the early stages of *mild bronchitis*. The dose for an adult is half a fluidounce to an ounce every three hours; for a child three years old, a teaspoonful.

MEDULLA SASSAFRAS is the pith of the indigenous sassafras, which yields a delicate mucilage with water. This mucilage, which is much used in diseases of the eye, is official (*Mucilago Sassafras*).

LINUM, or *Flaxseed*, is the seed of *Linum usitatissimum*, or common flax, and contains large quantities of mucilage and of oil: its *Compound Infusion* (*Infusum Lini Compositum*, $\frac{3}{4}$ ss to Oj: containing also liquorice root) is much used internally. It is often made with boiling water; but the application of too much heat causes the extraction of more or less of the oil, and thereby renders the preparation less palatable. The infusion should never, therefore, be boiled during its making. The addition of lemon and sugar renders it more palatable. It may be drunk *ad libitum* in pectoral catarrhs, in enteritis and dysentery, and in irritation of the kidneys or urinary passages. *Ground flax-seed* (*Lini Farina*) is especially adapted, by the oil and mucilage it contains, for making poultices.

The seeds of the *quince* (*Cydonium*, U. S. Pharmacopœia secondary list) also contain a large quantity of mucilage, which they readily yield to water, forming a thick tenacious solution.

TAPIOCA is the fecula of the root of *Janipha Manihot*, a native of South America. There are two varieties of the tapioca plant, the *sweet* and the *bitter cassava*. The latter is in its fresh state poisonous, from the prussic acid which it contains, but yields most, if not all, of the tapioca of commerce. The tapioca is obtained by allowing the expressed juice to stand, separating the powder which deposits, washing, and drying by heat, owing to the action of which the starch is rendered partially soluble in cold water.

Tapioca is in irregular, hard and rough, nearly tasteless grains, which the

microscope shows to be composed of broken, ruptured starch-granules. It constitutes an elegant farinaceous article of diet, and may be prepared in the following manner :

Soak two tablespoonfuls of very clean tapioca in two teacupfuls of cold water over-night ; in the morning add a little salt and one pint of milk, or water if milk is not allowed ; simmer it until quite soft ; stir well while cooling ; when done, pour into a bowl, and add sugar, wine, and nutmeg, according to taste or the exigencies of the case.

MARANTA (*Arrowroot*) is a fecula or starch, obtained from the rhizome of the *Maranta arundinacea*, a native of the West Indies. Arrowroot is produced in the West Indies, Africa, and in the Southern United States, especially Florida. It is obtained in the usual method by beating the root-stocks into a pulp, and the use of flowing cold water. The most esteemed variety is that which comes from the island of Bermuda.

Arrowroot occurs as a very light, tasteless, and odorless white powder, which the microscope shows to be composed of ovate, oblong, or irregularly-convex granules, from the seven-hundred-and-fiftieth to the two-thousandth of an inch long, marked with very fine rings and a circular hilum, which cracks in a linear or stellate manner. It is often adulterated with other starches, which are best detected by the microscope.

Arrowroot affords the most readily prepared and, in the sick-room, the most popular of all the farinaceous articles of diet. It may be prepared in the following manner :

Stir two teaspoonfuls of arrowroot in a teacupful of cold milk until a *perfectly smooth* mixture is made ; have on the fire a pint of milk, and, while this is boiling, add the arrowroot little by little, stirring constantly until cooked ; add sugar, nutmeg, and wine, according to taste or the exigencies of the case.

SAGO is a fecula prepared from the pith of *Sagus Rumphii* and other sago palms of Sumatra and the neighboring regions. It is first prepared in the usual manner, and then formed into a paste with water, rubbed into grains, and dried. *Pearl Sago* occurs in hard, roundish, somewhat translucent or sometimes opaque grains, about the size of a pin's head. *Common Sago* is in larger and browner grains, often mixed with some powder. It is composed of oval or ovate, often truncate and muller-shaped, often much broken, starch-granules. It is used exclusively as an article of diet, forming a jelly, which is best prepared as follows :

Wash the sago well in cold water ; put a small teacupful of it to soak in half a pint of water over-night, and in the morning put this mixture into one pint of hot water ; squeeze into it the juice out of a thinly-pared lemon, and allow to simmer slowly for twenty minutes ; then sweeten, add wine according to taste or the exigencies of the case, then pour into moulds to cool.

The U. S. Pharmacopœia, under the name of HORDEUM, or *Barley*,

recognizes the decorticated seeds of the common barley, the pearl barley of commerce. They contain starch and mucilage, and the *Decoction* (*Decoctum Hordei*) is official. *Barley-water* is used as a nutritious, demulcent drink in fevers and inflammatory conditions, especially when the gastric mucous membrane is involved. The U. S. Pharmacopœia directs that it should be prepared as follows:

Take of barley two troyounces; water, a sufficient quantity. Having washed away the extraneous matters which adhere to the barley, boil it with half a pint of water for a short time, and throw away the resulting liquid. Then, having poured on it four pints of boiling water, boil down to two pints, and strain.

CLASS XIV.—EMOLLIENTS.

TRUE emollients are perfectly bland, fatty substances which, when applied to the skin, soften it and render it more pliable. The action of these remedies is largely mechanical, and they probably soften the derm in precisely the same way as they affect a raw hide or a piece of leather. They are therefore especially useful when the skin has a tendency to crack or to chap. Whenever surfaces become sore by attrition, or, in other words, chafe, emollients are also useful in a mechanical method. Emollients often afford relief in simple inflammations of the skin under such circumstances that their action cannot be explained in any of the ways alluded to: indeed, they seem to exert a dynamic influence upon the nutrition of the parts concerned. To the best of our present knowledge, the oxygen or other constituent of the air acts as a stimulant to exposed surfaces and increases inflammations there present; and it is believed that fatty matters do good by keeping out the air. If this be so, the apparently dynamic influence of fatty matters is, after all, mechanical. Be these things as they may, clinical experience has demonstrated that fatty matters are of very great value in the treatment of superficial inflammations. It must be borne in mind that the blandest fat, when it becomes rancid, is very irritating, and will do far more harm than good. In the use of fatty emollients, the strictest attention must therefore be paid to the condition of the material employed. Any perfectly bland oily substance may be used as an emollient. *Mutton suet* and *goose-grease* are famous in domestic medicine, but are simply valuable because, if well prepared, they are less apt than some other fats to become rancid. It cannot be allowed that there is any difference in fats, unless it be in penetrating power: a very hard fat is of course not so readily applied as a softer one, and therefore only such fats as freely melt, or at least become very soft, at the temperature of the body, are to be used. *Cold Cream* (*Unguentum Aquæ Rosæ*), containing oil of sweet almonds, spermaceti, white wax, and rose-water, is the most elegant of all the officinal fatty emollients, unless it be glycerine, which is so much used that it merits a distinct notice.

GLYCERINA—GLYCERINE.

This is a thick syrupy liquid, colorless, and free from odor, but of a sweet taste. It was originally prepared by heating together a metallic oxide and

ordinary fats, as in the manufacture of lead-plaster. Chemically speaking, it belongs to the alcohols, and is known, according to the recent nomenclature, as *propenyl alcohol*. It is always set free during the process of saponification, when the stearic or other acids of the fat unite with some base to form salts, known as soap. Consequently, glycerine always forms a considerable part of the soap-maker's waste; but in this it is so mixed with impurities that, until recently, it was valueless. Now, by means of certain patent processes, the glycerine is said to be obtained pure from the waste products of soap-factories. It is also manufactured on a very large scale by the direct decomposition of fats by heated steam, in accordance with a plan patented by Mr. Tilghman. The great bulk of the glycerine in commerce is at present prepared in this way.

Under certain circumstances, not well understood, glycerine forms hard, brilliant crystals. In its usual liquid form it mixes in all proportions with water and alcohol, and possesses itself very great solvent powers, dissolving iodine, bromine, the alkalies, tannic and other vegetable acids, a large number of neutral salts, salicin, and very many other organic principles, so that it is largely used in pharmacy. The following table by Klever gives the amount of various drugs dissolved by one hundred parts of glycerine :

Acidum arseniosum.....20	Cupri acetat.....10	Potassii iodidum.....40
“ arsenicum20	“ sulphas.....30	Quinia 0.59
“ benzoicum.....10	Ferri et potassii tartras.. 8	Quiniæ tannas..... 0.25
“ boracicum.....10	“ lactas.....16	Sodii arsenias.....50
“ oxalicum.....15	“ sulphas.....25	“ bicarbonas..... 8
“ tannicum50	Hydrarg. chlor. corros.. 7.50	“ boras60
Alumen40	“ cyanidum.....27	“ carbonas.....98
Ammonii carbonas.....20	Iodinium 1.90	“ chloras.....20
“ chloridum.....20	Morphia 0.45	Sulphur..... 0.10
Antimonii et potas. tart. 5.50	Morphiæ acetat.....20	Strychnia 0.25
Atropia 3	“ murias20	Strychniæ nitras..... 4
Atropiæ sulphas.....33	Phosphorus 0.20	“ sulphas.....22.50
Barii chloridum.....10	Plumbi acetat.....25	Urea50
Brucia2.25	Potassii arsenias.....50	Veratria 1
Calcii sulphis 5	“ bromidum25	Zinci chloridum50
Cinchonia 0.50	“ chloras..... 3.50	“ iodidum.....40
Cinchoniæ sulphas..... 6.70	“ cyanidum32	“ sulphas.....35

Glycerine does not evaporate upon exposure, but is very hygroscopic, and absorbs water from the air. When pure, it is incapable of becoming rancid or of fermenting spontaneously. The acrid glycerine, formerly so abundant in commerce, and still met with, owes its irritant properties to contaminating substances, especially, it is said, to oxalic and formic acids. The former of these substances is apt to be created by the action of the sulphuric acid used during the purification of the glycerine, the latter by the reaction between the oxalic acid and the glycerine.

THERAPEUTICS.—Locally applied, glycerine is perfectly unirritating, and is much employed as an emollient. The chief disadvantage that attends its use

is its stickiness; on the other hand, its non-volatility and its hygroscopic properties give a persistency to its action which is often very advantageous. It enters largely into the composition of popular emollient ointments, or "creams," as they are called, and is often used itself for chapped hands, excoriations, and similar troubles. It is also employed by dermatologists to some extent in *chronic eczema*; in *seborrhœa*, whether affecting the hairy scalp or other parts, it is asserted to be especially useful, softening the masses of secretion, and, used in conjunction with such remedies as borax, zinc, and acetate of lead, diminishing the amount of secretion. When there is a want of sebaceous secretion, it is said also to act efficiently; in *scabies*, *pruritus*, and even *psoriasis*, glycerine is used, diluted with water, as a vehicle for more active remedies.

Upon the mucous membranes glycerine acts very much as it does upon the skin, and diluted with water is very useful in *coryza*, and even, by enemata, in *dysentery*; in *croup* or *laryngitis* it may with advantage be applied freely by means of a large camel's-hair brush to the orifice of the larynx, so as to run into the latter. It also forms an excellent basis for mouth-washes; or a paste may be made with it and borax, or similar substances, for use in ulcerations of the same cavity. The list of diseases in which this substance is employed might be very much lengthened; but the examples already given are sufficient to indicate the range of its application as an emollient and a vehicle.

There are certain persons upon whose skin and mucous membranes even the purest glycerine seems to act as an irritant. This influence is most intense when the glycerine is nearly or entirely free from water. It is, however, discernible even when the remedy is much diluted, and often prohibits its use. The existence of this idiosyncrasy to glycerine can be determined only by trial.

When administered internally in doses of one or two ounces, glycerine acts as a gentle and very uncertain laxative. I do not think that besides this action it has any perceptible effect. It was at one time proposed as a substitute for cod-liver oil in *cachectic diseases*, but, after pretty extensive trials by various clinicians, its employment as an alterative tonic has been universally abandoned. It has also been proposed and highly commended in *diabetes* (*London Lancet*, 1868; *Berliner Klinische Wochenschrift*, Aug. 1872). It appears to be harmless in this affection, and therefore may be used as a sweetening-material for coffee, tea, and other beverages; but there is no reason to believe that it exerts any control over the disease. The most important internal use of glycerine is as a harmless substance which has the power of disguising nauseous medicines. In this way it may be employed with castor oil, in emulsions of turpentine, in solutions of iron, and in various mixtures. It seems, as it were, to envelop the medicinal substances and prevent their acting on the palate.

POULTICES.—Under the head of Emollients I shall also speak of poultices, which are moist, soft, scarcely adhesive, perfectly bland plasters, used to a very

great extent to combat superficial inflammation. Poultices are much more powerful agents than the true fatty emollients, and are much more capable of being abused: the results of such abuse will be spoken of directly. A poultice may, of course, be stimulating and irritant if made of such a substance as mustard; but the emollient or true poultice is prepared out of some bland material, which is totally free from action upon the skin. I do not think that there is any difference whatever in the action of the various substances usually employed in the preparation of poultices, the latter depending for their remedial powers solely upon the warmth and water which they contain. Water, when pure and of a temperature approximating that of the body, is a sedative, checking all action, possibly by a direct influence, but probably by the merely mechanical acts of dilution of the pabulum and of separation of the germinal granules. It is also a relaxant, rendering all tissues soaked in it soft and yielding.

Poultices are sometimes applied in the early stages of phlegmonous and other superficial inflammations, for the purpose of checking the morbid action. Their influence is in such case simply one of sedation, and they are certainly not so efficient as the cold-water dressing. They are, however, especially useful in the advanced stages of inflammation, when suppuration has already commenced or is about to set in. Clinical experience has demonstrated that they now favor the formation of pus. It is hardly worth while to discuss how they do this, so long as the natural method in which pus is produced is a matter of dispute. If, as Cohnheim believes, pus is composed solely or largely of out-wandering white blood-corpuscles, it is evident that the relaxing influence of a warm poultice will greatly facilitate the escape of these bodies. Further, the poultice in the latter stages of a superficial phlegmon not only hastens the formation of pus in the inflammatory focus, but lessens irritation in the outlying parts by its sedative action, and so softens the tissues as to aid in the passage outwards and the discharge of the inflammatory products. When poulticing is persisted in for too long a time, the part becomes pale or white, swollen, relaxed, and has a sodden look. The granulations of the ulcer or abscess are large, pale, and very flabby, and all the vital actions are below the normal point. It is possible that even death of a part might be brought about by continuous poulticing. Be this as it may, after the discharge of pus, whenever the parts put on the aspect just spoken of, the poultice should be removed and stimulating applications be substituted.

So far, the use of poultices to combat external inflammations has alone been spoken of; but clinical experience has demonstrated their value in internal inflammations, even when such deep-seated tissues as the lungs are affected. Their action in these cases is somewhat different from that which they exercise over superficial inflammations. According to the dictates of experience, they should be applied very hot, and be frequently renewed; very often, too, a small amount of mustard or of some similar stimulating material is added to them with advantage. As a result, these poultices act

as gentle but deep-reaching counter-irritants, which in all likelihood affect not merely the blood-vessels of the skin, but also those of the subdermal tissue. When it is borne in mind that in all these cases the poultice is applied to a very large surface, it will be readily perceived that this counter-irritation is a powerful one. Thus, in *pleurisy*, or in *pneumonia*, the whole anterior or posterior surface of the chest is covered, or perhaps the whole chest is enveloped, by the jacket-poultice. In *peritonitis*, the poultice should be as large as the abdomen of the patient. Now, in either of the cases mentioned, it is plain that the amount of blood drawn to the surface must be considerable. I cannot help surmising that the water of the poultice in some cases actually soaks through and reaches the affected tissue so as to exert its direct sedative influence upon it. In lung-diseases of children, whose chest-walls are very thin, the value of poultices has seemed to me much greater than in corresponding affections in adults; and it is not illogical to believe that the difference may be dependent upon the inequality of the chest-walls.

Flaxseed meal is the most frequently-used substance for making poultices, for which purpose the large amounts of oil and mucilage which it contains especially fit it. Ground *slippery elm* makes a very elegant demulcent, mucilaginous poultice. Ordinary mush from *Indian meal* affords a cheap and very serviceable material, and *bread and milk* makes a popular, very mild and un-irritating, but expensive poultice. When a poultice is to be applied to affect internal organs, and consequently has to be large and capable of holding the heat for a long time, the choice of material lies exclusively between flaxseed and Indian meal. The former of these is the more adhesive, and makes the more manageable poultice; but popular belief, and I think with reason, attributes to mush a superior power of retaining heat. In either case the poultice should be put on as hot as it can be borne, and should be covered by a large piece of silk oil-cloth, which aids in retaining not only the moisture, but also the heat. The interval of renewal should be short, and should be governed solely by the rapidity with which the applied poultice grows cold.

CLASS XV.—DILUENTS.

By the term diluent is meant an indifferent substance which is absorbed and in its passage through the body simply dilutes the various fluids therein, and especially dilutes and renders less irritating the excretions. The only diluent is water, whether given in the form of mucilaginous or other drinks, or of simple water. The natural medicinal waters owe much of their value to the large quantity of water they contain. It is evident that when a quart of water more than usual is taken into the system, it must whilst it stays there lessen the concentration of the bodily liquids, and must finally in some way find an exit from the body.

The increased excretion even of water means increased action in the eliminating glands; and the water passing out of the blood must always carry with it more or less of the soluble matters contained in the same, so that whilst the percentage of solid matter in the urine or in the sweat may be lessened by large potations of water, the actual amount eliminated is no doubt increased. Hence water acts not only as a diluent, but also as a depurant.

It is especially in regard to the urinary organs that water is employed medicinally, and it is evident that there are three distinct objects to be attained. In the first place, it may be used simply to aid the re-establishment of completely or partially suppressed renal secretion.

During a recent visit to Bellevue Hospital, New York, I was informed that in *acute Bright's disease* water had been found to be a most efficient diuretic, increasing very remarkably the urinary excretion, relieving the irritation of the kidneys, and aiding the return to health. In these cases at least half a pint of water should be taken every two hours. I have tried it in chronic Bright's disease, and found it to work well in some cases; but in others, no increase of the urine taking place, the water accumulated in the system and added to the distress.

A second object to be attained by the free internal use of water is the dilution of the urine in cases in which it is very acid and irritant, or when it is so supersaturated with salts as to have a tendency to deposit them in the form either of gravel or of calculi.

A third use of water is as a depurant in chronic cases of digestive or other disorders, in which there is no organic disease, but habitual torpor of the emunctories,—cases in which the liver is said to be torpid, and in which there

is a foul tongue, habitual costiveness, and a scanty urine with a tendency to the formation of a lateritious deposit. In these cases a couple of tumblerfuls of water upon rising will often produce a stool after breakfast, as well as increase the flow of urine. Very generally a mild saline may be advantageously added in small quantity to the water. It is in these cases that such natural mineral waters as those of Saratoga are especially beneficial.

CLASS XVI.—PROTECTIVES.*

IN the present class are included those materials used by the physician as external applications to exclude the air and to protect inflamed dermal or other tissues. Sufficient has already been said in regard to the importance of the exclusion of the air, under the heading of Emollients. It is evident that the latter class of remedies, as well as demulcents, act as protectives; but the class of Protectives proper seems necessary for the consideration of certain remedies which act more plainly in a mechanical method, and which will defend the skin against external agencies.

First to be considered under the present class are certain plasters, used to protect the skin and raw surfaces from external influences. The *adhesive plaster* (*Emplastrum Resinæ*) is used enormously for mechanical purposes. It does, however, irritate the skin somewhat, and consequently is rarely employed where protection is the only object. Under the latter circumstances, the *lead plaster* (*Emplastrum Plumbi*) or the *soap plaster* (*Emplastrum Saponis*) is preferable. These substances are free from irritant properties, but are very slightly adhesive, and are scarcely used except to protect the skin from pressure or friction, as when *bed-sores* are threatened. They should be spread upon very soft kid. It is also very important that they be not so thick or hard as to lose their pliability. If they are stiff, by their movements during the motions of the body they often do more harm than good.

COLLODIUM—COLLODION.

When any finely-divided ligneous body, like raw cotton, is steeped for a few minutes in a mixture of nitric acid of the specific gravity of 1.5 and concentrated sulphuric acid, and then squeezed, thoroughly washed, and dried, it gains about seventy per cent. in weight, and is converted into a substance known as *pyroxylin*, or *gun-cotton*. The change consists in the substitution of nityl (NO_2) for a portion of the hydrogen. There are a number of varieties of gun-cotton. The true explosive gun-cotton, that which is alone adapted for gunnery, is *trinitrocellulose*. It is not so soluble as the less highly nitrated variety, and is not fit for use in medicine. The officinal pyroxylin

* This class was omitted by mistake in the enumeration on page 21.

(*Pyroxylin*) is one of the more soluble varieties of gun-cotton, but is not soluble to any extent in ether.

Gun-cotton is a perfectly inert substance, so far as the external surface of the skin is concerned, and probably has no effect upon the system when taken internally. It is not at all soluble in water, and its only use in medicine is in the manufacture of collodion. This substance is officinally prepared by dissolving two hundred grains of pyroxylin in a mixture of twelve and a half fluidounces of stronger ether and three and a half fluidounces of stronger alcohol. It is a colorless, slightly opalescent liquid, of a syrupy consistence, and smelling strongly of ether. By long standing it deposits a layer of fibrous matter, and becomes more transparent. This layer should be reincorporated, by agitation, before the collodion is used. When it is applied to the skin, and the menstrua are allowed to evaporate, collodion forms a colorless, transparent, flexible, and strongly contractile film, which adheres very closely, and cannot be readily removed by washing, motion of the part, or external mechanical force. As this coating is perfectly impervious to air and water, collodion is much used in surgery for various purposes. It is evident that care should be used in its application to abscesses and discharging wounds, lest it should prevent the discharge of pus. Small fresh wounds are often very advantageously dressed in the following manner, especially cuts on the fingers and about the head: If necessary, the hair should first be shaved off the part, and then a piece of coarse gauze or mosquito-netting, of suitable size and shape, should be laid so as to cover the wound and extend across each side from half an inch to an inch and a half, according to circumstances. One end should then be tightly fastened to the skin by repeated applications of the collodion with a camel's-hair brush. When the adhesion has become sufficiently firm, the gauze should be drawn so as to close the wound tightly, and whilst it is held in position the collodion should be applied all over it. As the collodion contracts during drying, the wound is more and more tightly closed, until at last a firm, perfectly tight dressing binds it close together. Of course in deep wounds this method of closure is not to be used.

The contraction of the collodion film is a great drawback to its use for certain purposes. It can be in great measure obviated and the film made more pliable by the addition of from eight to ten drops of castor oil to the ounce of the liquid. Under the name of *Collodium Flexible*, the U. S. Pharmacopœia directs a preparation very similar to that just spoken of. The official *flexible collodion* contains twenty grains of Canada turpentine, besides ten grains of castor oil, to the ounce; and probably the stimulant effect of the terebinthinate will make itself apparent upon some susceptible skins. Whenever collodion is used simply as a protective, one of these modified preparations is much preferable to the pure article.

Any principle which is soluble in a mixture of ether and alcohol may be added to collodion, and in this way medicinal substances may be applied to

the external surfaces. The films formed are often less firm and adhesive than those of the simple collodion.

LIQUOR GUTTÆ-PERCHÆ.—*Solution of Gutta-Percha* is made by dissolving small pieces of gutta-percha in purified chloroform. When it is applied to the skin, a thin elastic adhesive film is left, which protects the parts from the air. This is said to be a very elegant preparation for use in *small cuts, abrasions, chapped lips*, and the little injuries which come within the province of domestic medicine rather than in that of the professional art.

PART II.

NON-SYSTEMIC REMEDIES.

THESE are drugs which act not directly upon the human system or upon any of its tissues, but upon some extraneous material or entity, either in the cavities of the body or upon its exterior. Thus, an antacid neutralizes acid in the stomach, or an anthelmintic kills the tapeworm in the intestines, or a disinfectant destroys poisonous emanations in the exterior world and thereby wards off disease.

CLASS I.—ANTACIDS.

ANTACIDS are, strictly speaking, substances which are capable of neutralizing acid. The class, as here defined, contains those remedies which in medicine are used for the purpose of neutralizing an excess of acidity in the *primæ viæ*. They are almost solely employed in forms of *dyspepsia*. Without doubt, *cardialgia*, *gastric uneasiness*, "*heartburn*," and the rising of sour water in the mouth, are often the result of too much acid in the stomach, perhaps secreted by a perverted glandular action, but more probably in the great majority of cases formed by fermentative changes in the partially-digested food.

As this excessive acidity of the stomach causes gastric uneasiness and derangement, so will a similar condition of the intestinal canal cause pain and spasm and functional disturbance in the bowels. This is seen most frequently in infants, and is very often associated with a diarrhœa in which the passages have a green color, similar to that of spinach, and hence are sometimes spoken of as "*spinach-stools*." In *diarrhœa* of this character, as well as in *colic*, antacids are often of service by neutralizing the acid in the intestinal canal.

Clinical experience has demonstrated that dyspepsia is often permanently relieved by the use of alkalies when they are given steadily day after day, about twenty minutes after eating, for a long time. According to Dr. Thomas K. Chambers (*The Indigestions*, Am. ed., 1870, p. 67), this is dependent upon an effect pointed out by Claude Bernard, the augmentation of the acid gastric juice, and so of the normal peptic powers of the stomach. The same

authority further says, "The test of benefit being derived from an alkali is the dose not requiring to be increased as the patient goes on taking it. but, on the contrary, being diminished gradually, while relief from the recurrence of heartburn continues still to be experienced."

Sick headache is sometimes dependent upon gastric irritation produced by an excess of acid in the stomach. This true sick headache is generally to be distinguished from migraine by the early occurrence of the stomach-symptoms, either as heartburn, nausea, vomiting, or simple gastric distress, and by the facts that the pain comes on with an attack of blindness or of dizziness, and is not limited to any one spot, as the supraorbital or other neuralgic foci, but is felt all across the brows. In this form of cephalalgia, antacids often afford very great relief.

Various substances which have already been discussed in this work are excellent antacids, most of them uniting this to other medicinal properties. Thus, when a stimulating antacid is desired, as is very often the case in sick headache, half a drachm of the *aromatic spirit of hartshorn* may be taken, well diluted with water. Again, when a laxative antacid is desired, a teaspoonful to a tablespoonful of *magnesia* may be exhibited.

Potassa and the carbonates of potassium have already been dwelt upon with sufficient detail. They may be used as antacids; but, as they exert other powerful influences upon the system, they are, I think, not so generally useful as the soda preparations. Nevertheless, the *Solution of Potassa* (*Liquor Potassæ*) is largely employed as an antacid. It is a colorless water-like liquid, of a strong, acrid, alkaline taste, and is made by boiling a solution of the bicarbonate of potassium with lime. It contains only five and eight-tenths per cent. of the alkali, but acts upon animal and vegetable substances, and imparts a distinct soapy feel to the fingers when they are moistened with it and rubbed upon one another. It is capable, in overdose, of acting as an irritant poison. The dose is ten to twenty minims, well diluted.

SODIUM. (Na.)

Unlike potassium, sodium and its salts have very little influence upon the higher animals. Frogs, according to the experiments of Поджаеков (*Virchow's Archiv*, Bd. xxxiii. p. 507), are more susceptible to their action; for when a quantity double the lethal dose of potash was injected into the cellular tissue, the batrachians slowly succumbed, the chief symptoms being muscular tremblings and convulsions.

On the other hand, Grandeau (*Robin's Journal de l'Anatomie*, 1864) found that one hundred and seven grains of the carbonate of sodium injected into the vein of a dog produced only very slight symptoms, and that thirty-five grains of the nitrate similarly administered to a rabbit caused only some convulsive movements.

According to Guttman (*Virchow's Archiv*, Bd. xxxv.), however, the soda

salts thrown directly into the blood in very large amounts will slowly cause death, the agony being very prolonged, and, when the chloride is used, convulsions being developed.

Both Podocæpow and Guttman assert that even the largest doses do not sensibly affect the heart or the temperature; and the latter observer further declares that they are without influence upon the nerve-centres, the peripheral nerves, or the muscles.

If this be the case, however, it is difficult to perceive how they can cause death; and the earlier experiments of Podocæpow indicate that they do exert a very feeble action upon the peripheral nerves or the muscles.

Upon the blood the immediate influence of the soda salts is very slight, for Podocæpow asserts that one part dissolved in twelve parts of blood does not affect either the physical characters of the red corpuscles, or the intensity of the ozone reaction.

The effect of the continuous exhibition for a few days of large amounts of salt upon the human organism has been elaborately investigated by Dr. Münch (*Archiv des Vereins für Gemeinschaftliche Arbeiten*, Bd. vi. p. 369, 1863), and found to be very feeble. At first there was a slight diminution of excretion, and a corresponding gain of the body in weight; but after a time the excretions increased, and the weight of the body decreased. The variations in excretion affected chiefly the urine, but sometimes the perspiration and feces were also influenced. The urine was rendered alkaline, but its solid ingredients were scarcely at all affected.

Although a certain amount of the soda salts is a necessary food for the higher animals, yet it is very doubtful whether an habitual excess of them has any effect upon nutrition. The experiments of Plouriez (*Comptes-Rendus*, t. xxv., 1847) would seem, however, to indicate that the chloride at least exerts a tonic influence. The blood of this observer was analyzed after he had used daily one hundred and fifty grains of salt beyond the ordinary amount for three months, and again after he had taken only the usual quantity. The red corpuscles were very decidedly more numerous, the fibrin slightly more abundant, and the albumen decidedly less abundant at the first analysis than at the second.

As Bidder and Schmidt (*Constatt's Jahresbericht*, 1852) assert that the hydrochloric acid of the gastric juice is derived from the chloride of sodium, and as Rabuteau (*L'Union Médicale*, t. xii. p. 186, 1871) found that, in dogs with gastric fistula, both the quantity and the acidity of the gastric juice are decidedly increased by the use of salt meat, it would appear probable that common salt acts as a tonic by increasing the digestive power. On the other hand, it is well known that physiologists are still disputing as to whether free hydrochloric acid exists at all in the gastric juice.* Further, it is certain

* See Bernard's *Physiologie Expérimentale*, t. ii. p. 393, and, for a discussion of the subject, Prof. F. G. Smith's note to the American edition of *Marshall's Physiology*, p. 530; also Longet's *Physiologie*, Paris, 1861, t. i. p. 196.

that pepper or any similar condiment will, by its local action, increase the flow of gastric juice; and in Rabuteau's experiments the mere local irritation of the salt, or the difficult digestion of the preserved meat, may have given rise to the increased secretion by the stomach.

SODII CARBONAS—CARBONATE OF SODIUM ($\text{NaOCO}_2 - \text{CO}_3\text{Na}_2$).

This salt occurs in colorless crystals, which rapidly effloresce on exposure to the air, and fall into a white powder. The taste and reaction are strongly alkaline. It is very soluble in water, insoluble in the air. The chief native materials from which it is manufactured are common salt and the sulphate of sodium. As it occurs in commerce, the carbonate of sodium is sufficiently pure for ordinary use. When the carbonate of sodium is heated, its water of crystallization is driven off, and the officinal *Dried Carbonate* (*Sodii Carbonas Exsiccata*) is left. This may be given in pill-form.

LIQUOR SODÆ—SOLUTION OF SODA.

A colorless, exceedingly acrid liquid, containing five and seven-tenths per cent. of hydrate of soda, and having the specific gravity 1.071. It is made by the action of milk of lime upon a solution of the carbonate.

SODA ($\text{NaO} - \text{ONa}_2$).—*Caustic Soda* is prepared by the evaporation of the liquor, and occurs in grayish-white fragments, which deliquesce when exposed to the air; but as the fluid absorbs carbonic acid, and the soda in it is converted into an efflorescent carbonate, the liquid is after a time converted into a white powder.

Caustic soda is powerfully corrosive.

SODII BICARBONAS VENALIS—*Commercial Bicarbonate of Sodium*.—This is a white, opaque powder, containing variable amounts of soda not fully saturated with carbonic acid.

SODII BICARBONAS—*Bicarbonate of Sodium* ($\text{NaO}, 2(\text{CO}_2 - \text{CO}_3\text{NaH})$).—As directed by the U. S. Pharmacopœia, this is prepared by slowly passing six pints of water through sixty-four troyounces of the commercial bicarbonate upon a filter, the process being suspended so soon as the filtrate ceases to give a precipitate with the sulphate of magnesia. In this way the carbonate of sodium is washed out of the bicarbonate, because it is much more soluble in water. As the carbonate of sodium produces a precipitate with the sulphate of magnesium, whilst the bicarbonate does not, the magnesia salt offers a ready method of determining when the filtrate is free from the carbonate.

PULVERES EFFERVESCENTES.—*Effervescing or Soda Powders* contain in separate papers thirty grains of the bicarbonate of sodium, and twenty-five grains of tartaric acid. When these powders are dissolved separately, and their solutions mixed, active effervescence occurs, during which the remedy is to be

taken. These powders are said to be slightly laxative, and, I believe, are capable of rendering the urine alkaline.

THERAPEUTICS.—The fact that soda, in moderate amount, has no depressing action, and indeed very little, if any, influence upon the general system, renders it preferable to potash in cases of acidity of the *primæ viæ*. It is *par excellence* the alkali for acid dyspepsia. On the other hand, the circumstance clearly established by Dr. Roberts (*Urinary and Renal Diseases*, Am. ed., 1866, p. 240), that it is less powerful as a solvent of uric acid than its sister alkali, together with the property, believed to belong in a much greater degree to potash, of preventing the formation of uric acid, makes soda of very inferior value in *uric acid gravel* or *diathesis*. When in any case it is desirable simply to render the urine alkaline, and at the same time to avoid depressing the system generally, soda would, on theoretical grounds at least, seem preferable.

SODII ACETAS.—The *Acetate of Sodium* is a white, slowly efflorescent salt, which occurs in long prisms of a sharp, bitterish taste. It has been supposed to have the same remedial powers as the acetate of potassium; but this is certainly a mistake. It is probably capable of rendering the urine alkaline, but is rarely if ever used in medicine.

SODII NITRAS ($\text{NaO}, \text{NO}_5 - \text{NO}_3 \text{Na}$).—The *Nitrate of Sodium* occurs in colorless, rhomboidal crystals, slightly deliquescent, and wholly soluble in water. It is found native in the desert of Atacama, South America, and is officinal for the preparation of arseniate of sodium. It is scarcely used as a therapeutic agent.

CALX—LIME (CaO).

The therapeutic properties of lime are chiefly dependent upon its alkalinity and its astringency. It does not possess the latter property in an eminent degree, yet its preparations, when properly diluted, whiten and lessen the secretion of mucous membranes to which they are applied. It also appears to have, as it were, a sedative influence upon mucous membranes, lessening their irritability. Under the name of *Calx*, caustic or unslaked lime is included in the primary list of the U. S. Pharmacopœia. It is a powerful caustic, but, except in the form of Vienna paste, is rarely if ever used as such. It is more soluble in cold than in hot water. At 60°F . it requires about seven hundred times its weight of the liquid to dissolve. It is much more soluble in syrup than in pure water.

LIQUOR CALCIS—*Solution of Lime*.—Lime-water is simply a saturated solution of lime in water. It is a colorless liquid, of an alkaline taste, and nearly destitute of irritant properties. On exposure to the air it becomes turbid, or forms a crust upon the surface, or deposits a precipitate, owing to the absorption of carbonic acid from the air and the conversion of the lime into a carbonate.

THERAPEUTICS.—Lime-water is used internally exclusively for its effects upon the primæ viæ. In *vomiting* from almost any cause except acute gastritis, equal parts of it and milk afford an elegant, simple, and much-used remedy. If the vomiting be severe, all other food should be inhibited, and one or two tablespoonfuls of the mixture be given every half-hour; the quantity, as well as the proportion of milk, being increased as the stomach is able to bear it.

As lime-water when put in milk prevents the formation of dense coagula, it is often added with advantage to that fluid when used as food for infants, or for adults with weak digestion.

As an alkaline astringent, it is sometimes useful in the *diarrhœa* of children.

Externally, lime-water has been used as a wash in various skin-diseases, especially in *tinea capitis*: it is also applied to *ulcers*, and is said to have a very marked influence in lessening the amount of discharge. When mixed with an equal bulk of linseed or olive oil (*Linimentum Calcis*, U.S.P.: lime-water, eight fluidounces; linseed oil, seven troyounces), lime-water is a favorite application for recent burns: the thick, soapy liquid which is formed is sometimes spoken of as *Carron Oil*, from the name of the iron-works at which its reputation was first made.

Lime-water has the power of dissolving mucus, and also false membrane, and has therefore been introduced as a local remedy in *pseudo-membranous croup* and in *diphtheria*. It is sometimes used by causing the patient to inhale the vapors of slaking lime; but a better method is to pulverize lime-water by means of an atomizer, and direct the spray upon the back of the fauces while the patient is respiring deeply. The application should be made every two or three hours, and, in patients of sufficient age to allow of its being thorough, is often very serviceable.

CARBONATE OF CALCIUM ($\text{CaOCO}_2 - \text{CO}_3\text{Ca}$).

This salt is officinal in four different forms.

CRETA.—*Chalk* is the native, friable carbonate of calcium, a milk-white, soft solid, of an insipid, earthy taste; insoluble in water; wholly soluble, with effervescence, in dilute muriatic acid.

CRETA PRÆPARATA.—*Prepared Chalk* is chalk freed from impurities by pulverization, levigation, and elutriation; a white, perfectly smooth powder.

CALCI CARBONAS PRÆCIPITATA.—The *Precipitated Carbonate of Calcium* is a white powder, free from grittiness, which is made by precipitating chloride of calcium with carbonate of soda.

TESTA PRÆPARATA.—*Prepared Oyster-shell* is the powdered shell of *Ostrea edulis*, or common oyster, washed with warm water, and afterwards treated as in making prepared chalk.

THERAPEUTICS.—Carbonate of calcium in its different forms is used internally as an antacid and very mild astringent. As none of the salts which it forms are purgative, it, with the other preparations of lime, is the best antacid when diarrhœa is present. The crude chalk should never be used, but the other preparations are probably of equal value. Some practitioners claim, however, that the oyster-shell is more acceptable to delicate stomachs, on account of the animal matter which it contains; and under the name of *Cassillon's Powder*—a mixture of salep, tragacanth, sago, ãã three parts, prepared oyster-shell one part, and cochineal q. s. to color it—has been much used in obstinate summer diarrhœas. A drachm of this is boiled in a pint of milk, and the decoction taken as food *ad libitum*. The dose of carbonate of calcium is twenty grains to a drachm. *Chalk Mixture* (*Mistura Cretæ*, U.S.P.) is generally preferred to the powder. It contains thirty grains of the chalk to the ounce, and is given in doses of one to two tablespoonfuls. It is often combined with laudanum or paregoric and tincture of kino or catechu.

Externally, prepared chalk and the precipitated carbonate are used as desiccants and protective applications to *ulcers* and *chronic burns*; also in *excessive sweating* of the feet, and in *intertrigo* and other affections of the skin.

CLASS II.—ANTHELMINTICS.

THESE are medicines which kill or cause the expulsion of intestinal worms. They are sometimes divided into *vermicides*, those which kill, and *vermifuges*, those which expel; but there is little or no practical use in the division. It is of much greater importance to establish the relations between these drugs and the different species of entozoa, since clinical experience has demonstrated that an anthelmintic very efficient against one form of intestinal worm may be not injurious to another species. Therapeutically considered, the entozoa may be divided into the *Tapeworms* (*Tæniæ*), *Round-worms* (*Lumbrici*), and *Seat-worms* (*Ascarides*). The last of these differ from the others in that they are to be attacked solely by enemata.

It is obvious that the value of an anthelmintic depends not only upon its power of poisoning the articulate, but also upon its harmlessness as regards the patient. Thus, it is the eminent combination of these qualities which renders the infusion of quassia so valuable in cases of seat-worms, whilst carbolic acid, though very efficient, should never be used against the same parasite, since it has greatly imperilled, if it has not destroyed, the life of the patient when so employed.

There are certain general rules which govern the administration of anthelmintics, and which should not be lost sight of. They may be summed up as follows: Let the alimentary canal be as empty as possible, so that the drug may act with the greatest force upon the enemy. For this reason, anthelmintics are best administered early in the morning; and in obstinate cases the patient should be required to fast until dinner-time.

If the drug be not itself a purgative, from four to eight hours after its administration a brisk cathartic should be administered; or a purgative dose of calomel may be combined with it, as the bilious purging induced by the latter drug seems to be especially obnoxious to the entozoa.

SPIGELIA—PINKROOT.

The root of *Spigelia Marilandica*, a herbaceous perennial, growing in the Southern and Southwestern United States. It consists of a knotty head, with numerous fine, crooked, branching rootlets. The odor is faint and peculiar; the taste sweetish and slightly bitter. It contains, according

to the analyses of M. Feneulle and of R. H. Stablen, tannic acid, fixed and volatile oils, resin, and a bitter uncrystallizable body; but exactly upon what its virtues depend has not been determined.

PHYSIOLOGICAL ACTION.—Although there is sufficient testimony to show that pinkroot possesses decided narcotic powers, yet its action has been scarcely at all investigated. The observations of Drs. Hodge Thompson (*An Inaugural Dissertation*, quoted by Eberle), Eberle (*A Treatise on Materia Medica and Therapeutics*, i.), and Spalsburg (*Boston Medical and Surgical Journal*, April, 1855) have shown that in overdose it causes symptoms similar to those produced by one of the mydriatics. They are acceleration of the pulse, dilatation of the pupils, heat and dryness of the skin, flushing and a swollen appearance of the face, with, in Eberle's cases, talkative delirium. Two fatal cases* of poisoning by it are said to have been recorded by a Dr. Chalmers.

THERAPEUTICS.—*Spigelia* is a most efficient remedy in cases of the round-worm, and is, when given within the bounds of moderation, entirely safe. It appears to narcotize the worm, and requires the use of a brisk cathartic.

ADMINISTRATION.—One of the fluid extracts affords the best mode of administration.

Infusum Spigeliæ—*Infusion of Spigelia* (℥ss to Oj). Dose, f℥iv, repeated morning and evening.

Extractum Spigeliæ Fluidum—*Fluid Extract of Spigelia*. Dose, f℥ss.

Extractum Spigeliæ et Sennæ Fluidum—*Fluid Extract of Spigelia and Senna*.—A very valuable preparation, liked by children. Dose, for an adult, f℥ss; for a child two years old, f℥ss to f℥i, repeated every four hours until it purges.

AZEDARACH, the bark of the root of *Melia Azedarach*, or *Pride of China*, is used in the South as a remedy for the round-worm. It is said to possess poisonous properties similar to those of *spigelia*, yet it is affirmed that animals and children eat its fruit with impunity. It is usually given in decoction (℥ii to Ojss, boiled to a pint), the dose being for a child a tablespoonful every two or three hours, until the bowels are affected.

CHENOPODIUM—WORMSEED.

The fruit of *Chenopodium anthelminticum*, or *Jerusalem Oak*, a rank, odorous plant, growing about waste places in the suburbs of towns in the United States. It consists of minute, globular, light-brown seeds, about the size of a pin's head, of a nauseous odor and pungent taste, due to the volatile oil which they contain in large quantity.

Wormseed Oil (*Oleum Chenopodii*, U.S.P.) is of a light-yellow color,

* These cases appear to have been simply copied from book to book indefinitely. I have failed to find any reference to the original publication, and their authenticity remains doubtful.

becoming darker and less fluid by age, of a peculiar powerful odor and a hot burning taste.

THERAPEUTICS.—The oil of wormseed has been used in *hysteria*, but is now employed only as an anthelmintic against the lumbricus, and more rarely the tapeworm. It is very efficient, and ten drops of it on sugar may be given to a child three years old, before breakfast, dinner, and supper, for two days, followed by a brisk purge.

BRAYERA—KOOSSO.

The dried flowers and unripe fruit of *Brayera anthelmintica*, a tree, native of Abyssinia. They occur in compressed greenish-yellow clusters, of a fragrant balsamic odor and a taste which in a little while is acrid and disagreeable. Koosso contains a volatile oil, tannic acid, and a resin, discovered by Pavesi and known as *Koossin* or *Tenin*. This is crystalline, white or yellowish, slightly soluble in water, freely so in alcohol, and was shown by Dr. Bedall (*Sydenham Year-Book*, 1868, p. 476) to be the active principle of the drug, which yields about three per cent. of it.

THERAPEUTICS.—*Brayera* is a most efficient remedy against the tapeworm, and even in large doses causes no greater inconvenience to the patient than some nausea, abdominal pain, and looseness of the bowels. It is generally not necessary to administer any purgative with it, and the worm is discharged dead with the last watery passages. A half-ounce of the powdered flowers is given suspended in water in the morning, with the usual precautions as to diet, or from twenty to forty grains of koossin, wrapped up in a wafer, may be substituted for the crude drug. Care should be exercised in administering it to pregnant women, as it is stated to have produced abortion.

SANTONICA—SANTONICA.

Levant Wormseed consists of the unexpanded flowers and peduncles of *Artemisia Contra*, a composite of Asia Minor and other parts of the East. It consists of pale, greenish-brown, smooth heads of four or five tubular flowers, of very strong aromatic odor when rubbed, and a bitter disagreeable taste. It contains volatile oil, resinous matter, and a crystalline principle, *Santonin*, upon which its virtues depend.

SANTONINUM.—*Santonin*, or *Santoninic Acid*, occurs in colorless, pearly, four-sided, orthorhombic tables, soluble in from four thousand to five thousand parts of cold and two hundred and fifty parts of boiling water, freely soluble in alcohol and chloroform, moderately so in cold ether; insoluble, or nearly so, in glycerine. It has a neutral reaction, but unites with alkalies to form salts, and hence is freely soluble in alkaline solutions. When slowly heated, it sublimes, unchanged, at from 165° to 175° C. When rapidly heated, it is converted into a brownish-red oil, which becomes carmine-red upon the addition of caustic potash.

On exposure to sunlight, or, more slowly, even in the ordinary daylight, the colorless crystals of santonin acquire a golden-yellow tint. If this change is a chemical and not a mechanical one, the alteration must be very slight, since, according to Krauss, the yellow crystals conduct themselves in their chemical relations precisely as do the colorless crystals, and are precipitated, by the addition of acid to their alkaline solutions, as colorless crystals.

PHYSIOLOGICAL ACTION.—Santonin is said to have been first introduced into medical practice in 1830, by Dr. Alms, by whom and Dr. Kahler it was simultaneously discovered. It is at present used almost solely on account of its poisonous action upon entozoa, but certainly has a very great influence upon man and the higher animals. When it is given to dogs and rabbits in large doses it causes accelerated breathing, slowing of the pulse, universal trembling, cramps, free salivation, unconsciousness, convulsions, dilated pupils, and death. (See experiments of Manns, *Das Santonin*, Marburg, 1851; also of Rose, *Virchow's Archiv*, Bd. xvi., 1859; also of Theodore Krauss, *Ueber die Wirkungen des Santonins und Santonin-Natrons*, Inaug. Diss., Tübingen, 1869.) After death the lesions are not absolutely constant, but hyperæmia of the nerve-centres and congestion of the lungs and heart are nearly always present. According to Rose, dogs will recover after doses of from thirty to sixty grains, although five to six grains will produce very decided symptoms; and according to Krauss, thirty grains are required to kill a rabbit, even when they are dissolved in chloroform and given subcutaneously.

The symptoms caused by large doses of santonin in man are closely similar to those which it produces in the lower animals. There are several fatal cases of poisoning by it on record. Dr. Franceschi Giovanni (*Bulletin Thérapeutique*, t. lxxiv. p. 362) has seen a child, six or seven years old, killed by six grains of santonin, after suffering from hæmaturia;* and Dr. Ambroix is said to have recorded analogous cases. The only fatal case that I have met with in which a detailed account of the symptoms has been given is that of Dr. Grimm (*Schweizer Zeitschrift für Medicin, Chirurgie und Geburtshilfe*, 1852, p. 493). In this, a rather feeble child, five years old, took two one-grain doses of santonin, and was seized with convulsive tremblings, which increased in severity until they became severe convulsions, which were accompanied by unconsciousness, trismus, pallor of the face, cold sweats, dilated pupils, and rapid pulse and respiration. Thirteen or fourteen hours after the ingestion of the poison, whilst the patient lay on her back quiet, unconscious, with moderately dilated pupils and a slow, feeble pulse, death occurred suddenly. No post-mortem was allowed. The santonin is said to have been chemically pure and to have been used in other cases with its usual results.

Quite a number of very serious, but not fatal, instances of poisoning have occurred with the drug, and the symptoms have been similar to those in the

* This is probably a mistaken observation, the urine being only blood-colored, and not containing blood (see p. 520).

case just detailed; great pallor of surface, with a blue color around the eyes or involving the whole countenance, has been generally an early symptom; vomiting has not rarely been present, and sometimes has been accompanied by colicky pains. Besides these manifestations, giddiness, mental apathy or stupor, great coldness of the surface, profuse sweating, tremblings, mydriasis, and finally loss of consciousness, with convulsions, often violent and accompanied by opisthotonos and emprosthotonos, are the usual phenomena of santonin-poisoning.

A very curious symptom caused by santonin, even when in doses which can scarcely be called toxic, is xanthopsia, or "yellow-seeing," as the Germans term it. It was, I believe, first noticed by Calloud, and has since been spoken of by almost every writer upon the drug. Usually it consists of a very deep yellow tint imparted to the landscape and to every object looked at, an effect perhaps most comparable to that of looking through yellow glass; sometimes this yellow is replaced by green; and Heydloff states that he has seen patients in whom the tint was red, and others in whom it was blue. Dr. Edm. Rose has published exceedingly elaborate papers upon this chromatopsia; but, as the matter is of interest rather to the physicist and student of optics than to the clinician, I content myself with a reference to his memoirs, which may be found in *Virchow's Archiv* (xvi. p. 233; xviii. p. 15; xix. p. 522; xx. p. 245; xxviii. p. 30).

Santonin is eliminated probably in a more or less altered condition by the kidneys, and thus gives rise to a very characteristic symptom or sign of poisoning by the drug which has not as yet been spoken of,—namely, discoloration of the urine. The new color is a very marked yellow, which has at first an orange tint, but after very large doses becomes saffron-like, or sometimes even a purplish red, which has given origin to the idea that blood was present. According to Manns, the addition of an alkali to the yellow urine causes it to become red.

The exact nature of the eliminated principle has not been determined: it is, however, probably the result of an oxidation of the santonin, as is believed by Kletzinsky, who asserts that the drug receives in the system six atoms of oxygen.

The results of the ingestion of large doses of santonin show that it has a very powerful action on the organism; but as to what portions of the latter are especially affected we have no information. Rose believes that the chromatopsia is due to a peculiar action of the drug upon the nerve-centres; but it seems to me more probable that it is simply the result of a very faint staining of the humors and other parts of the eye by the drug, and is analogous to the similar phenomenon sometimes seen in jaundice. Like very many other substances which escape through the kidneys, santonin increases the flow of urine, and, according to Dr. Farquharson (*British Medical Journal*, 1872), increases slightly the elimination of urea.

THERAPEUTICS.—Dr. D. Dyce Brown (*British and Foreign Medico-*

Chirurg. Review, April, 1871), having noticed that a blind man to whom he was giving santonin for worms recovered to an extraordinary degree his vision, has recommended it in cases of loss of optic nerve power, and Dr. Ogston has used it with more or less complete success in a number of cases. Although Dr. Brown had apparently no knowledge of it, yet many years before his experience M. Guépin and M. Martin recommended the drug in *amaurosis* (*Ann. de Thérapeut.*, 1862). M. Guépin believes it to be especially useful in amaurosis following acute choroiditis and iritis.

Whether the use of santonin in diseases of the eye will or will not ever amount to anything, cannot at present be told; but in the past the drug has been employed almost solely as an anthelmintic in persons troubled with the *lumbricoid* or *round worms*.

It is a parasiticide, killing and not expelling the worm, and consequently should, when used, be combined with or followed by a brisk cathartic. Especial advantage is said to be derived from the joint use of it and calomel.

TOXICOLOGY.—It has been denied that santonin is poisonous, and asserted that the evil results which have followed its use have been due to the mixture of strychnia with the drug. The only ground for this opinion is that in one case, through the carelessness of an apothecary, strychnia was mixed with santonin, or else was sold instead of santonin, and death resulted. Dr. Walz, in an extended examination (*Jahresbericht für Pract. Pharmacie*, Bd. xv.), found that the santonin of the shops is pure; and it is inconceivable how strychnia could be mixed with it except purposely or by the grossest carelessness. Moreover, the symptoms which have been produced in alleged poisoning by santonin are very different from those caused by strychnia, and are in close accord with those which santonin induces in the lower animals. Finally, in some of the reported cases, seven of which are collected by Krauss, the santonin was examined and found to be pure. Under these circumstances, it would be absurd longer to deny to the drug poisonous properties. As to the proper specific treatment of poisoning, we are very much in the dark, as we have little knowledge of the physiological action of the drug. The stomach and bowels should be evacuated, and symptoms met as they arise.

ADMINISTRATION.—Santonin is best administered in the form of lozenges, which, if the unbroken crystals are used, can be rendered very pleasant to the taste, so that children will eat them as candy.

The dose of santonin for an adult is two to four grains; for a child two years old, one-fourth to one-half grain. Very alarming symptoms have been caused by two one-grain doses exhibited within three hours in a child eight years old (Grimm); in a child two and a half years old, four grains apparently came very near causing death (Dr. Berg, *Württemberg. Medic. Correspondenzblatt*, 1862); and in the fatal case noted on page 519, only two grains were taken by a child five years old. For young infants, santonin is hardly a safe remedy in any efficient dose. When a dose of any size is given, it should not

be repeated in less than eight hours, and the last dose should be accompanied by a purgative amount of calomel.

The *santonate of soda* has been proposed instead of santonin, but as an anthelmintic it has no advantage over the latter: since the object is to get as much of the remedy as possible in contact with the worm, and as to do this a slow, not rapid, absorption is necessary, the insolubility of santonin is an advantage.

FILIX MAS.—*Male Fern* is the rhizome of *Aspidium filix mas*, or male fern of Europe. The rhizome, when perfect, is six to twelve inches long, and covered with large, brown, imbricated scales. Its taste is bitter and astringent. Its therapeutic properties appear to reside in an *oleoresin* (*Oleoresina Filicis*, U.S.P.), which, as extracted by means of ether, is a dark, thick liquid, of a bitterish, nauseous, slightly acid taste.

THERAPEUTICS.—Male fern is employed almost exclusively against the *tapeworm*. In its administration it is necessary to regard strictly the general rules applying with greater or less force to all the remedies of the class, and which are especially imperative when a drug is employed against the tapeworm. The patient should live upon milk and a little bread for one day, and the following morning take a full dose (℥ss to ℥i) of the extract, fasting, and repeating it in two or three hours. At noon the patient may resume the use of food, and in the evening a brisk cathartic should be given.

PEPO—*Pumpkin Seeds*.—The seeds of the ordinary pumpkin are a most valuable remedy in cases of *tapeworm*, perhaps even more efficient than the male fern, and perfectly harmless. Two ounces of the seeds may be beaten up with sugar into an electuary, or with water into an emulsion, and be taken fasting in the morning, the patient having dieted the previous day. Some hours after their administration, a brisk purge should be given.

TURPENTINE, in doses of half a fluidounce, has been used in cases both of *tapeworm* and of *round-worm*. It is efficient, but is liable to produce unpleasant effects, and should be employed only when other remedies have been used without success or are not to be had. It should be given in combination with twice its bulk of castor oil.

POMEGRANATE RIND.—The bark of the pomegranate root (*Granati Radicis Cortex*, U.S.P.) is an efficient but very unpalatable anthelmintic against the *tapeworm*. The decoction of the fresh root (℥ii to Oj) is to be preferred; a pint of it to be taken in three doses, an hour apart, before breakfast.

MUCUNA—*Cowhage*.—The sharp, rigid hairs of the pods of *Mucuna pruriens*, an East India plant, were formerly used as an anthelmintic in cases of the *round-worm*, and are still officinal, although now very rarely if ever used. They are believed to kill the worm mechanically,—i.e., by piercing it. The pods were dipped into molasses, the hairs scraped off, and a tablespoonful of

the thick mass was given to an adult—a teaspoonful to a child—morning and evening, for three days, after which a brisk purge was administered.

ROTTLERA—*Kameela*.—The hairs of the fruit of *Rottlera tinctoria*, a plant cultivated in India as a dye-stuff. It is said to be an efficient parasiticide in cases of *tapeworm*. It is an orange-red, very inflammable, granular powder, mixing with water with some difficulty, and containing traces of a volatile oil and coloring resinoids, to one of which Dr. Anderson has given the name of *Rottlerin*. *Kameela*, in full doses, is actively purgative, indeed drastic, and sometimes causes also nausea and vomiting. It imparts its virtues to alcohol, and hence may be exhibited in the form of tincture. The dose of the powder is from one to two drachms suspended in syrup, given in the morning, and repeated in eight or ten hours if it do not purge.

CLASS III.—DIGESTANTS.

IN this class are put a few remedies which are used to aid the stomach in dissolving the various articles of food. There are, I think, really but three articles worthy of a position in the class: namely, pepsin, hydrochloric acid, and lactic acid. Prof. Geo. B. Wood claims, however, a position in it for yeast. The theory upon which the latter drug is employed is that in *diabetes* there is an excessive formation of sugar in the stomach, and that the yeast, by causing a fermentative alteration of this into acetic acid, will do good. With our present knowledge of the pathology of the disorder, it is certain that the sugar is not formed in the *primæ viæ*, but in the liver, or at least in some internal viscus; and, this being so, it is evident that the theory above spoken of falls to the ground. Further clinical experience has, I think, demonstrated the uselessness of yeast in diabetes, and at present the remedy is very rarely, if ever, exhibited in that disorder.

PEPSINA—PEPSIN.

As every one knows, there is secreted by the gastric glands a peculiar albuminous body, which has the power not only of coagulating albumen, but also, with the aid of acidulated water, of redissolving it. To this principle the name of pepsin has long been given. A discussion of its nature and properties would be better in place in a work upon physiology than in one on therapeutics, and I shall therefore say nothing further about them.

The dried stomach of calves has been used since time immemorial for the purpose of coagulating milk, by housewives, with whom it is customary to place the dried viscus in wine and to call the liquid thus formed, as well as the prepared stomach, *rennet*. It is stated by Dr. Jas. Gray (*Edinburgh Medical Journal*, Jan. 1853) that rennet-wine should be of such strength that one teaspoonful of it will coagulate a pint of milk. Rennet is said to have been long used in England as a domestic remedy in dyspepsia (*Medical Times and Gazette*, April, 1857). In South America, the inner coat of the gizzard of the ostrich is stated to be put to a similar use (E. S. Wayne, *American Journal of Pharmacy*, March, 1868); and in our own country the dried gizzards of chickens and turkeys are no less famous among medically-inclined housewives.

Dr. Corvisart, of Paris, is asserted to have been the first to propose the use of the active principle of the stomach in feeble digestion; and of latter years the manufacture and consumption of pepsin have become very great.

Various processes have been suggested for the preparation of the drug, but none of them yield a pure proximate principle, if indeed pepsin have really such nature, and be not an albuminous body of varying constitution.

By most of the methods of manufacture the pepsin is obtained in the form of a viscid fluid; and to change this into a powder requires the addition of a large quantity of starch, so that the powdered pepsin, as sold, contains a considerable percentage of foreign material.

There are in the market two chief brands of this powdered pepsin, Boudault's and Hawley's, and the testimony seems to be that they are about equal in value. Certainly neither of them is pure pepsin; and for the sake of those physicians who have opportunity to prepare or to have prepared their own material, I add a simple method of manufacture, as given by Prof. Lionel Beale,—a plan which he states to be the result of a large number of experiments. It is as follows:

Dissect off carefully the mucous membrane of a *perfectly fresh* pig's stomach, and place it on a flat board; cleanse it lightly with a sponge and water, so as to remove the particles of food and much of the mucus. Scrape it hard with an ivory knife, so as to squeeze out all the contents from the glands. The viscid mucus thus obtained contains all the pepsin, with much epithelium. Spread it upon a piece of glass so as to form a very thin layer, which is to be dried at a temperature of 100° (no higher) F. over hot water, or in vacuo over sulphuric acid. When dry, scrape from the glass, powder in a mortar, and transfer to a well-stoppered bottle.

Dr. Tuson found this pepsin to be odorless, nearly free from the disagreeable taste of the ordinary commercial preparation, and twenty-five times as strong as the latter (*Medical Times and Gazette*, vol. ii., 1872).

Whatever form of pepsin be used, if good effects are to be obtained from it it must be given with acid, unless indeed there be reason to believe that this constituent of the gastric juice is not wanting. Alcohol destroys the digesting power of pepsin, and therefore wines are inferior preparations of it. The reactions of pepsin with organic and inorganic matters are very complex, and not well understood: consequently I think the physician should eschew all elixirs or compound preparations of the drug, using only the powdered pepsin or a freshly-prepared digestive solution. If other remedies are to be given, it is no great hardship to write a second prescription and exhibit them by themselves. In making the digesting fluid, water and muriatic acid, or glycerine, water, and muriatic acid, should alone be employed with the pepsin.

Prof. Beale recommends the following formula for pepsin prepared by his method:

Take of the powder, five grains; strong muriatic acid, eighteen drops; water,

six ounces. Macerate at a temperature of 100° for an hour; filter, so as to form a perfectly clear fluid.

THERAPEUTICS.—It is a question of some importance to decide how far pepsin is valuable and reliable as a medicine. It is evident that any influence for good which it possesses is dependent upon its solvent power, and that this, therefore, is a measure of its value. Taking as a type Boudault's pepsin, because it is the commercial article of highest reputation, it is claimed only that it will dissolve, with the aid of muriatic acid, four times its weight of fibrin at the temperature of the body, and that fifteen grains of it should effect the solution of a drachm and a half of boiled white of egg in twelve hours; and yet grave doctors prescribe for men pepsin as an artificial solvent in doses of ten grains. Evidently one of two things is certain: either the present practice is ridiculously absurd, or else pepsin acts upon the stomach itself in some way as a stimulant.

Clinically, pepsin has been used with asserted advantage in the *loss of digestive power* in adults, whether primary, or occurring in the course of other affections. Probably four-fifths of the drug which has been given has been inert either originally or from the method of its administration; and I do not doubt that in the great majority of cases the good that has been achieved has been due, not to the pepsin, but to the regulation of the diet and habits of the patient and to the drugs which have been exhibited along with the animal ferment. I have frequently seen pepsin given to adults, and have never yet seen it do any distinct good. At the same time, I do not mean to deny to it the possession of any virtue; but I do believe that its value has been vastly over-estimated, that it has been given to adults in ridiculously small doses, and that at least half a drachm of the commercial article should be given at a dose.

The testimony as to the value of pepsin in diseases of young children is very strong. To such it is generally given in doses nearly as large as those usually exhibited in the cases of adults. If we represent the absolute digesting power of ten grains of pepsin as x , it is evident that x represents a proportionately much greater power in the *primæ viæ* of a child than in those of an adult. The use of pepsin in children is therefore much more rational than in adults; and my own experience is in close accord with what seem to me the dictates of common sense. Pepsin is certainly much more efficient in children than in grown persons, and in the *chronic indigestion* and consequent *diarrhæa* of young children it may be tried with great hope of benefit. To a baby six months old five grains may be given in a little acidulated water after each feeding.

Feeding by the Rectum.—It is sometimes of the greatest consequence to feed a patient in other ways than by the stomach; and therefore the question of nourishing by the rectum is one of the gravest importance. The solvent influence of the rectal juices is practically nothing, and formerly the attempt was made to supplement this complete lack of power by the use of soups. It is evident that very little nutriment can be introduced into the system by such

a plan; and consequently feeding by the rectum has hitherto been only a "forlorn hope," and has accomplished little good. Dr. W. O. Leube (*Deutsches Archiv für Klin. Medicin*, March 15, 1872) has proposed a new plan, which bears upon its face the marks of efficiency, and which, as he has proven by experiments made upon dogs, is capable of yielding to the blood largely of nutritive materials. Moreover, he has by it maintained life for four weeks in a patient who had been poisoned with iodine and whose stomach rejected all food. The plan consists simply in digesting the food before injecting it into the rectum. The method is carried out as follows. The pancreas of swine or cattle is carefully cleaned of fat, and 50 to 100 grammes thereof cut into very small pieces. In like manner 150 to 300 grammes of beef are prepared. Both substances are then put into a dish with about 50 to 150 cc. of lukewarm water, and stirred into a thick paste, and drawn in a clyster-pipe with wide opening. In many cases from 25 to 50 grammes of fat may be added to the mixture, also at times some starch. An hour before using this clyster, one of pure water should be administered, to clean out the intestines.

In very warm weather there is some trouble in obtaining and keeping sweet the pancreas. This difficulty may easily be avoided by making a glycerine extract, which is said to be quite equal in digestive power to the fresh pancreas, and will remain good for several weeks. The following is the manner of its preparation. The pancreas of a bullock (which is sufficient for three enemata) is finely chopped and rubbed with 250 grammes of glycerine; and to each third of this, when about to be used, are added from 120 to 150 grammes of finely-divided meat. It is important that this mass should be injected into the intestine as soon as it is made; for if it is allowed to stand, the meat swells and the operation is thereby rendered difficult.

The experiments of Dr. L. S. Joynes (*The Richmond and Louisville Medical Journal*, March, 1869) have shown that the stomach of the pig, placed in water acidulated with hydrochloric acid, will not only rapidly dissolve itself but also small pieces of beef. When life is to be maintained by rectal feeding, probably a preparation made in this way could with advantage be alternated with the solutions of Leube.

CLASS IV.—ABSORBENTS.

THIS class contains remedies which are used for the purpose of absorbing acrid and deleterious materials, such as offensive discharges on the exterior of the body, and acrid secretions, or the irritant products of the partial decomposition of food, in the alimentary canal. For the first purpose very fine dry earth and plaster of Paris are used to some extent in practice; but, as their employment is purely in the province of the surgeon, I shall say no more about them here.

CHARCOAL.

Charcoal is officinal in the U. S. Pharmacopœia in two forms:

CARBO LIGNI.—*Charcoal* prepared from wood.

CARBO ANIMALIS.—*Animal Charcoal*, prepared from bone.

Charcoal for medical purposes should be made out of a light porous wood, and that prepared from the young shoots of the willow or of the poplar is almost exclusively employed. It is a black, brittle substance, and should have more or less lustre. It has a very remarkable power of absorbing many times its own bulk of gases, and, when exposed to the air, increases rapidly in weight. It should therefore, when intended for medicinal purposes, be powdered so soon as it is burnt, and put in small, completely-filled, closely-sealed bottles.

Animal charcoal, or *bone-black*, formed as it is by the partial burning of bones, contains a large percentage of the phosphate and the carbonate of calcium. Although this does not interfere with its strictly medical employment, it does with many of its uses in pharmacy; and consequently the U. S. Pharmacopœia directs a *Purified Animal Charcoal* (*Carbo Animalis Purificatus*) to be prepared by digesting ordinary animal charcoal in dilute muriatic acid, by which all the lime salts are dissolved out, washing thoroughly with water, and heating to redness.

Bone-black possesses absorbing powers far greater than those of ordinary or wood charcoal, and takes up coloring-matters, alkaloids, and other substances from these solutions: it is therefore very extensively used in pharmacy, as well as in the refining of sugar, and in other processes of the arts.

THERAPEUTICS.—Charcoal is used externally as an absorbent and disinfectant dressing to *foul wounds* and *ulcers*. A *charcoal poultice* may be

made by adding one to three drachms (according to size) of powdered charcoal to a flaxseed poultice; or, as the British Pharmacopœia directs, two ounces of bread may be well mixed with ten ounces of warm water, an ounce and a half of flaxseed meal be thoroughly stirred in, and to the cataplasm thus formed two drachms of powdered charcoal be added, and one drachm be sprinkled on the surface.

Internally, charcoal is employed as an absorbent in those cases of acute or chronic *indigestion* in which there are offensive liquids and gases in the alimentary canal, giving rise to local symptoms and foul discharges: in this way it is sometimes employed in *dyspepsia*, *cardialgia*, and similar disorders. It is at least conceivable that charcoal, given very freely, should accumulate in the alimentary canal, and mechanically cause serious trouble: for this reason, its habitual employment is very generally combined with that of laxatives.

On account of its power of absorbing alkaloids, *purified animal charcoal* has been recommended in poisoning by substances of that class. Its action is, however, too slow for much good to be expected from its use. If given at all, it should be exhibited very freely, since at least half an ounce of it is said to be required for the absorption of one grain of the poison.

ADMINISTRATION.—The ordinary dose of charcoal is from half an ounce to one ounce. Except in a mechanical way, it is perfectly innocuous in any dose.

CLASS V.—DISINFECTANTS.

DISINFECTANTS are substances employed for the purpose of destroying noxious miasmata or effluvia. It is evident that the consideration of these materials belongs rather to the province of hygiene than to that of therapeutics, since their employment is hygienic rather than medicinal, preventive rather than curative. The importance of disinfection is, however, so great, and its relation to clinical medicine is so close, that a work like the present ought certainly to treat of it, although perhaps more briefly than of other matters.

All poisonous exhalations may be divided into those which are generated within and those which are generated without the body. To the former I shall apply the term *contagions*, to the latter the term *miasmata*.

Of the nature of these poisonous influences we have no positive knowledge. It is most probable that contagions are of similar character throughout, and that they differ in their nature from miasmata. They are undoubtedly formed within the body, and are probably connected with living matter. Their power of self-propagation—*i.e.*, their power of multiplying and increasing when taken into the system—seems to me irreconcilable with the idea that they are definite organic compounds, such as acids, alkaloids, salts, or neutral principles. Morphia, the poison of snake-bites, and other known lethal organic compounds, do not in any instance increase in the body of their victim. Of course, space is wanting here to discuss at length the nature of contagions; and, having given what seems to me at least very strong presumptive evidence that these matters are living, I must content myself with stating my belief that not only is proof wanting that they are distinct living entities, but that it is also improbable that they are of such nature. The reader will find my reasons for this opinion in *The American Journal of the Medical Sciences*, October, 1868, p. 333. It does seem to me that the present evidence upon the subject tends to show that contagions are minute particles of living germinal matter, capable of imparting certain properties to the germinal matter of animal systems similar to those from which they sprang, and thereby propagating themselves in these systems, at the same time producing general constitutional disturbances. The important practical deduction to be drawn from these conclusions is that contagions are to be destroyed by substances poisonous to germinal matter or protoplasm in its more imperfect

form of development, and that the best criterion we have at present of the power of any material over them is its action on the low forms of life.

It is evident that contagion is in no wise necessarily connected with bad odors, that its most deadly form may saturate the air of an apparently clean and sweet chamber, and that the most stinking cesspool may be completely free from it.

A question of very grave importance naturally arises here, as to the exact province of disinfection when applied to contagions; or, in other words, Is it possible to destroy them in the air? It is a law which holds everywhere, that the more imperfectly developed animal organisms are, the more tenacious are they of life; and it seems to me also a general law that highly specialized germinal matter is more readily destroyed than that of lower life-grade. Be this as it may, I think that experience demonstrates that all those substances actively poisonous to contagions are still more poisonous to the germinal matter of human beings, and that it may be laid down as a general proposition that it is impossible to destroy the contagion-germs in the confined air of a room while men or women remain in the apartment. When it comes to the general atmosphere, to attempt to disinfect it, to essay the destruction of a wide-spread poison, such as, for example, exists in a smallpox epidemic, is simply so childish as to need no discussion to reveal its utter futility. It is well known that contagion diluted beyond a certain point is powerless, and the admission of large quantities of *fresh* or *pure* air, or, in other words, *free ventilation*, is the only means at our command of disinfecting rooms in which there are human beings. In ventilating an apartment, it must always be remembered that it is not merely air, but pure air, that is required. To ventilate one room into another, as is sometimes done, is foolishness; to open a window with the wind blowing over a miasmatic swamp or an open sewer, may be but to invite disease. The opportunity to destroy disease-germs is to be found not after they have been dispersed into the air, but whilst they are still in connection with the various solids and fluids passed from the body or brought in contact with it. In such diseases as cholera, in which the poison-germs are probably eliminated in large quantities with the passages, it is of the most vital importance to act on the stools so soon as they leave the body; and the disinfectant should be placed in the receptacle *before*, not after, it is used. To allow any passage to exist for a moment undestroyed, or at least undeveloped, is most culpable neglect. In the same way it is proper to disinfect all the discharges and all the substances which are exposed to the exhalations from a patient suffering from a contagious disorder.

The nature of miasmata is not known at all. Over many of them, malaria for example, we have no control. Those which we are able to influence are almost all the results of animal or of vegetable decomposition, either alone or conjoined. After these poisons have been produced and diffused through the air, they are probably in great part beyond our reach. The only thing that can be done is to dilute them precisely as in regard to contagions; and hence

free ventilation is the only reliable disinfectant under these circumstances. Sulphuretted hydrogen gas, although hardly worthy of a place among miasmata, certainly exerts a deleterious influence upon the system, and is therefore to be destroyed when in the air. There are probably other allied gases which can be attacked in the air of an apartment. It must be borne in mind that the poison of cesspools, of close rooms, and of similar localities is not sulphuretted hydrogen, but some unknown compounds; that there is no reason for believing that these are destroyed with the gas; and that, although the air of an apartment may be rendered inodorous, it yet may be saturated with poison. The stinking gases are probably of use as indicators of the presence of more subtle poisons. Wide-open windows, great draughts of fresh air, are the only proper disinfectants for a close room or a noisome hospital-ward. The attempt so often indulged in to purify such apartments by little saucers of chlorinated lime or of carbolic acid would be exceedingly ludicrous if it were not for the frightful results of the false security engendered.

The gases which arise from cesspools and similar depositories no doubt act as carriers of the peculiar poisons produced in these places, and it may be that even after their dispersion destruction of the gas causes a deposition of the organic poison. This is, however, merely speculation: it is not proven. Moreover, it is very doubtful whether the deposition of a poison in an apartment is an advantage. It is certainly illogical to draw any reasons in favor of disinfection instead of ventilation from the idea that the gases when destroyed will drop the organic matters which they have floated into the air.

It is otherwise if the gases are destroyed at their places of emanation. Thus, if a sewer be belching forth deleterious gases and poison, chlorinated lime or other materials slowly generating disinfecting gases at the mouth, or in the sewer, may really be of service by destroying the noxious vapors before they find their way out and carry with them into the air the poisonous effluvia.

Although miasmata cannot be readily destroyed when once generated, yet in very many instances we have power to prevent their formation. In doing this the chief factor is *cleanliness*, and the best disinfectant is *water*. Water acts chiefly in two ways: first, by dilution; secondly, by destroying organic substances. In order for decomposition to produce serious poisoning, the mass of material must be considerable. If the matter is diffused through a large bulk of water, and this spread over a considerable surface, it is evident that the effect of dispersion is obtained. Further, when water containing organic matter is allowed to run away in its natural channels, oxidation and destruction of the impurities result. Without entering further into the subject, it is sufficiently evident that *air and water are the great disinfectants*, and that the most skilful use of chemical substances cannot take the place of ventilation and cleanliness.


Various classifications have been made of the substances used as disinfectants. Without laying any stress upon the matter, without claiming any scientific accuracy or value for the arrangement, I shall consider disinfectants

under the headings of—Oxidizing Agents; Desulphurating Substances; Antiseptic Materials; Absorbing Substances.

OXIDIZING DISINFECTANTS.

Probably the only absolutely trustworthy disinfectant would be one capable of completely oxidizing and destroying organic matter. Fire is of this character, and when ruthlessly applied certainly is absolutely efficacious. It is evident that the use of this disinfectant must be very limited.

POTASSII PERMANGANAS—PERMANGANATE OF POTASSIUM.



This salt is prepared by heating together ten parts of binocide of manganese and twelve of potassa. It occurs in slender, prismatic crystals of a dark-purple color, inodorous, of a sweetish, disagreeable taste, and very soluble in water, with which they form a solution varying from a purplish black to a beautiful reddish lilac, according to the strength. When kept dry and not exposed to the atmosphere, the permanganate of potash is a permanent salt, but whenever in solution it is brought into contact with an organic body it at once gives up its oxygen to the latter, and is converted into potassa and the black oxide of manganese.

The disinfectant power of this salt is beyond question; but at the same time this power is very limited, as the remedy acts only by yielding up its own oxygen. The salt is comparatively dear, and therefore cannot be used at all to disinfect large masses of materials. It is at present almost solely employed as an addition to water with which *ulcers*, *abscesses*, etc., are washed. In dilute solution its local influence is that of very mild stimulation, and is always beneficial. Whether permanganate of potash has the power of destroying specific poisonous compounds has not, that I am aware of, been determined: probably, however, it does affect the germinal matter of contagion, and it certainly arrests decomposition and destroys its products when used upon a wound. When employed in the form of powder, it even affects living tissues, acting as a mild caustic, and, as such, may often be applied to *sloughing ulcers* with advantage. In *ozæna*, in fetid *leucorrhæa*, *otorrhæa*, and similar affections, the permanganate makes a very useful wash, the strength of which may vary, according to circumstances, from one to twenty grains to the ounce. Internally this salt has been commended in various diseases, in doses of one or two grains. As it is evident that, even if many grains were taken into the stomach, immediate complete decomposition of them would there occur, the absurdity of this use of the drug needs only to be pointed out. There is an official *Liquor* (*Liquor Potassii Permanganatis*, grs. lxxiv to Oj), which has no advantages over extemporaneous preparations.

CHLORINUM—CHLORINE. (Cl.)

Chlorine gas is official only in the form of the *Aqua Chlorini*, which is prepared by heating together the black oxide of manganese and muriatic acid, and allowing the chlorine which is generated to pass through water until the latter is saturated. *Chlorine Water* is a greenish-yellow liquid, of a very sharp taste, and having a strong odor of chlorine.

When chlorine is brought into contact with organic substances and moisture, it unites with the hydrogen of the water and liberates nascent oxygen, which rapidly oxidizes and destroys the organic compound. When chlorine comes in contact with sulphuretted hydrogen, it removes its hydrogen and thereby destroys it. Again, chlorine in sufficiently concentrated solution is certainly fatal to germinal matter. From these facts it is readily perceived that chlorine is one of the most efficient of the disinfectants, destroying the compounds produced by decomposition, arresting putrefaction by destroying the organic bodies, and finally, if in sufficient concentration, killing the disease-germs.

The great drawback to the use of chlorine is the limited power any given quantity of it possesses. It is evident that after it has once oxidized organic matter, and been itself converted into muriatic acid, its disinfectant power has been in great measure lost, and it can exert no further influence. Further, its corrosive properties render it unfit for the purification of clothing, of fomites, and various other textile fabrics, since there is no good reason for believing that it is effective as a disinfectant unless in sufficient concentration to injure the bodies spoken of.

When it is desired to liberate chlorine in the air of a room, eighteen parts of finely-ground common salt with fifteen parts of finely-powdered black oxide of manganese should be introduced into a flask; then there should be added forty-five parts of concentrated sulphuric acid and twenty-one of water, previously mixed and *completely cooled*; and lastly, the flask is to be well shaken. When the evolution of gas ceases, it may be renewed by placing the flask in warm water. Of course, chlorine in the atmosphere can have no effect upon disease-germs unless in such quantity as to be completely inimical to human life. As its generation is less manageable than that of sulphurous acid, and as it is less efficient than the latter agent, the sulphurous acid is preferable whenever it is desired to disinfect a closed apartment.

Internally, chlorine water has been used in various diseases, especially in malignant *typhus*, but at present is very rarely if ever employed. It is stated to be stimulant and tonic to the stomach, and is thought by some to have an especial influence upon the liver. It has been employed in *chronic hepatic affections*; the dose is half a fluidrachm to two fluidrachms in three or four fluidounces of water.

Chlorine water is a powerful irritant, capable of producing severe inflammation of the skin or toxic *gastro-enteritis*. Properly diluted, it forms an

excellent stimulant, disinfectant, detergent wash for *foul ulcers*, and may also be used as a gargle in *malignant sore throat*.

CALX CHLORINATA—CHLORINATED LIME.

This is a grayish-white substance, occurring in powder or friable lumps, having a hot, acrid, astringent taste, and an odor resembling that of chlorine. It is made by the action of chlorine upon hydrate of calcium or slaked lime, and should contain at least twenty-five per cent. of chlorine. It probably varies in its chemical constitution, but, according to the most recent views, is chiefly composed of the hypochlorite and the chloride of calcium. When exposed to the air it slowly evolves hypochlorous acid, which, being an unstable compound, undergoes spontaneous decomposition, and finally sets free fourteen-fifteenths of its chlorine. When an acid is added to chlorinated lime, the chlorine gas is rapidly evolved. If a specimen of bleaching-powder be very moist, it generally contains an over-proportion of the deliquescent chloride of calcium, and is correspondingly unable to liberate chlorine.

LIQUOR SODÆ CHLORINATÆ.—*Solution of Chlorinated Soda*, or *Labarraque's Solution*, is an official preparation, made by triturating chlorinated lime with a solution of the carbonate of sodium. It is a greenish-yellow liquid, having a slight odor of chlorine and a sharp saline taste. It contains, among other substances, hypochlorite of sodium, and possesses the therapeutic and disinfectant properties of the chlorinated compound. Owing to its liquid form and comparative freedom from odor, it is the most elegant of all the preparations of its class for use in the sick-room. Properly diluted, Labarraque's solution may be employed for all the therapeutic purposes that chlorine water is used for. The dose is half a fluidrachm to two fluidrachms, in half a tumblerful to a tumblerful of water.

Iodine and *Bromine* both are capable of acting as disinfectants, by dehydrogenating water and liberating nascent oxygen. They are less readily applied than chlorine, and are very rarely used, especially as they are very costly substances.

DESULPHURATING DISINFECTANTS.

These are various metallic salts which are believed to act as disinfectants by uniting with the sulphur of sulphuretted gases and precipitating as sulphurets. As examples of such may be mentioned sulphate of zinc and nitrate of lead. Under the name of *Ledoyen's Disinfectant Solution*, a solution of the latter salt has been and still is used to a considerable extent as a disinfectant. Although it certainly destroys sulphuretted hydrogen with great rapidity, it does not seem to me a good material for the purposes to which it is applied. The reasons of this are, first, it has no action besides

that of a desulphurating body; second, it is a comparatively dear salt; third, it forms an intensely black precipitate, discoloring everything with which it comes in contact.

It is evident that chlorine precipitates sulphur; but it is not really a desulphurating compound, since it takes the hydrogen from the sulphur, and not the sulphur from the hydrogen. It is an oxidizer, and as such has been already considered.

It is very certain that at least some of the so-called desulphurating compounds act also in other ways, and it is probable that many of them are efficient oxidizers.

The power which the oxides of iron have of converting ordinary oxygen into ozone has already been spoken of (p. 78). This action is a slow, persistent one, and the oxidation which results is equally slow and persistent. As already stated, organic matter, if diffused through water and exposed to the air, is gradually destroyed by oxidation. It has been found that when water which is loaded with the products of decomposition is exposed to the action of iron plates, or even of iron pipes, this destruction of organic impurities is greatly hastened. According to Mr. G. Michaelis (*The Philadelphia Medical Times*, vol. iii. p. 621), even the most filthy water, under the influence simply of iron plates and the air, will become perfectly pure in forty-eight hours. The action is evidently one of oxidation, but is in its details complicated. According to the researches of Dr. Mankiwich, iron possesses the property of converting ammonia into nitric acid, and also facilitates, or even provokes, such decomposition in the organic matter as shall cause ammonia to be formed. The nitric acid thus generated is one of the most powerful oxidizing substances known, and as fast as formed attacks the organic matter. It is evident that in this process the iron acts as an intermediate agent between the air and the decomposing matter; that, unlike chlorine or permanganate of potassium, it does not itself undergo a conversion equivalent in chemical relation to the oxidation, and consequently that its power is not so limited as that of those compounds. The iron, however, undoubtedly suffers to some extent, and is largely oxidized; but the oxide formed has certainly the power of generating ozone, and very probably to as great an extent as the original metal. How soon the power of oxidation is finally lost, and what eventually becomes of the iron, in the presence of an overwhelming mass of organic matter, has not, that I am aware of, been determined.

FERRI SULPHAS.—*Sulphate of Iron*, or, as it is known in the impure form in which it is used, *Copperas*, is, for the purpose of averting the results of decomposition, probably the best disinfectant we have. Of course, in the choice of a disinfectant for use in cesspools, sewers, and similar positions, it is not so much the power in proportion to weight as to cost that is the important matter; and the excessive cheapness of copperas is combined with very great efficiency. I am informed that, by the quantity, the impure form,

the efficiency of which almost equals that of the pure drug, can be bought for about a cent a pound. The changes wrought by sulphate of iron thrown into a mass of decomposing matter are very complex and not absolutely known. In the first place, if sulphuretted hydrogen and ammonia be present, a sulphuret of iron is at once precipitated; again, a part of the salt is decomposed by the ammonia, and probably also by the stinking alkaloids which exist in the decomposing mass of a cesspool; further, by oxidation, in all probability, some of the protoxide of iron is changed into the sesquioxide, by which the ozonizing power of an iron compound is doubtless exerted long after the destruction of the original salt. The sulphuric acid set free from the iron very probably contributes to a feeble extent in the destructive reactions which are set up.

From what has been already stated, it is evident that the chief use of the copperas is in altering the course of putrefaction and in destroying its products. When a discharge which contains a disease-germ is to be disinfected, when living particles are to be killed, the sulphate of iron, to be efficient, must be in great excess. Indeed, proof of the completeness of its action in this respect is wanting, and consequently, when disease-germs are to be killed, one of the antiseptics should be added to it.

The exact nature of the changes which copperas produces when added to masses of decaying matter has not been fully determined, but practical use has certainly established its efficiency as a disinfectant. Without going more deeply into the subject, I shall quote, as an illustration rather than proof of this, the experiments of Albert Eckstein, who published (*Zeitschrift des Oester. Apotheker-Vereines*, Feb. 10, 1873) an account of his attempts to disinfect a privy which was used daily by one hundred persons.

The experiments are as follows:

1. Two pounds of the sulphate of iron in solution were used. After from two to three hours all bad smell had disappeared, but in twelve hours all the influence of the disinfectant was lost.

2. The sulphate of copper was employed in solution; result the same.

3. Two pounds of the sulphate of iron in crystals were thrown in; their effects lasted two days.

4. The sulphate of copper, the same.

5. Sulphurous acid in solution rapidly lost its effect, and was exceedingly irritating to the respiratory organs.

6. Two pounds of impure carbolic acid filled the house for two days with such a disagreeable smell that it was impossible to tell whether the original odor was destroyed or covered up.

7. Two pounds of sulphate of iron in a parchment sack exerted a disinfecting influence for three full days, and when the parchment sack was drawn up it contained only some dirty, odorless fluid.

8. Two pounds of the best chloride of calcium in the parchment sack disinfected the privy for at least nine days.

If a rapid effect is desired, or a mass of solid material is to be acted on, the copperas should be in solution: if a more persistent action is wanted, or if the mass is liquid, the method employed by Eckstein, of using a parchment sack, may be resorted to, or the salt in powder may be scattered over the surface of the material to be disinfected.

LIME is probably the oldest of all the disinfectants, and in some respects is efficient, although on the whole its general use is to be reprobated. It does not act merely as a desulphurating and coagulating agent: like the other strong alkalies, it causes catalytically a slow oxidation of organic matter. Thus, ozone of itself will not oxidize olein, but if potash be added to the mixture the reaction commences at once; and it is notorious that a compost-heap to which lime has been added rots—*i.e.*, oxidizes—much more rapidly than one which has none of the alkaline earth in it. This action of lime is, however, too slow for ordinary purposes: moreover, there are very serious objections to the use of lime as a disinfectant.

The poisonous principles contained in sewage, etc., whatever their nature may be, are probably volatile, and lime, acting as a strong base, sets free large quantities of ammonia in animal matter undergoing decomposition. It may be that volatile poisons, alkaloids, are liberated with the ammonia; but, whether they are or not, it is a well-known physical fact that a volatile substance in escaping carries off with it even non-volatile materials, and facilitates to a still greater degree the escape of principles only less volatile than itself.

Lime is, for the above reasons, not available for use as a disinfectant in cesspools and sewers unless it is added in large quantities day by day from the beginning, so as to keep the collection under its influence, and unless some absorbent is added with it to take up volatile principles. In the ordinary open privies of the country, a shovelful of a mixture of lime and plaster of Paris, or even of lime and earth, thrown in day by day, will at once tend to prevent odor, and at the same time prepare the contents for use as manure.

When spread upon walls in the form of whitewash, lime may act to some extent as an oxidizer; but probably its chief influence is as an absorbent, which takes up the deleterious emanations. A very striking example or illustration of this action of whitewash occurred some years since in the New York city hospital. A ward which stood isolated from the remainder of the institution had been used for the reception of cases of typhus fever from the shipping of the port. It was finally abandoned and allowed to stand unoccupied, with its windows wide open, for several months. At the end of this time, a gang of men were set to scraping the whitewash off the walls. Of these workmen a majority were seized with the ship-fever, and several died.

The *chloride of calcium*, which proved so efficient in the experiments of Eckstein, detailed on page 537, was brought into notice in 1872 by E. C. C. Stanford, Esq., of England, whose elaborate series of experiments seemingly proved that of all the chlorides it is the most powerful for its price, and is

decidedly superior to "*chloralum*." He recommends it in the form of solution containing twenty-five per cent. of the solid salt and acidified with twelve per cent. of hydrochloric acid. This increases its power, and is a harmless addition. As it does not make a black precipitate with sulphuretted hydrogen, the chloride of calcium, if it be equally effective, is better than copperas for ordinary household use. The solution of Mr. Stanford is said to be the best form of it for many purposes; but for others the solid form is preferred. Thus, there is always great difficulty in keeping urinals in houses free from offensive odor. This is said to be entirely obviated by putting a lump of chloride of calcium in the urinal; it lasts a long time, as it dissolves very gradually, and keeps the urinal perfectly free from odor.

At present, at least in our markets, the chloride of calcium is offered only in a pure form, and at a price which excludes it from common use as a disinfectant; but enormous quantities of it sufficiently pure for disinfecting purposes are said to be thrown away, as a waste product, in all alkali-works: its utilization would be a great benefit.

Further experimentation and investigation are needed before the proper value of the chloride as a disinfectant can be determined; but certainly the matter is worthy the attention of the owners of those manufacturing laboratories in which it is a waste product.

ANTIZYMOTICS.

For many years it has been believed that fermentation of every sort, whether alcoholic, acetic, or putrefactive, is due to the action of living organisms upon the material undergoing change; but latterly it has been asserted that the chemical alterations are not really produced by these forms. Without entering into any discussion of the matter, I think it will be admitted on all sides, first, that these living entities are the invariable accompaniments, under ordinary circumstances, of fermentative processes; second, that substances which poison or kill these germs likewise avert these processes. To the substances which have this power I shall apply the name of *antizymotics*, a word derived from the Greek *αντι*, against, and *ζυμωσις*, fermentation.

Antizymotics are used for the purpose of preventing decomposition when it is desired to preserve organic bodies, such as cadavers. They are not, in the strictest sense of the term, disinfectants, and it is evident that they do not hasten, but retard, the desired decomposition, and that they do not affect organic compounds already formed, and therefore ought not to be employed as disinfectants, at least alone, in the cases of cesspools and sewers.

The most important use of substances of the present class is to kill disease-germs,—to destroy the life of the living particles which constitute contagion. They are even more poisonous to the higher forms of animal life than to the infusoria and their congeners, and it is a natural deduction from

their action on living entities that they will affect disease-germs, whether these be, as some believe, distinct organisms, or merely detached particles of human germinal matter.

There are a large number of zymotics, of various power ; but I shall speak in detail of two only, sulphurous and carbolic acids.

ACIDUM SULPHUROSUM—SULPHUROUS ACID.

Sulphurous acid and its salts are most efficient in destroying the low forms of life which are connected with putrefaction and fermentation, and for this reason are excellent preservatives of organic matters. Owing to the ease and cheapness with which it is formed, and its great volatility, sulphurous acid is probably the best agent for disinfecting close apartments. What has already been said in regard to the disinfecting of the atmosphere of course holds good in regard to sulphurous acid ; but often it seems necessary to disinfect the walls, furniture, etc., of a chamber in which a person has passed through a contagious illness. When this is to be done, the method described below may be adopted. When it is to be carried into effect, of course the room should be made as tight as possible, the chimney-places, ventilators, windows, and doors of exit being closed, whilst all the drawers of furniture and the doors of closets are widely opened.

Take a large iron pot, or caldron, put in it a little stand, such as the cheap tripod used by chemists, place on this an iron plate containing flowers of sulphur thoroughly wet with alcohol or (probably better still) with turpentine ; underneath the plate set a tin alcohol lamp ; then put the whole on bricks in the middle of the room. Light the lamp underneath the dish, and if the sulphur does not take fire previously, when it begins actively to melt, ignite it. Leave the room at once, closing the door. Unless the apartment become densely filled with the fumes, far beyond what could be supported by a human being, the attempt at disinfection cannot be of any service. The room should be kept shut up until the sulphurous acid finds its escape through the cracks and disperses. Very often it may be well to repeat the process once or twice, and afterwards to leave the apartment as open and exposed to the air as possible for some days.

Sulphurous acid may be used in a saturated solution for the purpose of destroying disease-germs in the excretions of the sick ;* its action upon vegetable colors of course completely unfits it for many uses. The sulphites and the bisulphites are largely employed to arrest or control fermentation in various processes in the arts.

* For a very interesting account, by Dr. J. Hjaltelin, of the strangling of a smallpox epidemic in Iceland, see *British Medical Journal*, vol. ii., 1871, p. 519. I cannot help believing that the strict isolation and quarantining of the sick had far more to do with arresting the spread of the disease than had the sulphurous acid employed.

ACIDUM CARBOLICUM—CARBOLIC ACID.

Phenic Acid, or *Phenylic Alcohol*, is a substance obtained from coal-tar by distilling at a temperature of between 300° and 400° F., adding to the distillate a hot concentrated solution of potassa, and after this water, separating the light oily matters which rise to the top, and adding muriatic acid to the heavy alkaline bottom layer, when impure carbolic acid separates. This impure carbolic acid is of a dark color, and contains several congeneric bodies, especially xylie and cresylic acids. For disinfectant purposes these appear to be at least of equal value with the carbolic acid, and therefore the crude product of the above-detailed process is very largely used. Carbolic acid is separated from its allies and obtained in a pure state with some difficulty, by a process too complex to be discussed here. When finally procured, it occurs at ordinary temperatures in minute, colorless, transparent plates, or long rhomboidal needles, often fused into a mass, having a hot, corrosive, peculiar taste and a peculiar odor, resembling but entirely different from that of creasote. If, on exposure to the air, phenic acid becomes brown, it contains impurities. When opportunity is afforded, solid carbolic acid absorbs water from the atmosphere and melts into an oily-looking colorless liquid. It is inflammable, neutral to test-paper, but combining with bases; soluble in about twenty parts of water, very soluble in alcohol, acetic acid, ether, glycerine, and the volatile and fixed oils. Nitric acid converts it into picric acid.

CREASOTUM.—*Creasote* is a substance closely allied to carbolic acid, and is prepared from wood-tar, by a process similar to that by which the latter is obtained from coal-tar. It resembles closely carbolic acid solutions, but may at once be distinguished by its odor, which is much more smoky. Moreover, if a splinter of pine be dipped into carbolic acid, and then into muriatic acid, the wood, after the lapse of half an hour, assumes a distinct blue color. This does not occur with creasote. Again, carbolic acid does, and creasote does not, coagulate collodion; and by the action of nitric acid creasote is converted into oxalic acid, resinous matter, and but a small proportion of picric acid. Creasote is at present a very rare drug, the material usually sold under its name in the drug-stores being really a solution of carbolic acid. In their therapeutic value the two substances are probably about equal.

PHYSIOLOGICAL ACTION.—Carbolic acid is exceedingly poisonous to all forms of life, from the lowest to the highest. Much of its employment in medicine depends upon its action on infusoria and fungi; and yet its direct internal and external use in human medicine is quite large. Its physiological action is therefore to be viewed from two distinct stand-points: first, its influence upon the higher animals and man; second, its action on the lowest animal and vegetable forms.

Upon mammals, if not upon all vertebrates, carbolic acid acts as it does upon man. According to Dr. Isidor Neumann (*Archiv für Dermatologie und*

Syphilolog., Jahrgang i., 1869, p. 425), to Dr. Ernest Labée (*Archives Générales*, 6e sér., t. xviii. p. 451, 1871), and to Salkowski (*Pflüger's Archiv für Physiologie*, Bd. v., 1872), when a poisonous dose is given to a frog there is produced a paralytic condition which usually affects first the hind legs,* but eventually spreads to the front ones and involves all parts of the body. After a time there are developed tetanic convulsions, which are apparently reflex in their nature, and are said to be excited by external stimuli or irritations.

Carbolic acid acts upon mammals in very much the same way as upon the batrachian. According to W. Kempster (*American Journal of the Medical Sciences*, July, 1868), in the mouse and rat it causes intense muscular weakness, followed by violent convulsions and stupor. In the rabbit (Neumann, Salkowski), phenylic alcohol produces muscular weakness, often accompanied by tremblings and restlessness, at last giving place to violent convulsions. Before these have fairly set in, the animal is generally unable to stand; and during them he lies on his side, kicking into mid-air. Early in the poisoning the respiration is very much affected: and the death, which usually occurs in the midst of convulsions, appears to be owing to a disturbance of the respiration, since in acute cases the heart is found beating continuously immediately after death. According to the researches of Dr. Jules Lemaire (*De l'Acide Phénique*, 2e édition, Paris, 1865), in the dog symptoms very similar to those detailed above are caused by lethal doses of the drug; and Husemann (*Schmidt's Jahrbücher*, Bd. clv. p. 274) states that in mammals and in birds the characteristic phenomena of carbolic acid poisoning are clonic convulsions, sinking of the temperature, diminution of sensibility, dyspnoea, free salivation and secretion of tears, keratitis and conjunctivitis. According to the latter authority, albuminuria and hæmaturia are occasional phenomena.

Upon man, carbolic acid acts as upon other mammals. Reserving the details for the section on Toxicology, it is sufficient for our present purpose to state that the prominent symptoms induced by lethal doses are disturbance of respiration, coma, muscular weakness, and, in some cases, convulsions.

A closer investigation of the action of large doses of carbolic acid is best made by studying the effects upon the different systems *seriatim*.

Nervous system.—Upon the cerebrum phenylic alcohol appears to exert a direct influence, which, although not very intense in the lower animals, in the higher species, and especially in man, results in the early production of stupor.

According to the experiments of Salkowski (*loc. cit.*, p. 343), the convulsions are not of cerebral origin, since they take place after division of the cervical spinal cord; nor are they peripheral, since they do not occur in a limb whose connection with the spine has been severed by division of the nerve, and do take place in a leg which has been protected against the local action of the poison by tying the artery. Not being either cerebral or peripheral, these convulsions must be spinal.

* According to Lemaire, when a frog is allowed to swim in water impregnated with carbolic acid, the front legs are the first affected.

Dr. Ernest Labée agrees with Salkowski in regarding the convulsions as of central origin. Like the German investigator, he found that section of a nerve arrested the convulsions in the tributary muscles, and that tying the artery of a limb did not prevent their development. On the other hand, the experiments of the French investigator in regard to which centres are affected differed in their results from those of Salkowski. He found that the convulsions did not occur if the cord or the medulla had been previously divided, and that destruction of the whole encephalon had a similar result; but that if only the cerebral hemispheres and the optic lobes were removed, the convulsive phenomena developed in their usual manner. Admitting the truth of these asserted facts, the conclusion is inevitable that the convulsions produced by carbolic acid originate at the base of the brain, and are epileptiform.

No detailed accounts of the experiments of Labée have been published; and Salkowski (*loc. cit.*, p. 343) simply states that in a rabbit whose cord had been cut previous to the poisoning, convulsive movements (*Zuckungen*) occurred in the hind legs, although weaker than those in the front legs.

The experiments of the two investigators are, with our present light, irreconcilable; but it seems to me that the results of Labée are the most probable. The experiments of Salkowski are insufficient to carry complete conviction: indeed, he states only that convulsive movements were present to some extent after section of the cord. Further, according to his own researches, the nerves and muscles are not seriously affected by the poisoning; and, this being so, the co-existence of spinal convulsions and of paralysis is apparently inexplicable, and seems to demand the consentaneous existence of an exalted and a depressed state of the spinal motor ganglia.

As already stated, the muscles and nerves are not seriously implicated in carbolic acid poisoning; after death, galvanic stimulation of a muscle, either directly or through its nerve, causing vigorous contraction (Salkowski, *loc. cit.*, p. 338; Hoppe-Seyler, *loc. cit.*, p. 476).

Circulation.—The action of carbolic acid upon the circulation has not been thoroughly studied. According to Labée (*loc. cit.*, p. 464) and to Salkowski, in acute poisoning the heart is found pulsating regularly directly after death, but is finally arrested in diastole; and in slow poisoning death may be immediately produced by diastolic arrest. Salkowski asserts that the rapidity of the circulation in a frog's web can with the microscope be seen to be at first increased by the hypodermic injection of carbolic acid, and afterwards to be very greatly diminished; and the French observer states that the systoles in the early stages of the poisoning can be seen to increase in energy, and the vessels to contract. I do not think much weight is to be attached to such evidence as this; and the only manometrical studies which I have met with are those of Hoppe-Seyler (*Pflüger's Archiv für Physiologie*, 1872, Bd. v. p. 475).

This observer noted that the arterial pressure did not vary much under the action of the poison until the convulsions came on, when it rose very greatly,

and remained at a high level for a considerable length of time, after which it fell far below the normal point. I do not see that these facts prove that carbolic acid acts as a cardiac stimulant during any stage of the poisoning; the motor disturbance was in itself sufficient to account for the rise of the arterial pressure. Substances which arrest the heart in diastole are as a rule cardiac depressants, and any direct influence carbolic acid has upon the heart is very probably of such nature. This point needs further investigation before any positive opinion can be arrived at.

Respiration.—According to Salkowski (*loc. cit.*, p. 344), Labée, and other authorities, in the first stages of carbolic acid poisoning the respiration is remarkably increased in frequency. This acceleration Salkowski believes to be due partly to a stimulant action upon the peripheral vagi, and partly to a similar influence upon the respiratory centres.

He states that the respirations are very shallow, and that the diaphragm scarcely participates at all in them, but that if the cervical vagi be cut they become much slower, deep, and regular. On the other hand, if carbolic acid be given to an animal suffering from section of the pneumogastrics, the slow breathing is very much accelerated. From the former of these facts the German investigator draws the conclusion that the accelerated breathing produced by phenylie alcohol is in part due to a stimulation of the peripheral vagi, and from the latter fact that it partly arises from a similar action upon the respiratory centres.

The effects of carbolic acid are, in all probability, due to a direct action upon the organs concerned. Labée (*loc. cit.*, p. 464) states that, although changes occur in the blood-corpuscles when carbolic acid is added to the blood outside of the body, yet in the blood of animals killed by it no alterations can be detected in these minute bodies; and Hoppe-Seyler (*Pflüger's Archiv*, Bd. v. p. 476) confirms this observation of the French investigator.

Post-mortem examinations of animals killed by carbolic acid have yielded varying results. In Lemaire's investigation, nothing abnormal was found except intense injection of the alimentary mucous membrane, a pseudo-membranous and purulent inflammation of the bronchial tubes, with a disseminated lobular pneumonia or else congestion of the lungs and congestion of the nerve-centres. Prof. Bruckmüller, in Neumann's investigation (*loc. cit.*, p. 429), found the cells of the liver and kidneys in a state of fatty degeneration. This process, which seemingly was the counterpart of the changes in phosphorus-poisoning, was always more advanced in the kidneys than in the liver. Dr. Neumann states that it was found in a number of autopsies, and that it is a constant phenomenon; but Salkowski (*loc. cit.*, p. 273) was unable to find it in a number of examinations. In man, the post-mortem appearances are very much the same as in animals. If the acid has been ingested in a concentrated form, white, hardened spots are found upon the mucous membrane of the mouth, œsophagus, stomach, and even intestines. They are, of course, due to the local action of the poison, and are sometimes blackish in the

centre, or even blackish throughout, and very generally are surrounded by a red inflammatory zone. The liver, spleen, kidneys, and indeed all the organs, are found filled with dark, imperfectly-coagulated blood, such as is habitually found after death from asphyxia. According to Husemann, the fatty degeneration of the liver and kidneys is not either in man or in animals a constant or characteristic phenomenon of carbolic acid poisoning. Reuder found the renal epithelium degenerated in a man who had been fatally poisoned by the drug (*Journal de Pharmacie et de Chimie*, p. 456, Dec. 1871).

As the internal use of carbolic acid in such diseases as gangrene of the lungs is so closely connected with the question of its chemical history in the system and its elimination from the body, the latter is of very great interest. Since carbolic acid coagulates albumen, its absorption unchanged into the blood would seem a matter of doubt: yet in some form or other it certainly is absorbed, as is proven by the history of its elimination and by its having been found in the blood (Hoppe-Seyler, *Pflüger's Archiv*, Bd. v. p. 479). In exactly what form it circulates in the blood is not known; but most probably it is as an alkaline carbolate. Lemaire (*loc. cit.*, p. 77) states that it may be found in the breath of poisoned animals; but Hoffmann asserts that it is burnt up in the system, because he failed to detect it in any of the secretions. In this conclusion he is, however, certainly in error; for it has been distinctly proven that carbolic acid is rapidly eliminated from the system. It has been detected in the urine by Almén (*Zeitschrift f. Analyt. Chemie*, Bd. x. Heft vii.), by Patrouillard (*Journal de Pharmacie et de Chimie*, Dec. 1871, p. 459), by Salkowski (*Pflüger's Archiv*, Bd. v.), by Hoppe-Seyler (*loc. cit.*), by Waldenström (*Zeitschrift des Allgemein. Apothek.-Vereines*, Jan. 10, 1872), and by Hauxmann (*Ibid.*); and Hoppe-Seyler (*loc. cit.*, p. 480) has detected it in the saliva. It is probably eliminated in all the secretions. Salkowski believes that it is thrown off in the form of a carbolate, because, on the distillation of urine containing it, it does not pass over until the urine is strongly acidified. It is very probable that when in small amount carbolic acid escapes only in union with an alkali. In a fatal case of poisoning, Patrouillard (*loc. cit.*, p. 460), however, obtained an oily fluid, believed to be pure carbolic acid, by shaking the urine with ether, allowing the mixed fluids to separate, and removing the ethereal layer and evaporating.

Although, as stated, carbolic acid is largely eliminated from the system, a portion of it is probably burnt up in the body. A blackish urine is not a rare phenomenon in poisoning by phenic acid, especially when the acid has been absorbed from a wound or other external surface. The black coloring-matter of such urine is in all probability an educt from carbolic acid, formed by its partial oxidation. Hauxmann has proven that it is not altered hæmatin or any fixed coloring-principle, by finding that the urine is cleared up by heating after the addition of an acid; and his conclusion is corroborated by the observation of Dr. Stevenson, of Guy's Hospital (*British Medical Journal*, April, 1870), who found that the black urine does not contain more than a

normal proportion of iron. When carbolic acid is oxidized outside of the body, as by the action of the permanganate of potash, oxalic acid is formed; and Salkowski has found that when phenic acid is given to animals oxalic acid appears in the urine. The evidence therefore indicates very strongly that, partly by elimination and partly by oxidation, the system rapidly rids itself of carbolic acid. In conformity with this is the fact observed in several cases by Salkowski, that twenty-four hours after the administration of the drug indications of its presence are no longer to be found in the urine.

Städeler (*Ann. d. Chem. und Pharm.*, Bd. lxxvii. p. 17) discovered that when sulphuric acid was freely added to cow's urine the latter yielded upon distillation carbolic acid, and concluded therefrom that normal urine contains carbolic acid. His asserted fact has been corroborated by Buliginsky (*Hoppe-Seyler's Medicinisch-chemische Untersuchungen*, p. 234) and by Hoppe-Seyler (*Pflüger's Archiv*, 1872, Bd. v. p. 470), and is without doubt true not only of the urine of cattle, but also of that of men, dogs, horses, and probably other animals. Without discussing the matter in detail, it is sufficient for our purpose to state that Hoppe-Seyler has shown, however, that carbolic acid does not really exist in the urine, since it cannot be obtained from it by distillation, even after an excess of acetic acid has been added, and that it is formed during the process employed by Städeler, most probably, out of the indican of the urine.

The local action of carbolic acid is a very decided one. When applied to the skin, it produces at once a burning pain, and in a few minutes a peculiar white spot. If the application be removed, the pain continues for some minutes, and the white color changes to a dark or red stain, which gradually fades away as the skin desquamates. On a prolonged application, carbolic acid does not blister, but causes the formation of an eschar. A curious local action of carbolic acid, to which attention was drawn almost simultaneously by Dr. Erasmus Wilson (*Journal of Cutaneous Medicine*, June, 1870) and by Dr. J. H. Bill, U.S.A. (*American Journal of the Medical Sciences*, Oct. 1870), is due to the property which it has when applied in concentrated form of causing very great local anæsthesia. The complete loss of feeling is not confined to the tissue killed by the drug, but extends some little distance inwards.

THERAPEUTICS.—In the doses in which it is usually given, carbolic acid exerts no perceptible effect upon the system. Upon the idea that the so-called zymotic diseases are due to a sort of fermentation in the blood, that they are the result of changes set up in the vital fluid by living organisms similar to the yeast-plant or to the vibrio of putrefaction, carbolic acid has been quite largely used in such diseases. The zymotic theory is, however, at the very best merely a plausible speculation; and clinical experience with the acid in these diseases has certainly demonstrated its uselessness. It has been extensively employed in *typhus fever*, in *pyæmia*, in *smallpox*, and in other allied affections, but at present is rarely used and is of no value. The study of its

physiological action has failed to show the possession of any property which should render the medicine valuable in constitutional diseases, and clinical experience has borne this out: so that it is employed directly in medicine only for its local effects.

Internally it is of value in *nervous vomiting*, especially when there is an irritability of the gastric nerves. One or two drops of it, or of creasote, may be given in emulsion every three or four hours in such cases. The good which it achieves is probably dependent upon its local anæsthetic properties.

In *gangrene of the lung*, the internal administration of carbolic acid, combined with the use of a weak solution (ten drops to the ounce) by atomization, is sometimes of service.

Its external use is much more important than its internal employment; but it belongs to the domain of the surgeon rather than of the physician, and I will only discuss it briefly. As a *caustic*, carbolic acid is not available when large masses of tissue are to be destroyed, but may often be employed with advantage against *condylomata* and similar growths. Even in such cases, to be efficient, it must be in the most concentrated form. In *diphtheria*, *ulcerated sore throat*, and *aphthous stomatitis*, its concentrated solution in glycerine may be carefully applied, by means of a camel's-hair brush or a mop, as a mild caustic scarcely capable of destroying sound tissue. In various forms of *indolent ulcer*, in *ill-conditioned wounds*, carbolic acid affords a very useful stimulant application; in "*burns*," properly diluted, it is one of the very best remedies that can be used, relieving pain by its anæsthetic properties and at the same time lessening suppuration and facilitating cicatrization.

As a local anæsthetic, carbolic acid has not been used so widely as it seems to deserve. Dr. Bill (*loc. cit.*) has employed it in a number of cases of minor operations, always with the result of preventing or greatly mitigating pain. His plan in opening a felon is to soak the fingers for fifteen minutes in warm water containing three per cent. of the acid, and then to draw a brush dipped in the concentrated acid along the line of the incision. Sometimes, when a deep incision is necessary, a sensitive part is reached. Under these circumstances he is accustomed to brush out the wound anew with the anæsthetic. In operations requiring much cutting, this method is not available; but whenever only the skin is to be divided, as in opening *abscesses*, it appears to be very successful.

TOXICOLOGY.—The number of fatal cases of carbolic acid poisoning now on record is quite large, and the list is constantly growing. The symptoms, although varying within certain limits, are, on the whole, quite uniform. They almost always appear in a very short time after the ingestion of the poison. Sometimes the rapidity of the fatal result almost equals that of prussic acid poisoning. Thus, Dr. Taylor, U.S.N. (*Philadelphia Medical Times*, vol. ii. p. 284) records a case in which about an ounce is supposed to have been ingested, and in which the man fell unconscious within ten seconds after

taking the fatal draught, two minutes afterwards was totally unconscious, pulseless, with irregular distant gasping respirations, and in less than a minute later was dead, apparently from cardiac paralysis, since the impulse of the heart was entirely lost before the cessation of respiration. Generally some minutes elapse before the symptoms develop themselves: nausea, cold sweats, stupor deepening rapidly into insensibility and collapse, are the most frequent phenomena. During the period of insensibility, complete abolition of reflex movements and anæsthesia of the mucous membranes have sometimes been noted (case, *Journal de Pharmacie et de Chimie*, Dec. 1871): indeed, it is scarcely doubtful that in all cases both sensibility and reflex movements are profoundly affected. Convulsions are only exceptionally present. The symptoms of collapse are usually well developed, and the pulse is generally feeble and very frequent, but has been noted as being reduced to from forty to fifty per minute (case, *Medical Times and Gazette*, April, 1871). The respirations are mostly hurried and shallow in the advanced stages, or in very rapid cases they are irregular and suspended at intervals. Death, as has already been stated, may occur in a very few minutes, but usually the patient lives from one to ten hours, and life has been protracted for sixty hours (case, *Sydenham Year-Book*, p. 446, 1871-72; amount taken, one and a half ounces of the commercial acid). In some cases a great amendment has occurred and consciousness been restored, but after some hours rather sudden fatal collapse has come on (case, *British Medical Journal*, Feb. 1871). The minimum fatal dose of carbolic acid is not known; but half an ounce would probably prove fatal, since death from this quantity has been reported once in a man weakened by typhoid fever (*Medical Times and Gazette*, p. 474, vol. ii., 1870), and once in a healthy man (*Philadelphia Medical and Surgical Reporter*, Jan. 1870).

The free external use of carbolic acid is by no means devoid of danger: indeed, in more than one case it has caused death. Prof. R. Köhler reports a very interesting case (*Schmidt's Jahrbücher*, Bd. clv. p. 276). Two journeymen joiners, suffering from scabies, applied externally each about a half-ounce of carbolic acid, in watery solution. One of them was found dead. His fellow, who suffered from unconsciousness and drunken delirium ending in unquiet sleep, stated, after recovery, that directly after rubbing himself with the solution he had giddiness, that seven or eight minutes later his companion complained of burning, but that of what took place after this he knew nothing. Two fatal cases of poisoning by the external use of carbolic acid, and one of recovery after complete unconsciousness, are reported in the *Bulletin Thérapeutique*, t. lxxv. p. 285. It is scarcely necessary to refer in detail to cases in which serious results have followed the surgical use of carbolic acid.* A form of chronic poisoning, with loss of appetite, weak-

* Consult *British Medical Journal*, March 1, 1873,—death from absorption by a wound four inches long; also *New York Medical Gazette*, April, 1871; also *British Medical Journal*, 1868, p. 220,—two fatal cases.

ness, and a tendency to collapse, resulting from the continued surgical use of the acid, has been described by Dr. Wallace (*British Medical Journal*, April, 1870). In vol. ii., 1870, of the *Medical Times and Gazette*, is narrated a curious case of fatal poisoning from inflammation of the external parts and of the rectum, produced in a child by sitting upon a block on which some of the acid had been thrown. A serious result which has been known to follow the application of phenol to a slight wound of the finger is gangrene of the whole member (*L' Abeille Médicale*, Dec. 11, 1871).

The diagnosis of carbolic acid poisoning during life ought in most cases to be practicable. Although the symptoms simulate some forms of apoplexy too closely for the diagnosis to be made from them, yet very generally the odor of the drug can be perceived about the person of the victim, and close examination of the mouth will nearly always reveal traces of the local action of the acid, in the form of *white, hardened, or corrugated* patches of mucous membrane. These, in conjunction with the symptoms, are diagnostic. At post-mortem a strong odor of carbolic acid can almost always be perceived when the body is opened, and the mucous membrane of the stomach affords very reliable evidence as to the cause of death.

Blackish urine, if it be present, is also decisive as to the nature of the case: its absence does not, however, prove that the patient is not suffering from carbolic acid poisoning.

The treatment of carbolic acid poisoning is far from satisfactory. The symptoms are generally developed so rapidly that there is very little time to prepare and administer an antidote, even if we possessed a perfect one. I do not think there is any known substance which can be thoroughly relied on; but the recent experiments of Husemann indicate that the alkalis have some power in controlling the lethal action, provided they are exhibited in solution and in great excess. Lime is probably the best of them, especially if given in the form of the *saccharate*. The latter may be prepared as follows. Dissolve sixteen parts of sugar in forty parts of distilled water, and add five parts of caustic lime. Digest for three days, stirring from time to time, filter, and evaporate to dryness. The product thus obtained dissolves easily in water. Husemann's experiments indicate that the fats are of very little, if any, use; but Lemaire found that the solution of carbolic acid in oil is much less poisonous to animals than the acid itself,—probably because it is much more slowly absorbed. Consequently, the free ingestion of sweet oil or of castor oil should be practiced in a case of poisoning, and no time should be lost in emptying the stomach. Emetics are generally useless, the local action of the poison having so completely benumbed the sensibility of the stomach as to render it unsusceptible. For this reason, the stomach-pump is in these cases a necessity. After the viscus has been emptied, the usual measures for the relief of collapse may be instituted, but probably have very little influence upon the final result.

Carbolic Acid as an Antiseptic.

The action of carbolic acid upon man having been discussed, it is now proper to take notice of its influence upon the lower organisms, and of the uses which grow out of that influence. It is beyond dispute that carbolic acid has very great power in killing both animal and vegetable ferments, and in arresting the changes which they induce. The question which is at present most interesting and worthy of discussion is as to the degree of its influence, —as to the power of carbolic acid, when compared with other substances. There are several ways in which this power may be measured.

In August, 1870, Dr. John Dougall (*London Lancet*) published an account of a series of experiments which he had made as to the minimum quantity of various reagents required to arrest the movements of spermatozoa and to kill infusoria. His results are expressed, in fractional parts, in the following table :

Drugs.	Spermatozoa.	Infusoria.
Chloride of strychnia.....	1-30000	1-450
Chloride of arsenic.....	1-18000	1-8000
Nitric acid.....	1-18000	1-1500
Hydrochloric acid.....	1-15000	1-2000
Sulphuric acid.....	1-15000	1-1500
Alcohol.....	1-12500	1-750
Corrosive sublimate.....	1-7000	1-6000
Nitrate of silver.....	1-6500	1-6000
Strong acetic acid.....	1-10000	1-500
Oxalic acid.....	1-7500	1-1500
Chloride of zinc.....	1-7500	1-600
Picric acid.....	1-3700	1-450
Tartrate of antimony.....	1-4000	1-450
Hydrocyanic acid.....	1-5000	1-250
Carbolic acid.....	1-1000	1-750
Camphor.....	1-2500	1-400
Tincture of iodine.....	1-500	1-400
Solution of chloride of lime.....	1-500	1-300
Common salt.....	1-10	1-50

In these experiments of Dougall the action of the drugs upon animal life was alone tested. But the relations of drugs to low animal and vegetable forms are not identical, and the action of a drug upon infusoria, much less upon spermatozoa, is no criterion as to its influence upon fungi. Moreover, fermentation is caused not by infusoria but by fungi; and it is therefore very apparent that the experiments of Dougall are not decisive.

We are indebted to Dr. P. Grace Calvert for a series of experiments which in a measure supplement those of Dr. Dougall. Dr. Calvert, using a standard solution of albumen, added to equal quantities of it in test-tubes one-thousandth part of the substance whose power over putrefaction it was desired to test, and then, placing all of the test-tubes side by side in a rack, watched for the development of fungi and of vibrios. The following table expresses his results :

	Days required for the Development of		
	Fungi.	Vibrios.	Putrid Odors in Albumen, kept at 80°.
1. STANDARD SOLUTIONS.			
Albumen kept in laboratory for comparison..	18	12	16
2. ACIDS.			
Sulphurous acid.....	21	11	45
Sulphuric acid.....	9	9	16
Nitric acid.....	10	10	16
Arsenious acid.....	18	22	None.
Acetic acid.....	9	30	None.
Prussic acid.....	None.	9	35
3. ALKALIES.			
Caustic soda.....	18	24	72
Caustic potash.....	16	26	85
Caustic ammonia.....	20	24	26
Caustic lime.....	None.	13	14
4. CHLORINE COMPOUNDS.			
Solution of chlorine.....	22	7	16
Chloride of sodium.....	19	14	16
Chloride of calcium.....	18	7	11
Chloride of aluminium.....	21	10	16
Chloride of zinc.....	53	None.	38
Bichloride of mercury.....	81	None.	None.
Chlorinated lime.....	16	9	9
Chloride of potassium.....	19	17	38
5. SULPHUR COMPOUNDS.			
Sulphate of calcium.....	19	9	14
Protosulphate of iron.....	15	1	16
Bisulphite of calcium.....	18	11	16
Hyposulphite of sodium.....	18	11	11
6. PHOSPHATES.			
Phosphate of sodium.....	17	13	16
Phosphate of calcium.....	22	7	16
7.			
Permanganate of potassium.....	22	9	11
8. TAR SERIES.			
Carbolic acid.....	None.	None.	None.
Cresylic acid.....	None.	None.	None.
9. SULPHOCARBOLATES.			
Sulphocarbolate of potassium.....	17	18	35
Sulphocarbolate of sodium.....	19	18	26
Sulphocarbolate of zinc.....	17	None.	None.
10.			
Sulphate of quinine.....	None.	25	None.
Picric acid.....	19	17	26
Pepper.....	None.	8	16
Turpentine	42	14	35
11.			
Charcoal.....	21	9	None.

In comparing the results stated in the above table, the substances can be classed under four distinct heads, viz.: those which prevent the development of protoplasmic and fungus life; those which prevent the production of vibrio life but do not prevent the appearance of fungus life; those which permit the production of vibrio life but prevent the appearance of fungus life; and those which do not prevent the appearance of either protoplasmic or fungus life.

The first class contains only two substances,—carbolic and cresylic acids.

In the second class, also, there are only two compounds,—chloride of zinc and bichloride of mercury.

In the third class there are five substances,—lime, sulphate of quinine, pepper, turpentine, and prussic acid.

In the fourth class are included the remaining twenty-five substances.

The acids, while not preventing the production of vibrio life, have a marked tendency to promote the growth of fungus life. This is especially noticeable in the case of sulphuric and acetic acids.

Alkalies, on the contrary, are not favorable to the production of fungus life, but promote the development of vibrios.

Dr. Calvert also reversed, as it were, these experiments, by allowing the vibrios and fungi to develop in an albuminous fluid and then adding one-thousandth part of the reagent. He found that the various reagents used were divisible, according to their effects, into seven classes, as follows:

The first class includes those substances which completely destroyed the locomotive power of the vibrios immediately, and completely prevented their regaining it during the time the experiments were conducted:—cresylic acid.

The second class contains those compounds which nearly destroyed the locomotive power of all the vibrios present when added, and afterwards only one or two could be seen swimming about in each field:—carbolic acid, sulphate of quinine, chloride of zinc, and sulphuric acid.

The third class are those which acted injuriously on the vibrios on their addition, leaving only a small number retaining the power of swimming, but which allowed the vibrios gradually to increase in number, the field nevertheless containing less life after sixteen days than the standard albumen solution:—picric acid and sulphocarbolate of zinc.

The fourth class includes those substances which acted injuriously at first, but permitted the vibrios to regain their former locomotive power, so that the fluid after sixteen days contained as much vibrio life as the standard putrid albumen:—chloride of aluminium, sulphurous acid, and prussic acid.

The fifth class contains those compounds which acted injuriously at first, destroying the locomotive power of most of the vibrios, but which afterwards permitted the vibrios to increase more rapidly than in the standard albumen solution:—bleaching-powder, bichloride of mercury, chlorine solution, caustic soda, acetic and nitric acids, sulphate of iron, and the sulphocarbolates of potassium and sodium.

The sixth class contains those compounds which exercised no action on the animalcules, either at first or after sixteen days:—arsenious acid, common salt, chloride of calcium, chlorate of potassium, sulphate of calcium, bisulphite of calcium, hyposulphite of sodium, phosphate of calcium, turpentine, and pepper.

The seventh class includes those substances which favor the production of animalcules and promote putrefaction:—lime, charcoal, permanganate of potassium, phosphate of sodium, and ammonia.

On comparing these results with those obtained by Dougall, they are seen to be very much more favorable to the antiseptic use of carbolic acid,—so much more favorable, indeed, as to indicate either that one or the other set of experiments is not absolutely accurate. There is one very great fallacy which underlies the whole of Dr. Calvert's work, and which may in some measure account for the difference. It is this: the putrefaction of albumen does not represent, or is not equal to, the putrefaction of other organic bodies, especially of complex organic bodies. This is strikingly shown by Calvert's own trials with gelatine. In these experiments gelatine was used instead of albumen, all the processes being conducted as in the first set; and it was found that the protosulphate of iron completely prevented the development of either protoplasmic or vibrionic life, although when albumen was used it had very little effect upon either.

Both Dougall and Calvert's experiments show that the corrosive sublimate is very poisonous to the low forms of life.

With the evidence at our command, I do not think that we are in a position to decide at present as to the comparative value of carbolic acid in destroying disease-germs. We have, however, abundant evidence as to its absolute power. P. C. Plugge (*Hygien's Archiv*, 1872, Bd. v. p. 540) found that one part of carbolic acid in five hundred parts of water killed paramecia and colpoda instantly, but required about an hour to put an end to the motions of vibrios; one part in eight hundred of water killed the larger infusoria in two or three minutes, the smaller after a length of time; one part in a thousand of water destroyed the colpoda in from eight to fifteen minutes; the vibrios were still living at the end of twenty-four hours. One part in two thousand of water had no perceptible effect upon the larger organisms.

These experiments of Plugge are especially valuable because they were made with complex decaying organic solutions, such as infusion of hay and water out of stagnant pools and gutters. Dr. Plugge himself deduces from them the conclusion that a solution of carbolic acid weaker than one part in a hundred cannot be trusted to kill quickly all forms of life; and, as in disinfecting discharges, etc., it is necessary or desirable to kill at once the organic germs, it certainly is not proper to rely upon a solution of the strength of less than one per cent.

ABSORBING DISINFECTANTS.

The substances which are used to act as disinfectants by absorbing the products of decomposition are earth, charcoal, and plaster of Paris. All of these, when used in sufficient quantity, are very efficient in preventing the escape of foul gases; but none of them affect disease-germs, and none of them should for a moment be relied upon in cases of contagions. They, if properly employed, often not merely absorb the products of decomposition, but, by removing the moisture, check the decay. For this purpose they should, of course, be very dry when used.

APPENDIX.

WEIGHTS AND MEASURES OF THE UNITED STATES PHARMACOPEIA.

One Pound,	lb	=	12 Ounces	=	5,760 Grains.
One Ounce,	℥	=	8 Drachms	=	480 Grains.
One Drachm,	ʒ	=	3 Scruples	=	60 Grains.
One Scruple,	ʒ	.	.	.	20 Grains.
One Grain,	gr.	.	.	.	1 Grain.
One Gallon,	C	=	8 Pints	=	61,440 Minims.
One Pint,	O	=	16 Fluidounces	=	7,680 Minims.
One Fluidounce,	f℥	=	8 Fluidrachms	=	480 Minims.
One Fluidrachm,	fʒ	.	.	.	60 Minims.
One Minim,	ʒ	.	.	.	1 Minim.

WEIGHTS AND MEASURES OF THE METRICAL OR FRENCH SYSTEM.

MEASURES OF LENGTH.

One Myriametre	=	10,000 Metres.
One Kilometre	=	1,000 Metres.
One Hectometre	=	100 Metres.
One Decametre	=	10 Metres.
One METRE	=	the ten-millionth part of a quarter of the meridian of the earth.
One Decimetre	=	the tenth part of one Metre, or 0.1 Metre.
One Centimetre	=	the hundredth part of one Metre, or 0.01 Metre.
One Millimetre	=	the thousandth part of one Metre, or 0.001 Metre.

WEIGHTS.

One Myriagramme	=	10,000 Grammes.
One Kilogramme	=	1,000 Grammes.
One Hectogramme	=	100 Grammes.
One Decagramme	=	10 Grammes.
One GRAMME	=	the weight of a cubic Centimetre of water at 4° C.
One Decigramme	=	the tenth part of one Gramme, or 0.1 Gramme.
One Centigramme	=	the hundredth part of one Gramme, or 0.01 Gramme.
One Milligramme	=	the thousandth part of one Gramme, or 0.001 Gramme.

MEASURES OF CAPACITY.

One Myrialitre	=	10 cubic Metres, or the measure of 10 Milliers of Water.
One Kilolitre	=	1 cubic Metre, or the measure of 1 Millier of Water.
One Hectolitre	=	100 cubic Decimetres, or the measure of 1 Quintal of Water.
One Decalitre	=	10 cubic Decimetres, or the measure of 1 Myriagramme of Water.
One LITRE	=	1 cubic Decimetre, or the measure of 1 Kilogramme of Water.
One Decilitre	=	100 cubic Centimetres, or the measure of 1 Hectogramme of Water.
One Centilitre	=	10 cubic Centimetres, or the measure of 1 Decagramme of Water.
One Millilitre	=	1 cubic Centimetre, or the measure of 1 Gramme of Water.

RELATION OF WEIGHTS AND MEASURES OF THE U. S. PHARMACOPEIA TO EACH OTHER.

In distilled water at the temperature of 60°.

One Pound	=	0.7900031 Pint	=	6067.2238 Minims.
One Ounce	=	1.0533376 Fluidounces	=	505.6019 Minims.
One Drachm	=	1.0533376 Fluidrachms	=	63.2002 Minims.
One Scruple	=		=	21.0667 Minims.
One Grain	=		=	1.0533 Minims.
One Gallon	=	10.1265427 Pounds	=	58328.8862 Grains.
One Pint	=	1.2658178 Pounds	=	7291.1107 Grains.

One Fluidounce	=	0.9493633 Ounce	=	455.6944 Grains.
One Fluidrachm	=	0.9493633 Drachm	=	56.9618 Grains.
One Minim	=		=	0.9493 Grain.

RELATION OF MEASURES OF THE U. S. PHARMACOPŒIA TO CUBIC MEASURE.

One Gallon	=	231.	Cubic Inches.
One Pint	=	28.875	Cubic Inches.
One Fluidounce	=	1.80468	Cubic Inches.
One Fluidrachm	=	0.22558	Cubic Inch.
One Minim	=	0.00375	Cubic Inch.

RELATION OF WEIGHTS OF THE U. S. PHARMACOPŒIA TO METRICAL WEIGHTS.

<i>Fractions of a grain in Milligrammes.</i>		<i>Grains in equivalent metrical weights.</i>		<i>Drachms, Ounces, and Pounds in equivalent metrical weights.</i>	
Grain.	Milligrammes.	Grains.	Centigrammes.	Drachms.	Grammes.
$\frac{1}{64}$	= 1.012	1	= 6.479	1	= 3.887
$\frac{1}{50}$	= 1.079		Decigrammes.	2	= 7.775
$\frac{1}{30}$	= 1.295	2	= 1.295		Decagrammes.
$\frac{1}{48}$	= 1.349	3	= 1.943	3	= 1.166
$\frac{1}{40}$	= 1.619	4	= 2.591	4	= 1.555
$\frac{1}{36}$	= 1.799	5	= 3.239	5	= 1.943
$\frac{1}{30}$	= 2.159	6	= 3.887	6	= 2.332
$\frac{1}{25}$	= 2.591	7	= 4.535	7	= 2.721
$\frac{1}{24}$	= 2.699	8	= 5.183	Ounces.	
$\frac{1}{20}$	= 3.239	9	= 5.831	1	= 3.1103
$\frac{1}{16}$	= 4.049	10	= 6.479	2	= 6.2206
$\frac{1}{15}$	= 4.319	12	= 7.775	3	= 9.3309
$\frac{1}{12}$	= 5.399	15	= 9.718		Hectogrammes.
$\frac{1}{10}$	= 6.479		Grammes.	4	= 1.2441
$\frac{1}{8}$	= 8.098	16	= 1.036	5	= 1.5551
$\frac{1}{6}$	= 10.798	20	= 1.295	6	= 1.8661
$\frac{1}{5}$	= 12.958	24	= 1.555	7	= 2.1772
$\frac{1}{4}$	= 16.197	25	= 1.619	8	= 2.4882
$\frac{1}{3}$	= 21.597	30	= 1.943	9	= 2.7992
$\frac{1}{2}$	= 32.395	40	= 2.591	10	= 3.1103
		50	= 3.239	11	= 3.4213
		60	= 3.887	Pounds.	
				1	= 3.7324
				2	= 7.4648
					Kilogrammes.
				3	= 1.1197

RELATION OF METRICAL WEIGHTS TO WEIGHTS OF THE U. S.
PHARMACOPEIA.

<i>Metrical Weights.</i>	<i>Exact equivalents in grains.</i>	<i>Approximate equivalents in grains.</i>	<i>Metrical Weights.</i>	<i>Exact equivalents in grains.</i>	<i>Approximate equivalents in Troy Weight.</i>
Milligrammes.			Grammes.		
1 =	.0154	$\frac{1}{65}$	1 =	15.434	gr. xv.
2 =	.0308	$\frac{1}{32}$	2 =	30.068	3ss.
3 =	.0463	$\frac{1}{22}$	3 =	46.302	ʒij.
4 =	.0617	$\frac{1}{16}$	4 =	61.736	ʒi.
5 =	.0771	$\frac{1}{13}$	5 =	77.170	ʒiv.
6 =	.0926	$\frac{1}{11}$	6 =	92.604	ʒiss.
7 =	.1080	$\frac{1}{9}$	7 =	108.038	ʒvss.
8 =	.1234	$\frac{1}{8}$	8 =	123.472	ʒij.
9 =	.1389	$\frac{1}{7}$	9 =	138.906	ʒviij.
Centigrammes.			Decagrammes.		
1 =	.1543	$\frac{1}{6}$	1 =	154.340	ʒiiss.
2 =	.3086	$\frac{1}{3}$	2 =	308.680	ʒv.
3 =	.4630	$\frac{6}{13}$	3 =	463.020	ʒviiss.
4 =	.6173	$\frac{7}{11}$	4 =	617.360	ʒx.
5 =	.7717	$\frac{8}{4}$	5 =	771.701	ʒxiij.
6 =	.9260	$\frac{9}{10}$	6 =	926.041	ʒxv.
7 =	1.0803	1	7 =	1,080.381	ʒxviij.
8 =	1.2347	1 $\frac{1}{4}$	8 =	1,234.721	ʒxx.
9 =	1.3890	1 $\frac{1}{8}$	9 =	1,389.062	ʒxxiiij.
Decigrammes.			Hectogrammes.		
1 =	1.543	1 $\frac{1}{2}$	1 =	1,543.402	ʒiii ʒv.
2 =	3.086	3	2 =	3,086.804	ʒvj ʒiij.
3 =	4.630	4 $\frac{1}{2}$	3 =	4,630.206	ʒix ʒv.
4 =	6.173	6	4 =	6,173.609	ʒbi ʒviij.
5 =	7.717	7 $\frac{1}{2}$	5 =	7,717.011	ʒbi ʒiv.
6 =	9.260	9	6 =	9,260.413	ʒbi ʒviij.
7 =	10.803	11	7 =	10,803.816	ʒbi ʒx ʒiv.
8 =	12.347	12 $\frac{1}{2}$	8 =	12,347.218	ʒbij ʒi ʒv.
9 =	13.890	14	9 =	13,890.620	ʒbij ʒv.
Kilogramme.			Kilogramme.		
			1 =	15,434.023	ʒbij ʒviij.
Myriogramme.			Myriogramme.		
			1 =	154,340.023	{ ʒb xxvi. ʒix ʒiv.

TABLE OF THE PROPORTION BY MEASURE OF ALCOHOL (SP. GR. 0.825)
CONTAINED IN 100 PARTS OF DIFFERENT WINES, ETC.*

Lisa (mean).....	25.41	Teneriffe (C.).....	16.61	Lunel.....	15.52
Raisin wine (mean)...	25.12	Colares.....	19.75	Ditto (F.).....	18.10
Marsala [Sicily ma-		Lachryma Christi.....	19.70	Sheraz.....	15.52
deira] (mean)...	25.09	White Constantia.....	19.75	Ditto (C.).....	15.56
strongest (J.).....	21.10	Red Constantia.....	18.92	Syracuse.....	15.28
weakest (J.).....	19.90	Lisbon.....	18.94	Sauterne.....	14.22
Port, strongest.....	25.83	Ditto (C.).....	19.09	Burgundy (mean).....	14.57
mean.....	22.96	Bucellas.....	18.49	strongest (J.).....	13.20
weakest.....	19.00	Red madeira (mean)...	20.35	weakest (J.).....	10.10
strongest (C.).....	20.49	Cape muschat.....	18.25	Hock (mean).....	12.08
mean (C.).....	18.68	Cape madeira (mean).	20.51	strongest (J.).....	13.00
weakest (C.).....	16.80	Grape wine.....	18.11	weakest (J.).....	9.50
strongest (J.).....	23.20	Calavella (mean).....	18.65	Nice.....	14.63
weakest (J.).....	20.70	Vidonia.....	19.25	Barsac.....	13.86
White port (C.).....	17.22	Alba flora.....	17.26	Tent.....	13.30
Madeira, strongest....	24.42	Zante.....	17.05	Champagne (mean)...	12.61
mean.....	22.27	Madaga.....	17.26	Ditto (F.).....	12.20
weakest.....	19.24	White hermitage.....	17.43	Ditto, strongest (J.)...	14.80
strongest (C.).....	20.35	Roussillon (mean).....	18.13	weakest (J.).....	14.10
strongest (J.).....	19.70	Claret (strongest).....	17.11	Red hermitage.....	12.32
weakest (J.).....	19.00	mean.....	15.10	Vin de Grave (mean)..	13.37
Sercial madeira.....	21.40	weakest.....	12.91	Frontignac (Rives	
Ditto (C.).....	18.50	ditto (F.).....	14.73	Altes).....	12.79
Sherry, strongest.....	19.81	vin-ordinaire (C.)	10.42	Ditto (C.).....	12.29
mean.....	19.17	Château-Latour,		Côte rôtie.....	12.32
weakest.....	18.25	1825 (C.).....	9.38	Tokay.....	9.88
strongest (C.).....	19.31	first growth, 1811		Rudesheimer, first	
mean (C.).....	18.47	(C.).....	9.32	quality (C.).....	10.14
weakest (C.).....	16.96	strongest (J.).....	11.10	inferior (C.).....	8.35
Amontillado (C.)..	15.18	weakest (J.).....	9.10	Hambacher, first qual.	
strongest (J.).....	24.70	Malmsey madeira.....	16.40	(C.).....	8.88
weakest (J.).....	15.40	Ditto (C.).....	15.60	Catawba (Stearns)...	8 to 11
Teneriffe.....	19.79				
Cider, highest average	9.87	Ale (Edinburgh).....	6.20	Brandy.....	53.39
lowest average....	5.21	Ale (Dorchester).....	5.56	Rum.....	53.68
Perry, average of four		Brown stout.....	6.80	Gin.....	51.60
samples.....	7.26	London porter.....	4.20	Scotch whisky.....	54.32
Mead.....	7.32	London small beer....	1.28	Irish whisky.....	53.90
Ale (Burton).....	8.88				

* The analyses whose results are given in this table were mostly made by Mr. Brande. When no mark is attached, the quotation is upon his authority. When the mark (F.) is added, the analysis was made by Julia-Fontenelle; (C.) by Professor Christison; (J.) by Dr. H. Bence Jones.

THERE have been omitted from the body of the present work, by an oversight, several preparations of such importance as to require notice. These are appended here.

LIQUOR PLUMBI SUBACETATIS.—The *Solution of the Subacetate of Lead*, or *Goulard's Extract*, as it is sometimes called, is a colorless, limpid liquid, of a sweetish, astringent taste. It is made by boiling litharge in a solution of the acetate of lead, and has an alkaline reaction. When exposed to the air, it rapidly absorbs carbonic acid and deposits the carbonate of lead, the neutral acetate being left in solution. In its action upon the human organism, Goulard's extract resembles very closely the simple acetate of lead; but it is never used internally. Externally, it is a favorite application in cases of sprains or bruises, as well as in superficial inflammation. For this purpose it requires dilution, and from a fluidounce to four fluidounces of it may be added to a pint of water. When used upon a raw surface, the strength should not be so great. A sedative poultice, which is very highly recommended by some physicians in the early stages of inflammation, may be made by saturating crumbs of stale bread with Goulard's extract diluted with four to six times its bulk of water. This poultice must be applied cold. The official *Liquor Plumbi Subacetatis Dilutus*, or Dilute Solution of the Subacetate of Lead, is of the strength of three fluidrachms to the pint. It is too weak to be of much value.

SYRUPUS FERRI IODIDI.—The *Syrup of the Iodide of Iron* is made by shaking iron, iodine, and water together, and adding hot syrup. It is a transparent, greenish liquid, of a sweet, ferruginous taste. It deposits no sediment on keeping, and should not affect the color of starch. If it strikes a blue color with the latter substance, it contains free iodine. The syrup of the iodide of iron is a favorite remedy in those cases of anæmia in which there is a distinct scrofulous taint. It is believed to possess the peculiar alterative powers of iodine, conjoined with the tonic properties of iron. It is enormously used in *scrofulosis* occurring in anæmic children; but it certainly possesses no advantages over a ferruginous tonic and iodine when given separately but simultaneously. Indeed, it seems to me uncertain whether its use is as effective as the separate administration of iodine and iron. The dose of it for a child two years old is five to ten drops; for an adult, thirty to forty minims. As it affects the teeth very seriously, it should always be freely diluted when taken, and the mouth should be well washed after its administration.

VINUM ANTIMONII.—*Antimonial Wine* consists of two grains of tartar emetic dissolved in an ounce of sherry wine. The emetic dose is half a fluidounce. The *Unguentum Antimonii* (tartar emetic, one part in five) and the *Emplastrum Antimonii* are only employed externally, as counter-irritants. A small quantity of the first of these preparations is spread upon a linen rag

and laid upon the skin, or a little of it may be well rubbed in twice a day. The plaster is used in the ordinary manner. Whenever either of these preparations is persistently used, there comes on, sooner or later, a peculiar burning or tingling pain, which is very shortly followed by pustulation. The effect of the drug is very persistent as well as severe, so that the remedy is only applicable to a few cases in which an action of the kind spoken of is required. Care must be exercised not to continue the application too long, lest severe and obstinate ulceration be produced.

THE officinal suppositories of the U. S. Pharmacopœia are as follows :

Suppositoria Acidi Carbolici, each containing one grain.

Suppositoria Acidi Tannici, each containing five grains.

Suppositoria Aloës, each containing five grains.

Suppositoria Assafœtidæ, each containing the extract derived from the evaporation of forty minims of the tincture of assafœtida.

Suppositoria Belladonnæ, each containing one-half grain of the alcoholic extract.

Suppositoria Morphicæ, each containing one-half grain.

Suppositoria Opii, each containing one grain.

Suppositoria Plumbi, each containing three grains.

Suppositoria Plumbi et Opii, each containing three grains of the sugar of lead and one-half grain of opium.

ERRATA.

IN the formulæ for Alum, page 30, for $(\text{SO}_4)_2\text{AlK} + 12\text{OH}_2$, read $(\text{SO}_4)_2\text{AlK} + 12\text{H}_2\text{O}$; and for $(\text{SO}_4)_2\text{AlNH}_4 + 12\text{OH}_2$, read $(\text{SO}_4)_2\text{AlNH}_4 + 12\text{H}_2\text{O}$.

On page 37, for Pb_2NO_3 read Pb_2NO_3 ; also, the formulæ for the subcarbonate of bismuth should be altered to read $\text{BiO}_3\text{CO}_2\text{HO} = 2\text{Bi}_2\text{O}_3\text{CO}_3\text{H}_2\text{O}$; of the subnitrate, to $\text{BiNO}_3\text{H}_2\text{O}$.

On page 39, for $\text{ZnC}_2\text{H}_4\text{O}_2$, read $\text{ZnC}_2\text{H}_3\text{O}_2$.

On page 102, for NH_5NO_3 , read NH_4NO_3 .

On page 237, for $4\text{HO},\text{OH}$ and 2NO , read $2\text{H}_2\text{O}$ and N_2O .

GENERAL INDEX.

A.

- Absorbents, 528.
 Acacia, 493.
 Aceta, 14.
 Acetate of lead, 36.
 of morphia, 194.
 of potassium, 415.
 of zinc, 39.
 Acetated tincture of opium, 194.
 Acetic acid, 169.
 extract of colchicum, 136.
 Acetum, 168.
 destillatum, 168.
 lobeliae, 304.
 opii, 194.
 sanguinariae, 367.
 scillae, 408.
 Acid, benzoic, 448.
 copaivic, 426.
 eubie, 428.
 gentisic, 51.
 meconic, 200.
 Acidum aceticum, 169.
 aceticum dilutum, 169.
 arseniosum, 311.
 benzoicum, 448.
 carbolicum, 140.
 chromicum, 492.
 citricum, 167.
 gallicum, 26.
 hydrocyanicum dilutum, 155.
 muriaticum, 85, 492.
 muriaticum dilutum, 85.
 nitricum, 85, 491.
 nitricum dilutum, 86.
 nitromuriaticum, 87.
 nitromuriaticum dilutum, 87.
 nitrosum, 86.
 oxalicum, 169.
 phosphoricum dilutum, 347.
 phosphoricum glaciale, 347.
 sulphuricum, 82, 491, 530.
 sulphuricum aromaticum, 84.
 sulphuricum dilutum, 84.
 sulphurosum, 540.
 Acidum tannicum, 24.
 tartaricum, 165.
 valerianicum, 173.
 Aconella, 148.
 Aconite leaves, 147.
 root, 147.
 Aconiti folia, 147.
 radix, 147.
 Aconitia, 148-155.
 antidote for, 125.
 Adhesive plaster, 36.
 Administration, method of, 16.
 Æther, 242.
 fortior, 242.
 African pepper, 75.
 Alcohol, 102.
 as a sudorific, 439.
 dilutum, 102.
 fortius, 102.
 Alcoholic extract of bella-donna, 231.
 extract of conium, 308.
 extract of hyoscyamus, 234.
 Allium, 451.
 Allspice, 74.
 Aloe, 388.
 Barbadensis, 388.
 Capensis, 389.
 purificata, 390.
 Socotrina, 388.
 Aloin, 389.
 Alteratives, 310.
 Alum, 30, 370.
 Alumen, 30.
 exsiccatum, 30.
 Aluminii et ammonii sulphas, 30.
 Aluminii et potassii sulphas, 30.
 sulphas, 31.
 Amber, 177.
 American centaury, 53.
 columbo, 53.
 hellebore, 135.
 hemp, 200.
 Ammonia, 99, 485.
 Ammonia-alum, 30.
 Ammoniac, 448.
 mixture, 448.
 plaster, 448.
 Ammoniacum, 448.
 Ammonia murias, 444.
 Ammoniated copper, 39.
 mercury, 333.
 tincture of guaiac, 358.
 tincture of valerian, 173.
 Ammonii bromidum, 286.
 carbonas, 101.
 chloridum, 444.
 nitras, 102.
 sulphas, 102.
 valerianas, 173.
 Ammonio-ferric alum, 82.
 Amygdalin, 53.
 Amyli nitritum, 297.
 Amylic alcohol, 297.
 Anæsthetics, 235.
 Analgesics, 181.
 Angustura, 73.
 Animal charcoal, 528.
 Anise, 76.
 water, 77.
 Anisum, 76.
 Antacids, 509.
 Anthelmintics, 511.
 Anthemis, 72.
 Antimonial ointment, 560.
 plaster, 560.
 wine, 560.
 Antimonii et potassii tartaras, 130.
 dose of, 444.
 oxidum, 130.
 sulphuretum, 129.
 Antizymotics, 539.
 Aperient effervescent powders, 394.
 Apiol, 455.
 Apomorphia, 368.
 Aqua ammoniæ, 101.
 ammoniæ fortior, 101.
 Aquæ, 14.
 piceis liquidæ, 452.
 Arabin, 494.
 Argenti cyanidum, 165.
 nitras, 42.
 nitras fusa, 42.
 oxidum, 49.
 Argentinum, 41.
 Argyria, 48.
 Aritia, 55.
 Aromatic bitters, 72.
 powder, 73.
 spirit of ammonia, 101.
 sulphuric acid, 84.
 syrup of rhubarb, 388.

Aromatics, 71.
 Arrow-root, 497.
 Arseniate of sodium, 322.
 Arsenic, 311.
 antidotes for, 321.
 eating, 315.
 Arsenical paper, compound, 319.
 paste, 489.
 Arsenici chloridum, 322.
 iodidum, 322.
 Arsenicum, 311.
 Arsenious acid, 311.
 ointment, 489.
 Arsenite of potassium, 322.
 Assafetida, 173.
 mixture, 174.
 Assafoetida, 173.
 Astringents, 23.
 vegetable, 24.
 Atomization, 17, 441.
 Atropia, 206.
 antagonism to Calabar bean, 278.
 Atropiæ sulphas, 231.
 Aurantii amari cortex, 76.
 dulcis cortex, 76.
 flores, 76.
 Azedarach, 517.

B.

Balm, 77.
 Balsam of Peru, 451.
 of Tolu, 451.
 Balsamum Peruvianum, 451.
 Tolutanum, 451.
 Barbadoes aloes, 388.
 Barberry, 52.
 Bark of pomegranate root, 30.
 Barley, 497.
 Barytina, 141.
 Bassorin, 494.
 Bean of Saint Ignatius, 266.
 Bearberry, 423.
 Belladonna leaves, 206.
 plaster, 231.
 root, 206.
 Belladonnæ folia, 206.
 radix, 206.
 Benzoic acid, 448.
 Benzoin, 448.
 Benzoinum, 448.
 Berberina, 397.
 Berberis, 52.
 Bicarbonate of potassium, 416.
 of sodium, 512.
 Bichloride of methylene, 253.
 Bismuth, 37.
 Bismuthi subcarbonas, 37.
 subnitras, 37.
 Bismuthum, 37.
 Bitartrate of potassium, 416.
 Bitter orange peel, 76.
 polygala, 76.
 Black drop, 194.
 ginger, 75.
 hellebore, 403.

Black mustard, 482.
 oak, 29.
 pepper, 75.
 snakeroot, 180.
 wash, 334.
 Blackberry, 30.
 Blistering cerate, 481.
 Blisters, 477.
 Bloodroot, 367.
 Blue mass, 329.
 ointment, 329.
 pills, 329.
 Boneset, 52.
 Brandy, 102.
 Brayera, 518.
 Bromide of ammonium, 286.
 of lithium, 286.
 of potassium, 278.
 of sodium, 285.
 Brominated camphor, 177.
 Bromine, 492.
 Brominium, 492.
 Broom, 409.
 Brown mixture, 496.
 Buchu, 422.
 Burgundy pitch, 485.
 pitch plaster, 485.
 Burnt alum, 492.
 Butternut, 388.

C.

Calabar bean, 267.
 Calamus, 77.
 Calcii carbonas præcipitata, 514.
 phosphas præcipitata, 349.
 Calisaya bark, 55.
 Calomel, 330.
 as a purge, 380.
 Calumba, 52.
 Calx, 513.
 chlorinata, 535.
 Camphor, 174.
 hydrobromate of, 177.
 mixture, Hope's, 86.
 water, 176.
 Camphora, 174.
 Camphorated tincture of opium, 193.
 Canada erigeron, 424.
 fleabane, 424.
 pitch, 486.
 pitch plaster, 486.
 turpentine, 425.
 Canella, 76.
 Cannabin, 201.
 Cannabis Americana, 200.
 Indica, 200.
 Canquoin's paste, 490.
 Cantharidal collodion, 481.
 Cantharides, 478.
 cerate, 481.
 paper, 481.
 Cantharidin, 478.
 Cantharis, 478.
 Cape aloes, 389.
 Capsicin, 75.
 Capsicum, 75, 484.

Caraway, 76.
 Carbo, 528.
 animalis, 528.
 animalis purificatus, 528.
 ligni, 528.
 Carbolic acid, 541.
 Carbonate of ammonium, 101.
 of lead, 36.
 of magnesium, 379.
 of potassium, 414.
 of sodium, 512.
 of zinc, 39.
 Cardamom, 74.
 Cardamomum, 74.
 Cardiac sedatives, 129.
 stimulants, 99.
 Carminatives, 71.
 Carota, 424.
 Carrageen, 495.
 Carrot seed, 424.
 Carthagena bark, 55.
 Carum, 76.
 Caryophyllus, 73.
 Cascarella, 75.
 Cassava, 496.
 Cassia fistula, 370.
 Castillon's powder, 515.
 Castor, 172.
 oil, 381.
 Castoreum, 172.
 Catechu, 27.
 Cayenne pepper, 75.
 Cerata, 15.
 Cerate of carbonate of zinc, 39.
 Cerates, 15.
 Ceratum cantharidis, 481.
 sabinæ, 454.
 zinci carbonatis, 39.
 Cetraria, 495.
 Cetraric acid, 495.
 Cetrarin, 495.
 Cevadilla, 142.
 Chalk, 514.
 mixture, 514.
 Chamomile, 72.
 Champagne, 113.
 Charcoal, 528.
 Charta cantharidis, 481.
 sinapis, 484.
 Chartæ, 15.
 Chelerythrin, 367.
 Chemical food, 349.
 Chenopodium, 517.
 Chimaphila, 423.
 Chimaphilin, 423.
 Chinoidine, 70.
 Chloral, 287.
 Chlorate of potassium, 420.
 Chloride of ammonium, 444.
 of iron, 81.
 of mercury, corrosive, 331.
 of mercury, mild, 331.
 of zinc, 490.
 Chlorinated lime, 535.
 Chlorine, 534.
 Chlorine water, 534.
 Chlorinum, 534.

- Chloroformum purificatum, 246.
 venale, 246.
 Chondrus, 495.
 Chromic acid, 492.
 Chrysophanic acid, 387.
 Churrus, 201.
 Cigarettes, arsenical, 319.
 Cimicifuga, 179.
 Cinchona, 54.
 flava, 55.
 pallida, 55.
 red, 55.
 rubra, 55.
 Cinchonia, 55.
 Cinchonæ sulphas, 70.
 Cinchonidia, 55.
 Cinnamomum, 73.
 Cinnamon, 73.
 water, 73.
 Cissampelin, 422.
 Citrate of iron, 82.
 of iron and ammonium, 82.
 of iron and quinia, 82.
 of iron and strychnia, 82.
 of potassium, 415.
 Citric acid, 167.
 Classification, 20.
 Cloves, 73.
 Codeia, 197.
 Cod-liver oil, 341.
 Cohnheim's salt frog, 292.
 Colchiceina, 350, 352.
 Colchici radix, 349.
 semen, 349.
 Colchicia, 350, 355.
 Colchicum root, 349.
 seed, 349.
 Cold cream, 499.
 Collodion, 506.
 with cantharides, 481.
 Collodium, 506.
 cum cantharide, 481.
 flexile, 507.
 Colocynthis, 396.
 Colocynthin, 396.
 Colocynthis, 396.
 Columbo, 52.
 Commercial bicarbonate of sodium, 512.
 chloroform, 246.
 oxide of zinc, 39.
 Compound cathartic pills, 397.
 decoction of sarsaparilla, 357.
 extract of colocynthis, 396.
 fluid extract of sarsaparilla, 357.
 infusion of catechu, 28.
 infusion of flaxseed, 496.
 infusion of gentian, 51.
 infusion of rose, 29.
 iodine ointment, 338.
 mixture of iron, 80.
 mixture of liquorice, 496.
 pill of soap, 193.
 Compound pills of iron, 80.
 pills of rhubarb, 388.
 powder of ipecacuanha, 436.
 powder of jalap, 395.
 powder of rhubarb, 388.
 solution of iodine, 338.
 spirit of ether, 178.
 spirit of juniper, 424.
 spirit of lavender, 77.
 syrup of sarsaparilla, 357.
 syrup of squill, 451.
 tincture of benzoin, 450.
 tincture of cardamom, 74.
 tincture of cinchona, 71.
 tincture of gentian, 51.
 tincture of iodine, 338.
 Confection of opium, 193.
 of orange peel, 76.
 of rose, 29.
 of senna, 392.
 Confectiones, 14.
 Confections, 14.
 Conia, 305.
 Conii folia, 305.
 fructus, 305.
 Conium leaves, 305.
 seed, 305.
 Copaiba, 426.
 Copaivic acid, 426.
 Copper, 40.
 Coptis, 53.
 Coriander, 76.
 Coriandrum, 76.
 Cornus circinata, 53.
 Florida, 53.
 sericea, 53.
 Corrosive chloride of mercury, 331.
 sublimite, 331, 491.
 Corsican moss, 495.
 Counter-irritation, 474.
 Cowhage, 522.
 Cox's hive syrup, 452.
 Cranesbill, 29.
 Cream of tartar, 416.
 Creasote, 541.
 Creasotum, 541.
 Creta, 514.
 præparata, 514.
 Croton oil, 402.
 chloral hydrate, 296.
 Crotonol, 402.
 Cubeb, 428.
 Cubeba, 428.
 Cubebic acid, 428.
 Cubebin, 428.
 Cupri subacetate, 41.
 sulphas, 40, 370.
 Cuprum, 40.
 ammoniatum, 41.
 Cyanide of potassium, 165.
 of silver, 165.
 Cydonium, 496.
- D.**
- Dandelion, 359.
 Daphnin, 359.
 Datura, 233.
 Decoction, 14.
 Decoction of barley, 498.
 of seneka, 447.
 Decoction sarsaparillæ compositum, 357.
 sengæ, 447.
 uvæ ursi, 423.
 Delphinia, antidote for, 125.
 Deodorized tincture of opium, 194.
 Depresso-motors, 267.
 Dewee's emmenagogue mixture, 457.
 Diaphoretics, 431.
 Digestants, 524.
 Digitalin, 116.
 Digitalinum, 116.
 Digitalis, 116.
 as a diuretic, 408.
 Diluents, 504.
 Diluted acetic acid, 169.
 alcohol, 102.
 hydrocyanic acid, 155.
 muriatic acid, 85.
 nitric acid, 86.
 nitromuriatic acid, 87.
 phosphoric acid, 347.
 solution of subacetate of lead, 560.
 sulphuric acid, 84.
 Diospyros, 30.
 Disinfectants, 530.
 Distilled vinegar, 168.
 Diuretics, 404.
 Dogwood, 53.
 Donovan's solution, 340.
 Doses, rules for, 18.
 Dover's powder, 436.
 Dried alum, 30.
 carbonate of sodium, 512.
 sulphate of iron, 81.
- E.**
- Echolina, 459.
 Effects of medicines, primary, 15.
 of medicines, remote, 15.
 Effervescing draught, 415.
 powders, 512.
 Elaterin, 399.
 Elaterium, 399.
 Elixir of vitriol, 84.
 Emetia, 363.
 Emetics, 360.
 Emmenagogues, 453.
 tonic, 453.
 Emollients, 499.
 Emplastrum, 15.
 Emplastrum belladonnæ, 231.
 plumbi, 36.
 resinæ, 36.
 saponis, 36.
 Emulsin, 53, 483.
 Emulsions, 53.
 Endermic administration, 17.
 Epispastics, 474.
 Epsom salts, 392.
 Ergot, 458.

Ergota, 458.
 Ergotin, 459.
 Ergotism, 469.
 Erigeron, 424.
 Canadense, 424.
 Errhines, 473.
 Escharotics, 487.
 Eserina, 267.
 Ether, 242.
 Ethereal oil, 178.
 Ethyl oxide, 242.
 Eupatorium, 52.
 Exeito-motors, 254.
 Expectorants, 440.
 Extract of aconite, 155.
 of American hemp, 205.
 of belladonna, 231.
 of black hellebore, 403.
 of butternut, 388.
 of Calabar bean, 267.
 of cinchona, 71.
 of colocynth, 396.
 of conium, 306.
 of dandelion, 359.
 of digitalis, 128.
 of gentian, 51.
 of hyoseyamus, 234.
 of ignatia, 266.
 of Indian hemp, 200.
 of jalap, 396.
 of liquorice, 496.
 of logwood, 28.
 of May-apple, 38.
 of nux vomica, 254.
 of opium, 193.
 of quassia, 21.
 of rhatany, 29.
 of rhubarb, 388.
 of stramonium leaves, 233.
 of stramonium seed, 233.
 of taraxacum, 359.
 of uva ursi, 423.
 of valerian, 172.
 Extracta, 14.
 fluida, 14.
 Extractum cannabis Ameri-
 cane, 200.
 cannabis Indicæ, 200.
 hæmatoxyli, 28.
 juglandis, 388.

F.

Fennel, 76.
 water, 76.
 Ferri citras, 82.
 et ammonii citras, 82.
 et ammonii sulphas, 82.
 et ammonii tartras, 82.
 et potassii tartras, 82.
 et quiniæ citras, 82.
 et strychniæ citras, 82.
 ferrocyanidum, 82.
 lactas, 82.
 oxidum hydratum, 80.
 phosphas, 82.
 pulvis, 79.
 pyrophosphas, 82.
 subcarbonas, 80.

Ferri sulphas, 81.
 sulphas exsiccata, 81.
 Ferrocyanide of iron, 82.
 Ferrum, 77.
 redactum, 79.
 Figs, 378.
 Filix mas, 522.
 Flaxseed, 496.
 meal, 496.
 oil, 496.
 Fleabane, 424.
 Flexible collodion, 507.
 Fluid extract of American
 hellebore, 140.
 of belladonna root, 233.
 of buchu, 422.
 of cimicifuga, 179.
 of cinchona, 71.
 of colchicum root, 355.
 of colchicum seeds, 356.
 of columbo, 52.
 of conium seed, 308.
 of dandelion, 359.
 of ergot, 471.
 of gentian, 52.
 of ginger, 74.
 of ipecacuanha, 367.
 of lupulin, 179.
 of pareira brava, 423.
 of rhatany, 29.
 of rhubarb, 388.
 of sarsaparilla, 357.
 of senna, 392.
 of serpentaria, 73.
 of spigelia, 517.
 of spigelia and senna, 517.
 of squill, 408.
 of valerian, 173.
 of veratrum viride, 140.
 of wild-cherry, 54.
 Fœniculum, 76.
 Fowler's solution, 322.
 Fraxea, 53.
 Fumigations, mercurial, 329.
 Fusel oil, 297.

G.

Gaduin, 342.
 Galactagogue, 382.
 Galla, 27.
 Gallic acid, 26.
 Galls, 27.
 Gambia, 27.
 Gamboge, 401.
 Gambogia, 401.
 Garlic, 451.
 Gaultheria, 77.
 Gentian, 51.
 Gentiana, 51.
 Gentiopikrin, 51.
 Gentsic acid, 51.
 Geranium, 29.
 German chamomile, 72.

Gillenia, 368.
 Ginger, 74.
 Glacial phosphoric acid, 347.
 Glauber salts, 393.
 Glycerina, 499.
 Glycerine, 499.
 Glycerita, 14.
 Glycerite of gallic acid, 27.
 of tannic acid, 26.
 Glycyrrhiza, 496.
 Goldthread, 53.
 Goose-grease, 499.
 Goulard's extract, 560.
 Granati fructus cortex, 30.
 radicis cortex, 30.
 Gray powder, 329.
 Green iodide of mercury, 333.
 Guaiac, 357, 457.
 yellow, 358.
 Guaiaci lignum, 357.
 resinæ, 357.
 Guaiacic acid, 358.
 Guaiacin, 358.
 Guaiaconic acid, 358.
 Guaiacresinic acid, 358.
 Guaiacum wood, 357.
 Gum arabic, 493.
 Gun cotton, 506.
 Gunjah, 201.
 Gutta-percha, 508.

H.

Hæmatin, 28.
 Hæmatoxylin, 28.
 Hæmatoxylin, 28.
 Hashish, 201.
 Helleborein, 403.
 Helleborin, 403.
 Helleborus, 403.
 Hemlock leaves, 305.
 pitch, 486.
 pitch plaster, 486.
 Hiera pieræ, 76.
 Hoffman's anodyne, 178.
 Honey, 14.
 Hops, 178.
 Hordeum, 497.
 Humulus, 178.
 Huxham's tincture, 71.
 Hydragogues, 375.
 Hydrargyri chloridum corro-
 sivum, 491.
 chloridum mite, 330.
 iodidum rubrum, 333.
 iodidum viride, 333.
 oxidum flavum, 333.
 oxidum rubrum, 333.
 sulphas flava, 333.
 sulphuretum rubrum, 333.
 Hydrargyrum, 322.
 ammoniatum, 333.
 cum creta, 329.
 Hydrate of chloral, 287.
 Hydrated oxide of iron, 80.
 Hyoseyami folia, 233.
 Hyoseyami semen, 233.

Hyoscyamia, 233.
Hyoscyamus leaves, 233.
 seed, 233.
Hypodermic injections, 16.
 injections of calomel, 331.

I.

Ice-land moss, 195.
Ignatia, 266.
 Indian hemp, 300.
 Indications for the use of medicines, 15.
Infusa, 14.
 Infusion of *angustura*, 73.
 of *buchu*, 422.
 of *capsicum*, 75.
 of *cascarilla*, 73.
 of *chamomile*, 72.
 of *cloves*, 73.
 of *columbo*, 52.
 of *digitalis*, 128.
 of *ginger*, 74.
 of *hops*, 179.
 of *juniper*, 424.
 of *pareira brava*, 423.
 of *quassia*, 51.
 of *red cinchona*, 71.
 of *rhatany*, 29.
 of *rhubarb*, 388.
 of *senna*, 292.
 of *serpentaria*, 72.
 of *spigelia*, 517.
 of *valerian*, 173.
 of *wild-cherry*, 54.
 Inunctions, mercurial, 328.
 Iodide of arsenic, 322.
 of potassium, 338.
Iodine, 334.
 ointment, 338.
Iodinium, 334.
Iodoform, 340.
Iodoformum, 340.
Ipecacuanha, 444.
 Irish moss, 495.
 Iron, 77.
 by hydrogen, 79.
 Quevenne's, 79.

J.

Jalap, 395.
Jalapa, 395.
 Jamaica ginger, 74.
Jervina, 141.
Juglans, 388.
 Juice of *conium*, 308.
 of *dandelion*, 359.
Juices, 14.
Juniper, 424.
Juniperus, 424.

K.

Kamecla, 523.
 Kentish ointment, 485.
Kino, 28.
Kinovic acid, 55.
Koossin, 518.

Koosso, 518.
Krameria, 29.

L.

Labarraque's solution, 535.
 Lactate of iron, 82.
Laudanum, 193.
 deodorized, 194.
Lavandula, 77.
 Lavender, 77.
 Lead, 32.
 acetate of, 36.
 carbonate of, 36.
 nitrate of, 37.
 plaster, 36.
 poisoning, 32.
 water, 36.
Ledoyen's disinfectant solution, 37.
 Lemon-juice, 168.
 Lemons, essential salt of, 169.
Lichenin, 495.
Lignum-vitæ, 357.
Limæ, 513.
 lacto-phosphate of, 349.
 liniment, 514.
 water, 514.
Limonia succus, 168.
Lini farina, 496.
 Liniment of ammonia, 485.
 of camphor, 176.
 of turpentine, 485.
Linimenta, 15.
Linimentum saponis, 177.
Linum, 496.
Liquor ammonii acetatis, 437.
 arsenici chloridi, 322.
 arsenici et hydrargyri iodidi, 340.
 bismuthi et ammonii citratis, 38.
 calcis, 514.
 ferri subsulphatis, 81.
 ferri tersulphatis, 81.
 gutta-perchæ, 508.
 hydrargyri nitratis, 491.
 iodinii compositus, 338.
 magnesi citratis, 393.
 plumbi subacetatis, 560.
 plumbi subacetatis dilutus, 560.
 potassæ, 488.
 potassii arsenitis, 322.
 potassii citratis, 415.
 sodæ chlorinata, 50.
 sodii arseniatis, 322.
 zinci chloridi, 491.
Liquores, 14.
Liquorice, 496.
 root, 496.
Litharge, 35.
Lobelia, 303, 443.
Lobelic acid, 304.
Lobelina, 304.
Logwood, 28.
 Lugol's solution, 338.
Lupulin, 178.
Lupulina, 178.
Lyctonia, 148.

M.

Mace, 73.
Macis, 73.
 Magendie's solution, 195.
Magnesia, 379.
Magnesi carbonas, 379.
 sulphas, 392.
 Male fern, 522.
Manna, 378.
Maranta, 497.
Materia medica, 13.
Maticin, 430.
Matico, 430.
Matricaria, 72.
 May-apple, 397.
 Meadow saffron, 349.
Measures, 555.
Meconia, 200.
Mecconic acid, 182, 200.
Medulla sassafras, 496.
Mellita, 14.
Mentha piperita, 77.
 viridis, 77.
Mercurial ointment, 329.
 plaster, 392.
 purgatives, 383.
Mercury, 322.
 ammoniated, 333.
 with chalk, 329.
Methenyl chloride, 246.
Methylene bichloride, 253.
Mezereon, 359.
 ointment, 359.
Mezereum, 359.
 Mild chloride of mercury, 330.
 Milk of assafetida, 174.
Misturæ, 14.
 Mixture of assafetida, 174.
 of citrate of potassium, 415.
Molasses, 377.
 Monsef's solution, 81.
Morphia, 194.
Morphiæ acetas, 194.
 murias, 194.
 sulphas, 194.
Moschus, 171.
Mucilage of gum arabic, 494.
 of sassafras pith, 496.
 of slippery-elm bark, 494.
 of tragacanth, 494.
Mucilagines, 14.
Mucuna, 522.
 Muriate of ammonia, 444.
 of morphia, 194.
Muriatic acid, 85.
Muscaria, antidote for, 125.
Musk, 171.
Mustard, 482.
 as an emetic, 369.
 paper, 484.
 Mutton suet, 499.
Mydriasis, 218.
Mydriatics, 206.
Myristica, 73.
Myronic acid, 483.
Myrrh, 452.
Myrrha, 452.

N.

Napellina, 148.
 Narcein, 195.
 Narcotina, 198.
 Neutral mixture, 415.
 Nitrate of ammonium, 102.
 of calcium, 417.
 of lead, 37.
 of potassium, 417.
 of silver, 42.
 of sodium, 513.
 Nitre, 417.
 Nitric acid, 85.
 Nitrite of amyl, 297.
 Nitrites, action of, on the blood, 301.
 Nitrogen monoxide, 237.
 Nitromuriatic acid, 87.
 Nitrous oxide, 237.
 Nutgall, 27.
 Nutmeg, 73.
 Nux vomica, 254.

O.

Oil of amber, 177.
 of anise, 76.
 of cajeput, 76.
 of Canada erigeron, 424.
 of caraway, 76.
 of cinnamon, 73.
 of cloves, 73.
 of copaiba, 428.
 of cubeb, 430.
 of fennel, 76.
 of gaultheria, 77.
 of juniper, 424.
 of lavender, 77.
 of mustard, 483.
 of nutmeg, 74.
 of peppermint, 77.
 of pimento, 74.
 of rosemary, 77.
 of rue, 455.
 of sassafras, 359.
 of savine, 454.
 of spearmint, 77.
 of succinum, 177.
 of succinum, rectified, 177.
 of turpentine, 114, 425, 484.
 of valerian, 172.
 of wormseed, 517.

Ointment, 15.

 of ammoniated mercury, 333.
 of belladonna, 231.
 of benzoin, 450.
 of carbonate of lead, 36.
 of galls, 27.
 of nutgall, 27.
 of oxide of zinc, 39.
 of red iodide of mercury, 333.
 of red oxide of mercury, 333.
 of rose-water, 499.
 of stramonium, 233.

Ointment of tannic acid, 26.
 of veratria, 147.
 of yellow oxide of mercury, 333.
 Olea destillata, 14.
 Oleoresin of black pepper, 75.
 of capsicum, 76.
 of cubeb, 430.
 of fern, 522.
 of ginger, 74.
 of lupulin, 179.
 Oleoresinæ, 14.
 Oleum phosphoratum, 97.
 Opianine, 200.
 Opium, 181.
 poisoning by, 191.
 Orange flowers, 76.
 flower water, 76.
 Oxalate of iron, 82.
 Oxalic acid, 169.
 Oxide of antimony, 130.
 of iron, hydrated, 80.
 of lead, 35.
 of silver, 49.
 of zinc, 39.
 Oxytocics, 458.
 Oyster-shell, 514.

P.

Pale cinchona, 55.
 rose, 29.
 Papaverina, 199.
 Paregoric, 193.
 Pareira, 422.
 brava, 422.
 Parsley, 455.
 Pellitory, 472.
 Pepo, 522.
 Peppermint, 77.
 water, 77.
 Pepsin, 525.
 Pepsina, 525.
 Permanganate of potassium, 133.
 Persimmon, 30.
 Petroselinum, 455.
 Pharmacology, 13.
 Pharmacopœia, 13.
 Pharmacy, 13.
 Phenic acid, 541.
 Phenylc alcohol, 541.
 Phosphate of calcium, 347.
 of iron, 82.
 of sodium, 393.
 Phosphoric acid, 347.
 Phosphorus, 89.
 antidote to, 96.
 Physostigma, 267.
 Physostigma, 267.
 Pill of carbonate of iron, 80.
 Pills of aloes, 391.
 of aloes and assafetida, 391.
 of aloes and mastic, 391.
 of aloes and myrrh, 391.
 of assafetida, 174.
 of copaiba, 428.
 of mercury, 329.

Pills of opium, 193.
 Pilula ferri carbonatis, 80.
 saponis composita, 193.
 Pilulæ, 15.
 catharticæ compositæ, 397.
 Pimenta, 74.
 Pimento, 74.
 Pinkroot, 516.
 Piper, 75.
 Piperin, 75.
 Pipsissewa, 423.
 Pix Burgundica, 485.
 Canadensis, 486.
 liquida, 452.
 Plaster of ammoniac with mercury, 448.
 of Canada pitch, 486.
 of pitch with cantharides, 485.
 Plumbi acetas, 36.
 carbonas, 36.
 nitras, 37.
 oxidum, 35.
 Plumbum, 32.
 Podophyllin, 398.
 Podophyllum, 397.
 Polygalic acid, 447.
 Pomegranate rind, 30, 522.
 Poppy, 181.
 Porphyroxin, 367, 199.
 Port wine, 102.
 Potassa, 488.
 Potassa cum calce, 489.
 Potassii acetas, 415.
 bicarbonas, 415.
 bitartas, 416.
 bromidum, 278.
 carbonas, 414.
 carbonas impura, 414.
 carbonas pura, 414.
 chloras, 420.
 citras, 415.
 cyanidum, 165.
 et sodii tartas, 394.
 iodidum, 338.
 nitras, 417.
 permanganas, 533.
 sulphas, 417.
 sulphuretum, 381.
 tartas, 417.
 Potassium, 411.
 Potentilla tormentilla, 30.
 Poutices, 501.
 Powder of aloes and canella, 76.
 Precipitated carbonate of calcium, 514.
 carbonate of iron, 80.
 carbonate of zinc, 39.
 phosphate of calcium, 349.
 sulphur, 380.
 Prepared chalk, 514.
 oyster-shell, 514.
 Propenyl alcohol, 500.
 Protectives, 506.
 Prunus Virginianæ, 53.
 Prussic acid, 155.
 Pseudacantha, 148.

Pseudomorphine, 200.
 Puccin, 367.
 Pulveres, 15.
 effervescentes, 512.
 effervescentes aperientes, 394.
 Pulvis aromaticus, 73.
 ipeacacuanhæ compositus, 436.
 Pumpkin-seed, 522.
 Purgative enemas, 454.
 Purgatives, 375, 381.
 Purgings cassia, 370.
 Pyrethrum, 472.
 Pyrophosphate of iron, 82.
 Pyroxylin, 507.
 Pyroxylon, 507.

Q.

Quassia, 51.
 Quassin, 51.
 Quercitron, 29.
 Quercus alba, 29.
 tinctoria, 29.
 Quevenne's iron, 79.
 Quince-seed, 496.
 Quinia, 55.
 as an abortifacient, 65.
 dihydroxyle, 65.
 Quiniæ sulphas, 55.
 Quinidia, 55.

R.

Rectified oil of amber, 177.
 Red cinchona, 55.
 iodide of mercury, 333.
 oxide of mercury, 333.
 precipitate, 333.
 rose, 29.
 sulphuret of mercury, 333.
 Reduced iron, 79.
 Resin, 113.
 of jalap, 395.
 of May-apple, 398.
 of scammony, 397.
 plaster, 36, 113.
 Resinæ, 14.
 Rhabarbin, 387.
 Rhatany, 29.
 Rhein, 387.
 Rheum, 386.
 Rhodeoretin, 395.
 Rhubarb, 386.
 Rhus glabrum, 421.
 Rochelle salt, 394.
 Roman chamomile, 72.
 Rosa centifolia, 29.
 Gallica, 29.
 Rosemary, 77.
 Rose-water, 29.
 Rosmarinus, 77.
 Rottlera, 523.
 Rottlerin, 523.
 Rubefacients, 482.
 Rubus, 30.
 Rue, 455.
 Ruta, 455.

S.

Sabadilla, 142.
 Sabbatia, 53.
 Sabina, 454.
 Saccharate of lime, 549.
 Sage, 77.
 Sago, 497.
 Sal ammoniac, 444.
 Salicin, 53.
 Salix, 53.
 Salseparin, 356.
 Salt of sorrel, 169.
 Salvia, 77.
 Sanguinaria, 367.
 Sanguinarina, 367.
 Santonate of soda, 522.
 Santonica, 518.
 Santonin, 518.
 Santoninum, 518.
 Saponin, 447.
 Sarsaparilla, 356.
 Sarsaparillin, 356.
 Sassafras, 359.
 pith, 496.
 Savine, 454.
 Scammonium, 396.
 Scammony, 396.
 Scilla, 406, 451.
 Scillitin, 406.
 Sclerotium, 458.
 Scoparius, 409.
 Scudamore's mixture, 354.
 Sculein, 406.
 Seidlitz powders, 394.
 Senega, 447.
 Seneka, 447.
 Senna, 391.
 Septfoil, 30.
 Serpentaria, 72.
 Sesquichloride of iron, 82.
 Sherry wine, 102.
 Shore oil, 341.
 Sialagogues, 472.
 Silver, 39.
 Simaruba, 51.
 Sinapis alba, 482.
 nigra, 482.
 Slippery-elm bark, 494.
 Smilacin, 356.
 Snake-root, 179.
 Soap liniment, 176.
 plaster, 36.
 Socotrine aloes, 388.
 Soda, 512.
 powders, 512.
 Sodii acetat, 513.
 arsenias, 322.
 bicarbonas, 512.
 bicarbonas venalis, 512.
 bromidum, 285.
 carbonas, 512.
 nitras, 513.
 phosphas, 393.
 sulphas, 393.
 Sodium, 510.
 Solution of acetate of ammonium, 437.
 Solution of arsenite of sodium, 322.

Solution of arsenite of potassium, 322.
 of chloride of arsenic, 322.
 of chloride of zinc, 491.
 of chlorinated soda, 535.
 of citrate of potassium, 415.
 of gutta-percha, 508.
 of iodide of arsenic and mercury, 340.
 of lime, 513.
 of nitrate of mercury, 491.
 of permanganate of potassium, 533.
 of persulphate of iron, 81.
 of potassa, 488.
 of soda, 512.
 of subacetate of lead, 560.
 of subsulphate of iron, 81.
 of sulphate of morphia, 195.
 of tersulphate of iron, 81.
 Spanish flies, 478.
 Spartein, 409.
 Spearmint, 77.
 water, 77.
 Spice plasters, 484.
 Spigelia, 516.
 Spirit of ammonia, 101.
 of anise, 76.
 of camphor, 176.
 of cinnamon, 73.
 of juniper, 424.
 of lavender, 77.
 of Mindererus, 437.
 of nitrous ether, 437.
 of peppermint, 77.
 of spearmint, 77.
 Spiritus, 14.
 ætheris compositus, 178.
 ætheris nitrosi, 410, 437.
 frumenti, 102.
 vini Gallici, 102.
 Squill, 406.
 as an emetic, 369.
 as an expectorant, 451.
 St. Ignatius' bean, 266.
 Star anise, 76.
 Sticking plaster, 36.
 Straits oil, 341.
 Stramonii folia, 231.
 semen, 231.
 Stramonium leaves, 231.
 ointment, 203.
 seed, 231.
 Stronger alcohol, 102.
 ether, 242.
 water of ammonia, 101.
 Strychnia, 254.
 Strychniæ sulphas, 254.
 Subacetate of copper, 41.
 Subcarbonate of bismuth, 37.
 Subcarbonate of iron, 80.
 Sublimed sulphur, 380.

Subnitrate of bismuth, 37.
 Succci, 14.
 Succinum, 177.
 Succus conii, 308.
 Sugar of lead, 36.
 Sulphate of aluminium, 31.
 of aluminium and potassium, 30.
 of ammonium, 102.
 of atropia, 231.
 of cinchona, 70.
 of copper, 40.
 of iron, 81, 536.
 of iron and ammonium, 30.
 of magnesium, 392.
 of morphia, 194.
 of potassium, 417.
 of quinia, 55.
 of sodium, 393.
 of strychnia, 266.
 of zinc, 39.
 Sulpho-sinapisin, 483.
 Sulphur, 380.
 lotum, 380.
 ointment, 381.
 precipitatum, 380.
 sublimatum, 380.
 Sulphurated antimony, 129.
 Sulphuret of antimony, 129.
 of potassium, 381.
 Sulphuric acid, 82, 530.
 Sulphurous acid, 540.
 Sumach, 421.
 Suppositoria, 15.
 Suppositories of aloes, 561.
 of assafetida, 561.
 of belladonna, 561.
 of carbolic acid, 561.
 of lead, 561.
 of lead and opium, 561.
 of morphia, 561.
 of opium, 561.
 of tannic acid, 561.
 Sweet spirit of nitre, 410, 437.
 Syrup, 14.
 of acacia, 494.
 of ginger, 74.
 of iodide of iron, 560.
 of ipecacuanha, 367.
 of red rose, 29.
 of rhatany, 29.
 of rhubarb, 388.
 of sarsaparilla, compound, 349.
 of seneka, 447.
 of squill, 408, 451.
 of the phosphates, compound, 357.
 of Tolu, 451.
 of wild-cherry, 54.
 Syrupi, 14.

T.

Table of relation of weights and measures of the U. S. Pharmacopœia to each other, 556.

Table of solvent power of glycerine, 500.
 Table of weights and measures of the metrical system, 555.
 Table of weights and measures of the U. S. Pharmacopœia, 555.
 Tamarind, 378.
 Tamarindus, 378.
 Tannic acid, 24.
 Tapioca, 496.
 Tar, 452.
 ointment, 452.
 water, 452.
 Taraxacum, 359.
 Tartar emetic, 130, 370, 444.
 Tartaric acid, 165.
 Tartrate of antimony and potassium, 130.
 of iron and ammonium, 82.
 of iron and potassium, 82.
 of potassium, 417.
 of potassium and sodium, 394.
 Tenuin, 518.
 Tereb chloride of formyl, 246.
 Terebinthina, 103, 425.
 Canadensis, 425.
 Testa, 514.
 preparata, 514.
 Thebaia, 199.
 Therapeutics, 13.
 Tinctura opii, 193.
 opii acetata, 194.
 opii camphorata, 194.
 opii deodorata, 194.
 Tincture of aconite root, 185.
 of assafetida, 174.
 of belladonna, 231.
 of benzoin, 450.
 of black hellebore, 403.
 of bloodroot, 368.
 of cantharides, 478.
 of capsicum, 76.
 of cardamom, 74.
 of castor, 172.
 of catechu, 28.
 of chloride of iron, 81.
 of cinchona, 71.
 of cinnamon, 73.
 of colchicum, 355.
 of columbo, 52, 308.
 of conium, 306.
 of cubeb, 430.
 of digitalis, 128.
 of galls, 27.
 of ginger, 74.
 of guaiac, 358.
 of hemp, 205.
 of hops, 179.
 of hyoscyamus, 234.
 of iodine, 338.
 of kino, 28.
 of lobelia, 304.
 of lupulin, 179.
 of nutgall, 27.
 of nux vomica, 254.

Tincture of opium, 193.
 of opium, acetated, 194.
 of quassia, 51.
 of rhubarb, 388.
 of rhubarb and senna, 388.
 of sanguinaria, 368.
 of serpentaria, 72.
 of squill, 408.
 of stramonium, 233.
 of Tolu, 451.
 of valerian, 173.
 of valerian, ammoniated, 173.
 of veratrum viride, 140.
 Tonics, 50.
 Tormentil, 30.
 Tormentilla, 30.
 Tragacanth, 494.
 Tragacantha, 494.
 Trinitro-cellulose, 506.
 Troches, 15.
 Turpentine, 425, 113.
 Turpeth mineral, 333.

U.

Ulmus, 494.
 fulva, 494.
 Unguenta, 15.
 Urea, elimination of, by skin, 425.
 Ursin, 423.
 Uva ursi, 423.

V.

Valerian, 172.
 Valeriana, 172.
 Valerianate of ammonium, 173.
 Valerianic acid, 173.
 Vegetable acids, 165.
 Veratria, 142.
 ointment, 147.
 Veratroidia, 136.
 Veratrum album, 140.
 viride, 135.
 viride, resin of, 137.
 Verdigris, 41.
 Vesicatories, 477.
 Vienna paste, 489.
 Vina, 14.
 Vinegar, 168.
 distilled, 168.
 of bloodroot, 367.
 of lobelia, 304.
 of opium, 194.
 of squill, 408.
 Vinum antimonii, 560.
 portense, 102.
 Xericum, 102.
 Virginia snakeroot, 72.
 Viridia, 136.
 Vitriolated tartar, 394.

W.

Warming plaster, 485.
 Washed sulphur, 380.
 Water of ammonia, 101.

Weights and measures of the
metrical system, 555.
and measures of the
United States Pharmacopœia, 555.
Whisky, 102.
White arsenic, 311.
hellebore, 140.
lead, 36.
mustard, 482.
oak, 29.
pepper, 75.
precipitate, 334.
precipitate ointment,
334.
turpentine, 113.

White vitriol, 39.
Wild carrot, 424.
cherry, 53.
Wine of aloes, 391.
of colchicum root, 355.
of colchicum seed, 355.
of ergot, 471.
of ipecacuanha, 307.
of lobelia, 443.
of opium, 194.
Wormseed, 517.

Y.

Yellow cinchona, 55.

Yellow oxide of mercury,
333.
sulphate of mercury, 333.

Z.

Zinc, 39.
Zinci acetat, 39.
carbonas præcipitata, 39.
chloridum, 490.
oxidum, 39.
oxidum venale, 39.
sulphas, 39.
Zincum, 39.
Zingiber, 74.

INDEX OF DISEASES.

A.
Abscess :
 tannic acid, 26.
 alcohol, 110.
 carbolic acid, 541.
Acne :
 arsenic, 318.
 iodine, 337.
Acne rosacea :
 solution of nitrate of
 mercury, 491.
Adynamic fevers :
 ammonia, 100.
 alcohol, 110.
 digitalis, 126.
 camphor, 176.
 opium, 190.
 chloride of potassium,
 421.
 acetate of ammonium,
 437.
 spirit of nitrous ether,
 438.
Alcoholic intoxication :
 ammonia, 101.
Amaurosis :
 strychnia, 261.
Amenorrhœa :
 ginger, 74.
 carbonate of iron, 80.
 aloes, 390.
 turpentine, 426.
 Dover's powder, 436.
 alcohol, 439.
 seneka, 447.
 emmenagogues, 453.
 myrrh, 453.
 aloes, 454.
 hellebore, 454.
 savine, 454.
 rue, 455.
 apiol, 456.
 guaiac, 457.
Anæmia :
 iron, 77.
Aneurism :
 digitalis, 125.
 ergot, 467.
Angina pectoris :
 nitrite of amyl, 302.
 chlorate of potassium,
 421.
Animal poisoning :
 ammonia, 100.

Animal poisoning :
 alcohol, 110.
Aortic disease :
 digitalis, 124.
Ascites :
 elaterium, 400.
 chlorate of potassium,
 421.
Asthma :
 belladonna, 223.
 stramonium, 232.
 anæsthetics, 237.
 ether, 246.
 chloral, 294.
 nitrite of amyl, 302.
 lobelia, 304.
Atheroma :
 digitalis, 125.

B.
Bed-sores :
 protectives, 506.
Bilious fever :
 quinia, 69.
 calomel, 385.
 podophyllum, 398.
Biliousness :
 nitro-muriatic acid, 88.
 calomel, 385.
 potassium salts, 416.
Blood, determination of, to the
head, 400.
Bright's disease :
 gallic acid, 27.
 tincture of the chloride
 of iron, 81.
 diuretics, 404.
 diaphoretics, 436.
 water, 504.
Bronchial catarrh :
 alteratives, 443.
 benzoic acid, 450.
Bronchitis :
 oil of turpentine, 115.
 tartar emetic, 134, 444.
 cimicifuga, 180.
 opium, 190.
 lobelia, 304.
 sanguinaria, 368.
 copaiba, 428.
 cubebs, 429.
 expectorants, 440.
 astringents, 442.

Bronchitis :
 ipecacuanha, 444.
 chloride of ammonium,
 446.
 seneka, 447.
 ammoniac, 448.
 garlic, 451.
 squill, 451.
 tar, 452.
 Burgundy pitch, 485.
 demulcents, 493.
 liquorice, 496.
Bronchorrhœa :
 gallic acid, 27.
 alum, 31.
 astringents, 442.
Buboes :
 iodoform, 341.

C.
Cancer :
 iodoform, 337.
Carcinoma oris :
 nitric acid, 86.
Carcinoma of the stomach :
 bismuth, 38.
Cardiac disease, chronic :
 digitalis, 123.
 veratrum viride, 139.
Cardiac dropsy :
 squill, 407.
 digitalis, 408.
Cardialgia :
 antacids, 503.
 charcoal, 529.
Caries :
 cod-liver oil, 345.
Cataract :
 phosphorus, 90.
 atropia, 228.
Catarrh of the bladder :
 juniper, 424.
Cerebral excitement :
 bromide of potassium,
 283.
Chancres :
 iodoform, 341.
 escharotics, 487.
 chloride of zinc, 490.
 solution of nitrate of
 mercury, 491.
 corrosive sublimate, 491.
 nitric acid, 491.

Cholera :
sulphuric acid, 83.
camphor, 176.

Cholera infantum :
sulphuric acid, 83.
chlorate of potassium, 421.

Chordee :
camphor, 176.
arsenic, 319.

Chorea :
oxide of zinc, 39.
Calabar bean, 275.
chloral, 293.
conium, 308.
arsenic, 319.

Cirrhosis :
nitro-muriatic acid, 88.

Colic :
ginger, 74.
assaftetida, 174.
opium, 190.
belladonna, 223.
ether, 246.
antacids, 503.

Colica pictorum :
alum, 31.
sulphuric acid, 83.
belladonna, 223.
chloroform, 252.
salts, 393.

Colitis :
nitrate of silver, 47.
sulphate of magnesium, 393.

Colliquative sweats :
gallic acid, 27.
alum, 31.
sulphuric acid, 83.
belladonna, 224.
ergot, 467.

Condylomata :
nitric acid, 490.
chromic acid, 492.
carbolic acid, 547.

Congestion, hepatic :
nitric acid, 86.
nitro-muriatic acid, 88.

Congestion of cord :
ergot, 468.

Congestion, renal :
juniper, 424.

Conjunctivitis :
alum, 31.
sulphate of copper, 40.
nitrate of silver, 45.

Constipation :
alum, 31.
belladonna, 223.
strychnia, 262.
Calabar bean, 275.
castor oil, 382.
podophyllum, 398.
gambooge, 402.
croton oil, 403.

Convulsions :
anæsthetics, 237.
bromide of potassium, 283.
chloral, 293.

Convulsions :
garlic, 451.

Coryza :
cubebæ, 430.
errhines, 473.
glycerine, 501.

Croup, membranous :
alum, 31.
glycerine, 501.

Croup, pseudo-membranous :
mercury, 326.
lime-water, 514.

Cystitis, chronic :
buchu, 422.
pareira, 422.
uva ursi, 422.
turpentine, 425.
copaiba, 427.
cantharides, 430.

D.

Debility :
prunus Virginiana, 54.

Delirium tremens :
veratrum viride, 139.
hops, 178.
opium, 189.
bromide of potassium, 283.
chloral, 292.
croton oil, 403.

Diabetes :
opium, 190.
glycerine, 501.
digestants, 524.

Diabetes insipidus :
opium, 190.

Diarrhœa :
tannic acid, 26.
catechu, 28.
hæmatoxylin, 28.
alum, 31.
acetate of lead, 36.
bismuth, 38.
aromatics, 72.
oil of cajeput, 76.
sulphuric acid, 83.
nitrous acid, 86.
camphor, 176.
opium, 190.
belladonna, 224.
strychnia, 262.
castor oil, 382.
rhubarb, 387.
ergot, 467.
antacids, 503.
lime-water, 514.
pepsin, 526.
charcoal, 529.

Diarrhœa, chronic :
bismuth, 38.
sulphate of copper, 40.
sulphate of iron, 81.
nitro-muriatic acid, 88.
magnesia, 379.
copaiba, 428.

Dilatation of heart :
digitalis, 123.
carbolic acid, 547.

Diphtheria :
tincture of the chloride of iron, 82.
muriatic acid, 85.
chlorate of potassium, 421.
lime-water, 514.

Diuresis :
turpentine, 425.

Dropsy :
elaterium, 400.
gambooge, 402.
scoparius, 410.
spirit of nitrous ether, 438.
seneka, 447.
blisters, 477.

Dysentery :
acetate of lead, 36.
nitrous acid, 86.
opium, 190.
iodine, 337.
ipecacuanha, 366.
calomel, 386.
chlorate of potassium, 421.
copaiba, 428.
ergot, 467.
flaxseed, 496.
glycerine, 501.

Dysmenorrhœa :
camphor, 176.
belladonna, 223.
acetate of ammonium, 437.
apiol, 456.
guaiaac, 457.

Dyspepsia :
ginger, 74.
piperin, 75.
muriatic acid, 85.
nitric acid, 86.
alcohol, 111.
assaftetida, 174.
strychnia, 262.
Calabar bean, 275.
magnesia, 379.
antacids, 509.

E.

Eczema :
arsenic, 318.
glycerine, 501.

Effusion, pericardial :
squill, 407.

Effusion, pleural :
squill, 407.

Empyema :
iodine, 337.

Endocarditis :
mercury, 326.

Enteritis :
nitrate of silver, 47.
opium, 190.
magnesia, 393.

Enteritis, obstructive :
calomel, 386.
ulmus fulva, 494.
flaxseed, 496.

Epilepsy :

oxide of zinc, 39.
ammoniated copper, 41.
nitrate of silver, 47.
camphor, 176.
anæsthetics, 237.
Calabar bean, 276.
bromide of potassium, 283.
bromide of sodium, 285.
bromide of lithium, 286.
bromide of ammonium, 287.
nitrite of amyl, 302.

Epistaxis :

tannic acid, 25.
ergot, 467.

Erysipelas :

nitrate of silver, 46.
sulphate of iron, 81.
tincture of the chloride of iron, 81.
belladonna, 224.
iodine, 337.

Exophthalmic goitre :

iodine, 334.

F.

Failure of heart :

ammonia, 100, 101.
digitalis, 123.

Faucitis :

nitrate of silver, 45.

Fecal accumulation :

Epsom salts, 393.

Felon :

nitrate of silver, 46.

Fetid expectoration :

carbolic acid, 443.

Fibroid tumors of uterus :

ergotin, 468.

Fissure of the anus :

benzoic acid, 450.

Flooding :

ergot, 466.

Fragilitas ossium :

phosphates, 348.

Furuncles :

phosphorus, 91.

G.

Galactorrhœa :

ergot, 468.

Gangrene :

nitric acid, 86.
carbolic acid, 547.

Gangrene, hospital :

bromine, 492.

Gastralgia :

alum, 31.
prussic acid, 163.
arsenic, 319.

Gastric catarrh :

chloride of ammonium, 446.

Gastric pain :

bismuth, 38.

Gastric ulcer :

nitrate of silver, 46.
oil of turpentine, 115.

Gastritis :

nitrate of silver, 46.

Glands, enlarged :

conium, 308.
mercury, 329.
iodine, 337.

Gleet :

tincture of the chloride of iron, 81.
turpentine, 425.
cantharides, 430.

Gonorrhœa :

bismuth, 38.
nitrate of silver, 46.
tartar emetic, 133.
pareira, 422.
copaiba, 427.
cubeb, 429.
matico, 430.

Gout :

ether, 246.
arsenic, 320.
iodide of potassium, 339.
colechicum, 354.
rubefacients, 482.

Graves's disease :

iodine, 336.

Gums, retraction of :

iodine, 337.

H.

Hæmatemesis :

tannic acid, 26.
gallic acid, 27.
sulphate of iron, 81.
oil of turpentine, 116.

Hæmaturia :

gallic acid, 27.
turpentine, 426.

Hæmoptysis :

alum, 31.
acetate of lead, 37.
sulphate of iron, 81.
opium, 190.
astringents, 442.
ergot, 467.

Headache :

magnesia, 379.
ergot, 468.
antacids, 510.

Hemiplegia :

strychnia, 261.

Hemorrhage :

astringents, 24.

Hemorrhage from bowels :

tannic acid, 26.
oil of turpentine, 116.
erigeron, 425.

Hemorrhoids :

tannic acid, 25.
stramonium, 232.
iodoform, 341.
sulphur, 380.
aloes, 390.
cubeb, 429.

Hepatic abscess :

chloride of ammonium, 446.
escharotics, 487.

Hepatitis :

nitro-muriatic acid, 88.
mercury, 326.
chloride of ammonium, 446.

Hiccough :

musk, 172.
oil of amber, 177.
belladonna, 223.
ether, 246.

Hydrocephalus :

iodide of potassium, 339.

Hydrophobia :

escharotics, 487.

Hypertrophy of heart :

digitalis, 123.
aconite, 154.

Hypertrophy of uterus :

ergot, 468.

Hypochondriasis :

alcohol, 111.

Hysteria :

antispasmodics, 171.
musk, 172.
valerian, 173.
valerianic acid, 173.
assaftida, 174.
camphor, 176.
anæsthetics, 237.
ether, 246.
oil of wormseed, 518.

I.

Impotence :

turpentine, 426.

Incontinence of urine :

belladonna, 223.
strychnia, 262.
chloral, 293.
turpentine, 426.

Inflammations :

tartar emetic, 133.
mercury, 326.

Intermittent fever :

quinia, 69.
piperin, 75.
arsenic, 316, 317.
chloride of ammonium, 446.

Iritis :

atropia, 228.
mercury, 326.

Irritable heart :

digitalis, 124.

Irritation of bladder :

buchu, 422.
pareira, 422.

Ich :

sulphur, 380.

J.

Jaundice :

lemon-juice, 168.
potassium salts, 416.

Jaundice, catarrhal :
 ipecacuanha, 366.
 calomel, 385.
Joints, enlarged :
 conium, 308.
 mercury, 329.
 iodine, 337.
 cod-liver oil, 345.
Joints, chronic inflammation of :
 blisters, 477.

K.

Keratitis :
 atropia, 228.

L.

Laryngismus stridulus :
 belladonna, 223.
Laryngitis :
 nitrate of silver, 45.
 mercury, 326.
 narcotics, 443.
 alteratives, 443.
 glycerine, 501.
Lead-paralysis :
 strychnia, 261.
Leucorrhœa :
 tannic acid, 26.
 ammonio-ferrie alum, 82.
 iodine, 337.
 permanganate of potassium, 533.
Lichen :
 arsenic, 318.
Locomotor ataxia :
 nitrate of silver, 47.
Lumbago :
 iodide of potassium, 338.
 cod-liver oil, 346.
 sulphur, 380.
Lupus :
 phosphorus, 90.
 arsenic, 322.

M.

Malarial neuralgia :
 quinia, 69.
Malarial poisoning :
 quinia, 67.
 cinchonia, 70.
 arsenic, 316.
 potassium salts, 416.
 apiol, 456.
Malignant pustule :
 escharotics, 487.
Mania :
 croton oil, 403.
 blisters, 477.
Mania a potu :
 veratrum viride, 139.
 valerian, 173.
Melancholia :
 alcohol, 111.

Menorrhagia :
 aloes, 390.
 erigeron, 425.
 acetate of ammonium, 437.
 rue, 455.
 savine, 455.
 ergot, 467.
Mercurial sore mouth :
 tannic acid, 25.
 opium, 190.
 belladonna, 224.
Metallic poisoning :
 iodide of potassium, 339.
Metritis :
 ergot, 468.
Mitral disease :
 digitalis, 124.
Muscular rheumatism :
 eupatorium, 53.
 aconite, 153.
 opium, 191.

Myelitis :
 nitrate of silver, 47.

N.

Narcotic poisoning :
 mustard, 369.
Nephritis :
 belladonna, 224.
Nervous cough :
 belladonna, 223.
Nervous exhaustion :
 phosphorus, 90.
Nervous irritability :
 prussic acid, 163.
Neuralgia :
 subcarbonate of iron, 80.
 phosphorus, 90.
 alcohol, 111.
 veratria, 147.
 aconite, 154.
 valerianic acid, 173.
 cannabis Indica, 204.
 belladonna, 223.
 ether, 246.
 chloroform, 252.
 bromide of potassium, 283.
 croton chloral, 297.
 arsenic, 319.
 iodide of potassium, 338.
 cod-liver oil, 346.
 chloride of ammonium, 447.
 ergot, 468.
 epispastics, 476.
Neuralgia, intermittent :
 quinia, 246.
 arsenic, 317.
 apiol, 456.
Night-pains, syphilitic :
 iodoform, 340.
Nymphomania :
 bromide of potassium, 285.

O.

Onychia maligna :
 corrosive sublimate, 491.
Ophthalmia :
 iodine, 337.
Osmidrosis :
 tannic acid, 26.
Osteomalacia :
 phosphorus, 91.
 phosphates, 348.
Otorrhœa :
 permanganate of potassium, 533.
Over-secretion :
 astringents, 24.
Ozœna :
 iodine, 337.
 permanganate of potassium, 533.

P.

Paralysis agitans :
 conium, 308.
Paraplegia, myelitic :
 nitrate of silver, 47.
 phosphorus, 90.
Parasitic skin-diseases :
 iodine, 337.
Pemphigus :
 arsenic, 318.
Pericarditis :
 iodide of potassium, 339.
Periostitis, syphilitic :
 phosphates, 349.
Peritonitis :
 veratrum viride, 139.
 aconite, 153.
 opium, 190.
 mercury, 326.
 blisters, 477.
 poultices, 503.
Pernicious fever :
 quinia, 69.
 mustard, 369.
 epispastics, 476.
Phthisis :
 prunus Virginiana, 54.
 alcohol, 111.
 cannabis Indica, 204.
 iodine, 337.
 cod-liver oil, 345.
 phosphates, 349.
Pityriasis :
 oil of cajeput, 76.
Pleuritis :
 mercury, 326.
 iodine, 337, 339.
 blisters, 477.
 poultices, 503.
Pneumonia :
 alcohol, 110.
 oil of turpentine, 115.
 tartar emetic, 133.
 veratrum viride, 139.
 aconite, 153.
 musk, 172.
 belladonna, 224.
 mercury, 326.

Pneumonia :
 blisters, 477.
 poultices, 503.
Post-partum hemorrhage :
 ergot, 466.
Pregnancy :
 phosphates, 349.
Priapism :
 hops, 178.
Prolapsus of the rectum :
 strychnia, 262.
Pruritus :
 glycerine, 501.
Psoriasis :
 oil of cajeput, 76.
 phosphorus, 90.
 arsenic, 318.
 iodine, 334.
 glycerine, 501.
Pyralism :
 opium, 190.
 belladonna, 224.
Puerperal eclampsia :
 chloroform, 251.
 chloral, 292.
 nitrite of amyl, 303.
Puerperal fever :
 quinia, 66.
 oil of turpentine, 116.
Puerperal peritonitis :
 mercury, 327.
Purpura hemorrhagica :
 oil of turpentine, 116.
Pyæmia :
 quinia, 66.
 tincture of the chloride
 of iron, 82.
 carbolic acid, 546.
Pyelitis, chronic :
 buchu, 422.
 uva ursi, 423.
 juniper, 424.
 turpentine, 425.
 copaiba, 427.
 cantharides, 430.

R.

Rachitis :
 phosphates, 348.
Relaxation :
 astringents, 23.
Relaxation of uvula :
 pellitory, 472.
Remittent fever :
 quinia, 69.
 arsenic, 316.
Retention of urine :
 strychnia, 262.
Rheumatism :
 quinia, 67.
 oil of cajeput, 76.
 veratria, 147.
 arsenic, 319.
 iodine, 337.
 iodide of potassium, 338.
 iodoform, 340.
 cod-liver oil, 346.
 colchicum, 354.
 guaiac, 358.

Rheumatism :
 sulphur, 380.
 alcohol, 439.
 Burgundy pitch, 485.
Rheumatism, cerebral :
 nitrite of amyl, 303.
Rheumatism, inflammatory :
 quinia, 67.
 lemon-juice, 168.
 cimicifuga, 180.
 bromide of ammonium,
 287.
 potassium salts, 415.
 nitrate of potassium,
 419.
 Dover's powder, 436.
Rheumatoid arthritis :
 arsenic, 320.
 iodide of potassium, 339.
Rickets :
 phosphorus, 91.

S.

Scarlet fever :
 aconite, 153.
 belladonna, 224, 229.
 chlorate of potassium,
 421.
Sciatica :
 iodide of potassium, 338.
 cod-liver oil, 346.
 sulphur, 380.
Serofulosis :
 alcohol, 111.
 sarsaparilla, 356.
 pipsissewa, 423.
 iodine, 334.
 cod-liver oil, 344.
 phosphoric acid, 347.
 phosphates, 348.
 syrup of iodide of iron,
 560.
Serofulous tumors :
 emplastrum ammoniaci,
 448.
Scurvy :
 lemon-juice, 168.
 chlorate of potassium,
 421.
Seat-worms :
 quassia, 51.
 oil of cajeput, 76.
 vinegar, 169.
Shock :
 epispastics, 476.
Sick stomach :
 ipecacuanha, 366.
Singultus :
 chloral, 293.
Skin, tubercular affections of :
 arsenic, 322.
 Donovan's solution, 340.
 magnesia, 379.
Sleeplessness :
 opium, 190.
 chloral, 292.
Smallpox :
 opium, 190.
 carbolic acid, 546.

Softening of the brain :
 phosphorus, 90.
Sore nipples :
 tannic acid, 26.
Sore throat :
 tannic acid, 25.
 alum, 31.
 nitrate of silver, 45.
 belladonna, 224.
 chlorate of potassium,
 421.
Spasm of œsophagus :
 anæsthetics, 237.
Spermatorrhœa :
 turpentine, 426.
Spongy gums :
 tannic acid, 25.
Sprains :
 vinegar, 169.
Stomatitis :
 chlorate of potassium,
 421.
 carbolic acid, 547.
Strangury :
 opium, 191.
Subinvolution of uterus :
 ergot, 468.
Sunburn :
 vinegar, 169.
Suppression of urine :
 digitalis, 408.
Syncope :
 digitalis, 125.
 ether, 146.
Syphilis :
 nitric acid, 86.
 nitro-muriatic acid, 88.
 mercury, 327.
 iodide of potassium, 339.
 cod-liver oil, 346.
 sarsaparilla, 356.
 guaiac, 358.
 chlorate of potassium,
 421.
Syphilitic nodes :
 emplastrum ammoniaci
 cum hydrargyro, 448.

T.

Tabes mesenterica :
 cod-liver oil, 345.
Tape-worm :
 ether, 246.
 spigelia, 516.
 chenopodium, 517.
 brayera, 518.
 santonin, 519.
 pepo, 522.
 rottlera, 523.
 mucuna, 523.
Tetanus :
 cannabis Indica, 204.
 Calabar bean, 275.
 bromide of potassium,
 284.
 chloral, 292.
 nitrite of amyl, 303.
Tinea capitis :
 lime-water, 514.

Tonsillitis :
tincture of capsicum, 76.
Toothache :
oil of cloves, 73.
pellitory, 472.
Trismus nascentium :
Calabar bean, 275.
chloral, 293.
Tuberculosis :
cod-liver oil, 345.
Tympanitis :
assafetida, 174.
Typhoid fever :
quinia, 67.
alcohol, 110.
oil of turpentine, 115.
digitalis, 126.
veratrum viride, 138.
Typhus fever :
alcohol, 110.
belladonna, 224.
chloral, 292.
chlorine water, 534.
carbolic acid, 546.

U.

Ulcer of cervix uteri :
solution of nitrate of
mercury, 491.
Ulcer of cornea :
atropia, 228.

Ulceration, chronic :
iodoform, 341.
Ulceration of bowels :
oil of turpentine, 115.
Ulcers :
alum, 31.
nitrate of silver, 46.
sulphuric acid, 83.
nitric acid, 86.
conium, 308.
lime-water, 514.
permanganate of potas-
sium, 532.
Ununited fracture :
phosphates, 349.
Uræmia :
elaterium, 400.
Urethral fever :
aconite, 153.
bromide of potassium,
284.
Urethritis :
nitrate of silver, 45.
Uric acid calculus :
potassium salts, 416.
Uric acid diathesis :
benzoic acid, 450.
sodium salts, 513.
Uric acid gravel :
potassium salts, 416.
Uterine inertia :
ergot, 465.

Uterus, cancer of :
iodoform, 341.

V.

Venereal warts :
nitric acid, 86.
Vesical catarrh :
benzoic acid, 450.
Vomiting :
bismuth, 38.
aconite, 154.
prussic acid, 163.
opium, 191.
bromide of potassium,
283.
lime-water, 511.
carbolic acid, 547.

W.

Whooping-cough :
assafetida, 174.
camphor, 176.
belladonna, 223.
bromide of ammonium,
287.
chloral, 294.
conium, 308.
arsenic, 319.





R
REYNOLDS
BINDERY, INC.
SEPT 1973

QV W875t

61841020R



NLM 05063383 0

NATIONAL LIBRARY OF MEDICINE